

**Table 2-1. Composition of Red Dog lead and zinc concentrates**

Element/Compound	Concentrate	
	Lead	Zinc
Aluminum oxide <sup>a</sup>	2,600	1,300
Antimony <sup>a</sup>	1,600	400
Arsenic <sup>a</sup>	400	200
Barium <sup>a</sup>	2,400	2,700
Cadmium <sup>a</sup>	1,200	3,300
Calcium oxide <sup>a</sup>	600	500
Copper <sup>a</sup>	600	1,400
Iron <sup>a</sup>	53,000	50,000
Lead <sup>a</sup>	579,000	32,000
Manganese oxide <sup>a</sup>	100	100
Silicon oxide <sup>a</sup>	38,000	35,000
Sulfate <sup>a</sup>	4,000	4,500
Sulfur (total) <sup>a</sup>	205,000	317,000
Zinc <sup>a</sup>	108,000	552,000
Bismuth	6	8
Chloride	50	50
Chromium	677	537
Cobalt	77	98
Fluoride	64	56
Gallium	11	26
Germanium	17	79
Gold	0.22	0.209
Manganese	12	10
Mercury	18	94
Molybdenum	43	20
Nickel	45	16
Selenium	28	3
Silver	420	137
Strontium	11	10
Thallium	70	19
Tin	23	49
Vanadium	23	13

**Source:** Teck Cominco (2002)

**Note:** All units are expressed in ppm (i.e., mg/kg dry weight basis).

<sup>a</sup> Values are based on dry weight; actual lead and zinc concentrates contain approximately 8.5 and 9.5 percent water, respectively.

Table 2-2. DMTS-related spills from DEC database

Spill Number	Spill Date	Spill Name	Quantity Released	Quantity Unit ID	Substance Type	Clean Up
93389921001	07/29/93	36,000 gal diesel (Tank #2)	36,000	gallons	Diesel	yes
98389915802	06/07/98		70	gallons	Diesel	yes
98389933101	11/27/98		55	gallons	Diesel	yes
97389925701	09/14/97		30	gallons	Diesel	yes
98389915702	06/06/98		20	gallons	Diesel	yes
98389908201	03/23/98		15	gallons	Diesel	yes
99389935401	12/20/99		10	gallons	Diesel	yes
00389907901	03/19/00		10	gallons	Diesel	yes
00389923201	08/19/00		10	gallons	Diesel	yes
96389929601	10/22/96		15	gallons	Engine lube oil	yes
98389921901	08/07/98		10	gallons	Engine lube oil	yes
01389919601	07/15/01		90	gallons	Hydraulic oil	yes
98389921601	08/04/98		45	gallons	Hydraulic oil	yes
01389903101	01/31/01		30	gallons	Hydraulic oil	yes
97389917301	06/22/97		25	gallons	Hydraulic oil	yes
00389915601	06/04/00		25	gallons	Hydraulic oil	yes
97389922401	08/12/97	Port old CSB-column 47	20	gallons	Hydraulic oil	yes
99389933702	12/03/99		20	gallons	Hydraulic oil	yes
95389934901	12/15/95		20	gallons	Hydraulic oil	no
97389931801	11/14/97		15	gallons	Hydraulic oil	yes
96389933401	11/29/96		10	gallons	Hydraulic oil	yes
98389915203	06/01/98		10	gallons	Hydraulic oil	yes
97389927803	10/05/97		65	gallons	Other <sup>a</sup>	yes
98389921301	08/01/98	Port CSB	76,000	pounds	Lead concentrate	yes
99389902101	01/21/99	Port Road mile 9	60,000	pounds	Lead concentrate	yes
00389928301	10/09/00	Port Road mile marker 31	52,000	pounds	Lead concentrate	yes
99389910101	04/11/99	Port CSB	150	pounds	Lead concentrate	yes
98389931402	11/10/98		200	pounds	Other <sup>b</sup>	yes
98389903801	02/07/98	Zinc at mile marker 27.25	140,000	pounds	Zinc concentrate	yes
00389936301	12/28/00	Port Road mile marker 45	88,100	pounds	Zinc concentrate	yes
97389923301	08/21/97	Port site entrance to racetrack	70,000	pounds	Zinc concentrate	yes
98389932501	11/21/98	Port Road mile 41.75 by MS 11	70,000	pounds	Zinc concentrate	yes
99389900601	01/06/99	Port Road mile 45	50,000	pounds	Zinc concentrate	yes
97389900201	01/02/97	40,000 lb zinc at mile marker 27	40,000	pounds	Zinc concentrate	yes
98389901701	01/17/98	Port Road mile 35 (near MS9)	37,760	pounds	Zinc concentrate	yes
98389919301	07/12/98	Mile post 42	26,500	pounds	Zinc concentrate	yes
96389915901	06/07/96	6,743 lb zinc conc. at MS-2	2,000	pounds	Zinc concentrate	yes
98389910701	04/17/98	Port Road 150 ft south of Tutak Bridge	800	pounds	Zinc concentrate	yes
00389923402	08/21/00	Port - Conveyor P-10 drive house	750	pounds	Zinc concentrate	yes
01389920101	07/20/01	Red Dog Mine zinc spill MP 38.3	20,000	pounds	Zinc concentrate	yes

**Note:** CSB - concentrate storage building  
DEC - Alaska Department of Environmental Conservation

Data were provided by DEC from its Prevention and Early Response & Preparation database.

Table includes spills greater than or equal to 10 gallons or 10 pounds.

Database does not include spills that occurred prior to 1995, except for July 29, 1993, spill.

<sup>a</sup> Uncertain, but possible match with process water spill at the mine mill on October 5, 1997.

<sup>b</sup> Uncertain, but possible match with 1 gallon spill of ethylene glycol at the mine (Hagy 2003, pers. comm.)

Table 2-3. Relative importance of potential human exposure pathways<sup>a</sup>

Metal	Human Exposure Pathways			Cumulative PRG
	Inhalation	Dermal	Ingestion	
Aluminum	2,882,040		78,214	76,142
Antimony			31	31
Arsenic	588	4	0.4	0.4
Barium	294,086		5,475	5,375
Cadmium (cancer)	1,405			1,404
Cadmium (noncancer)		698	39	37
Chromium VI (cancer)	30			30
Chromium VI (noncancer)	4,529		235	223
Cobalt (cancer)	903			903
Cobalt (noncancer)	11,734		1,564	1,380
Copper			3,129	3,129
Fluoride		16,760	4,693	3,666
Iron			23,464	23,463
Lead				
Manganese	28,820		1,877	1,762
Mercury			23	23
Molybdenum			391	391
Nickel			1,564	1,564
Selenium			391	391
Silver			391	391
Strontium			46,929	46,924
Thallium			5	5
Tin			46,929	46,924
Vanadium			548	547
Zinc			23,464	23,463

**Note:** Units are in mg/kg.

EPA - U.S. Environmental Protection Agency

PRG - preliminary remediation goal

<sup>a</sup> The screening values listed above are U.S. EPA (2003c) Region 9 PRGs for residential soil. This table is not meant to provide screening concentrations for the DMTS risk assessment. Rather, the PRGs listed above are provided to illustrate the relative contribution of inhalation, dermal contact, and ingestion exposure. The PRGs were derived assuming a risk level of  $1 \times 10^{-6}$  for cancer and a hazard quotient of 1.0 for noncancer endpoints. Higher PRGs indicate relatively lower contribution to risk, and vice versa. These PRGs suggest that dermal contact is at least an order of magnitude lower risk than ingestion, and that inhalation is several orders of magnitude lower risk than ingestion.

Table 2-4. Subsistence resource categories and representative receptors

Subsistence Resource <sup>a</sup>	Inupiat Name <sup>b</sup>	Scientific Name	Representative Receptors
<b>Fish</b>			<b>Fish</b>
Bluecod			<b>Note:</b> Risk to fish evaluated by comparison of chemical concentrations in sediment and water to effects ranges reported in the literature
Bullhead	kanayuq	<i>Myoxocephalus quadricornis</i>	
Burbot/mudshark	tittaaliq	<i>Lota lota</i>	
Char (arctic)	aqalukpik	<i>Salvelinus alpinus</i>	
Dolly Varden		<i>Salvelinus malma</i>	
Grayling (arctic)	suluppaugaq	<i>Thymallus arcticus</i>	
Herring (Pacific)	uqsruqtuuq	<i>Clupea pallasii</i>	
Pike (northern)	siilik	<i>Esox lucius</i>	
Salmon (king)		<i>Oncorhynchus tshawytscha</i>	
Salmon (pink)	qalugruaq	<i>Onchorynchus gorbuscha</i>	
Salmon (silver)		<i>Oncorhynchus kisutch</i>	
Sheefish	sii	<i>Stenodus leucichthys</i>	
Smelt (rainbow)	ijhuabniq	<i>Osmerus mordax dentex</i>	
Tomcod (arctic cod)	uugaq	<i>Boreogadus saida</i>	
Whitefish	qalupiat, quptik, qaalbiq, qalusraaq	<i>Coregonis</i> spp., <i>Prosopium</i> spp.	
<b>Shellfish</b>			<b>Shellfish</b>
Crab	putyuun	<i>Paralithodes</i> spp., <i>Lithodes aequispinus</i> , <i>Cancer magister</i> , <i>Chionoecetes</i> spp.	<b>Note:</b> Risk to shellfish evaluated by comparison of chemical concentrations in sediment and water to effects ranges reported in the literature
Shrimp	putuguqsiuyuk	<i>Pandalus</i> spp., <i>Pandalopsis</i> spp.	
<b>Sea Mammals</b>			<b>Sea Mammals</b>
Seal (bearded)	ugruk	<i>Erignathus barbatus</i>	Bearded seal (invertebrate-eater)
Seal (ribbon)	qaigullik	<i>Phoca fasciata</i>	Ringed seal (fish- and invertebrate-eater)
Seal (spotted)	qasigiaq	<i>Phoca largha</i>	
Walrus	aiviq	<i>Odobenus rosemarus</i>	
Whale (Beluga)	sisuaq	<i>Delphinapterus leucas</i>	
Whale (bowhead)	abviq	<i>Balaena mysticetus</i>	
<b>Large Mammals</b>			<b>Large Mammals</b>
Bear (black)	iyyabriq	<i>Ursus americanus</i>	Caribou (plant-eater)
Bear (brown/grizzly)	akjaq	<i>Ursus horribilis</i>	Moose (plant-eater)
Caribou	tuttu	<i>Rangifer tarandus</i>	Polar bear (marine animal-eater)
Moose	tiniikaq	<i>Alces alces</i>	
Muskox	imummak	<i>Ovibus moschatus</i>	
Sheep (dall)	ipnaiq	<i>Ovis dalli dalli</i>	
<b>Small Mammals</b>			<b>Small Mammals</b>
Beaver	aqu	<i>Castor canadensis</i>	Arctic fox (terrestrial animal-eater)
Fox (arctic)	qujhaaq	<i>Alopex lagopus</i>	Muskrat (freshwater plant-eater)
Fox (red)	kavviaq	<i>Vulpes fulva</i>	River otter (freshwater fish-eater)
Muskrat	kigvaluk	<i>Ondatra zibethicus</i>	Tundra shrew (terrestrial invertebrate-eater)
Otter (river)	pamiuqtuuq	<i>Lutra canadensis</i>	Tundra vole (terrestrial plant-eater)
Porcupine	ixuqutaq	<i>Erethizon dorsatum</i>	
Rabbit (Alaska hare)	ukallisugruk	<i>Lepus othus</i>	
Rabbit (snowshoe hare)	ukalliuraq	<i>Lepus americanus</i>	
Squirrel (ground)	siksrik	<i>Citellus parryi</i>	
Squirrel (red/tree)	saqalataayiq	<i>Tamiasciurus hudsonicus</i>	
Wolverine	qapvik	<i>Gulo luscus</i>	
<b>Birds</b>			<b>Birds</b>
Crane (sandhill)	tatirgaq	<i>Grus canadensis</i>	Black-bellied plover (marine insect-eater)
Ducks	qaugak	Multiple species	Brant (marine plant-eater)
Geese		Multiple species	Common snipe (freshwater invertebrate-eater)
Grouse (spruce)		<i>Falci pennis canadensis</i>	Green-winged teal (freshwater plant-eater)
Owl (snowy)	ukpik	<i>Nyctea scandiaca</i>	Lapland longspur (terrestrial insect-eater)
Ptarmigan (mountain)			Red-throated loon (freshwater and marine fish-eater)
Ptarmigan (willow)	aqargiq	<i>Lagopus lagopus</i>	Snowy owl (terrestrial animal-eater)
Swans (tundra, formerly whistling)	qugruk	<i>Cygnus columbianus</i>	Willow ptarmigan (terrestrial plant-eater)

Table 2-4. (cont.)

Subsistence Resource <sup>a</sup>	Inupiat Name <sup>b</sup>	Scientific Name	Representative Receptors
<b>Vegetation</b>			<b>Vegetation</b>
Blueberry	asriavik	<i>Vaccinium uliginosum</i>	<b>Note:</b> Risk to plants evaluated by comparison of chemical concentrations in soil and plants to effects ranges reported in the literature
Coast greens sura (fresh willow leaf)		<i>Salix</i> spp.	
Cow parsnip		<i>Heracleum lanatum (Umbelliferae)</i>	
Cranberry (bog)	qunmun asriaq	<i>Vaccinium oxycoccus</i>	
Cranberry (highbush)	uqpifeaq	<i>Viburnum edule</i>	
Cranberry (lowbush)	kikmieeq	<i>Vaccinium vitis idaea</i>	
Crowberry (black berry)	paunbaq	<i>Empetrum nigrum</i>	
Eskimo (labrador/tundra) tea		<i>Ledum decubens</i>	
Eskimo/wild potato	masru	<i>Hedysarum alpinum</i>	
Herbal tea		Species unknown	
Matsu sura		Species unknown	
Raspberry	tuunbaum asriaq	<i>Rubus pendantus</i>	
Salmonberry/cloudberry	aqpik	<i>Rubus chamaemorus</i>	
Sourdock	quabaq	<i>Rumex arcticus</i>	
Spring tea		Species unknown	
Stinkweed		<i>Thlaspi arvense</i>	
Wild celery	ikuusuk	<i>Angelica lucida</i>	
Wild onions/chives	paatitaaq	<i>Allium schoenoprasum sibiricum</i>	
Wild rhubarb	qusrimmak	<i>Polygonum alaskanum</i>	
Wild tea		Species unknown	

<sup>a</sup> Sundet (2002a,b, pers. comm.).

<sup>b</sup> Webster and Zibell (2003).

Table 2-5. Summary of preliminary assessment endpoints, representative receptors, and measurement endpoints<sup>a</sup>

Environment	Assessment Endpoint	Representative Receptor	Measurement Endpoint
Terrestrial	Structure and function of <b>terrestrial plant communities</b>	Terrestrial plant communities	Range of CoPC concentrations in soil relative to ecological screening benchmarks
Terrestrial	Structure and function of <b>terrestrial fauna communities</b>	Soil fauna communities	Range of CoPC concentrations in soil relative to ecological screening benchmarks
Terrestrial	Survival, growth, and reproduction of <b>terrestrial avian herbivore populations</b>	Willow ptarmigan	Range of modeled total dietary exposures relative to avian TRVs
Terrestrial	Survival, growth, and reproduction of <b>terrestrial mammalian herbivore populations</b>	Tundra vole; caribou; moose	Range of modeled total dietary exposures relative to mammalian TRVs
Terrestrial	Survival, growth, and reproduction of <b>terrestrial avian invertevore populations</b>	Lapland longspur	Range of modeled total dietary exposures relative to avian TRVs
Terrestrial	Survival, growth, and reproduction of <b>terrestrial mammalian invertevore populations</b>	Tundra shrew	Range of modeled total dietary exposures relative to mammalian TRVs
Terrestrial	Survival, growth, and reproduction of <b>terrestrial avian carnivore populations</b>	Snowy owl	Range of modeled total dietary exposures relative to avian TRVs
Terrestrial	Survival, growth, and reproduction of <b>terrestrial mammalian carnivore populations</b>	Arctic fox	Range of modeled total dietary exposures relative to mammalian TRVs
Freshwater Aquatic	Structure and function of <b>freshwater aquatic and wetland plant communities</b>	Freshwater aquatic and wetland plant communities	Range of CoPC concentrations in freshwater sediment and water relative to ecological screening benchmarks
Freshwater Aquatic	Structure and function of <b>freshwater aquatic invertebrate communities</b>	Freshwater aquatic invertebrate communities	Range of CoPC concentrations in freshwater sediment relative to ecological screening benchmarks
Freshwater Aquatic	Structure and function of <b>freshwater fish communities</b>	Freshwater fish communities	Range of CoPC concentrations in freshwater relative to ecological screening benchmarks
Freshwater Aquatic	Survival, growth, and reproduction of <b>freshwater avian herbivore populations</b>	Green-winged teal	Range of modeled total dietary exposures relative to avian TRVs
Freshwater Aquatic	Survival, growth, and reproduction of <b>freshwater mammalian herbivore populations</b>	Muskrat	Range of modeled total dietary exposures relative to mammalian TRVs
Freshwater Aquatic	Survival, growth, and reproduction of <b>freshwater avian invertevore populations</b>	Common snipe	Range of modeled total dietary exposures relative to avian TRVs

Table 2-5. (cont.)

Environment	Assessment Endpoint	Representative Receptor	Measurement Endpoint
Freshwater Aquatic	Survival, growth, and reproduction of <b>freshwater avian piscivore populations</b>	Red-throated loon	Range of modeled total dietary exposures relative to avian TRVs
Freshwater Aquatic	Survival, growth, and reproduction of <b>freshwater mammalian piscivore populations</b>	River otter	Range of modeled total dietary exposures relative to mammalian TRVs
Marine	Structure and function of <b>marine aquatic and wetland communities</b>	Marine aquatic and wetland plant communities	Range of CoPC concentrations in marine sediment and water relative to ecological screening benchmarks
Marine	Structure and function of <b>marine aquatic invertebrate communities</b>	Marine aquatic invertebrate communities	Range of CoPC concentrations in marine sediment and water relative to ecological screening benchmarks
Marine	Structure and function of <b>marine fish communities</b>	Marine fish communities	Range of CoPC concentrations in marine water relative to ecological screening benchmarks
Marine	Survival, growth, and reproduction of <b>marine avian herbivore populations</b>	Brant	Range of modeled total dietary exposures relative to avian TRVs
Marine	Survival, growth, and reproduction of <b>marine avian invertevore populations</b>	Black-bellied plover	Range of modeled total dietary exposures relative to avian TRVs
Marine	Survival, growth, and reproduction of <b>marine mammalian invertevore populations</b>	Bearded seal	Range of modeled total dietary exposures relative to mammalian TRVs
Marine	Survival, growth, and reproduction of <b>marine avian piscivore populations</b>	Red-throated loon	Range of modeled total dietary exposures relative to avian TRVs
Marine	Survival, growth, and reproduction of <b>marine mammalian piscivore populations</b>	Ringed seal	Range of modeled total dietary exposures relative to mammalian TRVs
Marine	Survival, growth, and reproduction of <b>marine mammalian carnivore populations</b>	Polar bear	Range of modeled total dietary exposures relative to mammalian TRVs

**Note:** CoPC - chemical of potential concern  
 TRV - toxicity reference value

<sup>a</sup> A refined version of this table was developed following CoPC screening and is presented later in this document.

**Table 3-1. Target chemical list**

---

---

Aluminum
Antimony
Arsenic
Barium
Cadmium
Chromium
Cobalt
Copper
Fluoride
Iron
Lead
Manganese
Mercury
Molybdenum
Nickel
Selenium
Silver
Strontium
Thallium
Tin
Vanadium
Zinc

---

---



Table 3-2. Overview of prior studies

Lead Organization	Study Type	Citation	Study Dates	Analytical Data Available					
				Moss	Soil	Water	Sediment	Plants	Fish
<b>Pre-Mine/Baseline</b>									
Teck Cominco	Environmental baseline study	Dames & Moore (1983a,b)	1981–1983			•	•		•
General Crude Oil and Minerals	Environmental baseline study	Ward and Olson (1980)	1978–1979			•			•
Alaska Department of Environmental Conservation	Aquatic baseline study	EVS and Ott Water (1983)	1982						•
U.S. Fish and Wildlife Service	Baseline study for Selawik NWR	Mueller et al. (1993)	1987–1988			•	•		•
<b>Post-Mine</b>									
Teck Cominco	Port site monitoring	ENSR (1990, 1991, 1993, 1996); RWJ (1997)	1990–1996			•	•	•	
	Transportation corridor monitoring	ENSR (1991)	1991–1992			•	•		
	Vegetation and soil monitoring	RWJ (1998)	1992, 1993, 1997			•			
	Fugitive dust study	Exponent (2002a); DEC et al. (2002)	2001	•	•	•			•
	Kivalina drinking water study	RWJ (1997); DHSS (2001); (Kulas 2003, pers. comm.)	1991–2003			•			
	Supplemental road sampling	Exponent (2002b)	2002			•			
	Caribou evaluation	Exponent (2002c)	1996, 2002						•
	Port site characterization	Exponent (2003a)	2002			•	•	•	
	Phase I risk assessment field sampling program	Exponent (2003e) and Appendix A of this document	2003			•	•	•	
Alaska Industrial Development and Export Authority	Sediment quality survey	Cominco et al. (1999)	1998			•		•	
Alaska Department of Environmental Conservation	Subsistence foods investigation	E&E (2002); DHSS (2001)	2001			•			•
Alaska Department of Fish and Game	NPDES monitoring, expanded scope	Weber-Scannell and Ott (2001)	1994–2001			•			•
	Juvenile fish tissue study	Morris and Ott (2001); DHSS (2001)	1993, 1998–2001						•
National Park Service	DMTS road dustfall study	Ford and Hasselbach (2001)	2000	•	•				
		Hasselbach (2003a, pers. comm.)	2003	•	•				
Kivalina Village	Kivalina drinking water sampling	DHSS (2001)	1995, 1996, 2001			•			
United States Geological Survey	Cape Krusenstern trace elements study	Brabets (2003, pers. comm.)	2002			•	•		
	Willow study	Gough (2003, pers. comm.)	2002						•
	Soil study	Kelley and Hudson (2003)	2002			•			

Table 3-3. Analytical data summary for screening chemicals of potential concern

Environment	Medium	Site/ Reference	Survey Name	Numbers of Samples by Analyte <sup>a</sup>																					
				Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Copper	Fluoride	Iron	Lead	Manganese	Mercury	Molybdenum	Nickel	Selenium	Silver	Strontium	Thallium	Tin	Vanadium	Zinc
Terrestrial	Soil	Site	PHASE1RA, PSCHAR, FUGDST01, SUPPRSS, TECK03	51	40	75	40	478	40	40	40	12	51	479	40	12	40	40	30	40	20	12	27	40	479
		Reference	PHASE1RA, FUGDST01	10	5	10	5	10	5	5	5	5	10	10	5	5	5	5	5	5	5	5	5	5	10
	Tundra Soil	Site	PHASE1RA, PSCHAR, ENSR92	31	25	31	25	224	25	25	25	12	31	264	25	12	25	25	25	25	17	12	17	25	264
		Reference	PHASE1RA	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10
Stream	Sediment	Site	PHASE1RA	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	
		Reference	PHASE1RA	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
	Water	Site	TECK03, TECK01, USGS02	230	14	14	14	229	18	14	18	31	230	230	18	14	14	14	29	14	14	29	14	14	230
		Reference	PHASE1RA	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
Tundra Pond	Sediment	Site	PHASE1RA	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	
		Reference	PHASE1RA	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
	Water	Site	PHASE1RA	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	
		Reference	PHASE1RA	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	
Lagoon	Sediment	Site	PHASE1RA, PSCHAR	8	8	8	8	34	8	8	8	11	8	26	8	8	8	8	8	8	8	8	11	8	26
		Reference	PHASE1RA, PSCHAR, ENSR91, ENSR92, ENSR95, ENSR96	3	3	3	3	13	3	3	3	3	3	28	3	3	3	3	3	3	3	3	3	3	28
	Water	Site	PHASE1RA, PSCHAR	8	8	8	8	14	8	8	8	8	8	14	8	8	8	8	8	8	8	8	8	8	14
		Reference	PHASE1RA, PSCHAR	3	3	3	3	5	3	3	3	3	3	5	3	3	3	3	3	3	3	3	3	3	5
Marine	Sediment	Site	PHASE1RA, PSCHAR, CORPS00, DMTP98	18	17	69	69	129	69	18	69	16	18	129	18	16	18	18	17	69	17	17	17	41	129
		Reference	PHASE1RA, DMTP98, BASLIN82	15	9	21	21	21	21	9	21	9	15	21	9	9	9	15	15	21	9	9	9	9	21
	Water	Site	PHASE1RA	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
		Reference	PHASE1RA	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6

**Note:** Survey names and citations: PHASE1RA Exponent (2003e) and Appendix A of this document  
 PSCHAR Exponent (2003c)  
 FUGDST01 Exponent (2002a)  
 SUPPRSS Exponent (2002b)  
 TECK03 Teck Cominco (2003)  
 TECK01 Exponent (2002a)  
 ENSR91 ENSR (1992)  
 ENSR92 ENSR (1993)  
 ENSR95 ENSR (1996)  
 ENSR96 RWJ (1997)  
 DMTP98 Cominco et al. (1999)  
 CORPS00 Corps (2001)  
 BASLIN82 Dames & Moore (1983a)  
 USGS02 Brabets (2003, pers. comm.)

<sup>a</sup> The numbers of samples shown are for the data to be used in the assessment, processed according to the data usability criteria listed in Section 3.2 of the main text.

Table 3-4. Statistical comparison of site and reference soil data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	<i>p</i> -values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	10	1,640	12,400	6,963	4,351	51	1,180	16,600	7,392	3,281	no	0.38	0.42
Antimony											≥50% ND Site		
Arsenic	10	4.2	35.0	12.6	9.8	75	1.3	93.6	12.2	15.1	no	0.76	0.47
<b>Barium</b>	5	109	622	249	213	40	357	7,090	2,137	1,830	<b>yes</b>	0.0002	<0.0001
<b>Cadmium</b>	10	0.24	3.6	1.1	1.1	478	0.40	388	25.2	37.8	<b>yes</b>	<0.0001	<0.0001
Chromium	5	4.9	19.3	11.8	5.7	40	4.9	24.0	15.0	5.1	no	0.12	0.16
Cobalt	5	7.3	20.6	13.5	5.1	40	4.2	27.0	11.3	5.0	no	0.86	0.32
<b>Copper</b>	5	14.3	46.5	23.7	13.0	40	9.8	109	36.0	20.3	<b>yes<sup>c</sup></b>	0.067	0.17
<b>Fluoride</b>	5	0.30	0.50	0.42	0.084	12	0.40	1.3	0.73	0.30	<b>yes</b>	0.018	0.025
Iron	10	5,750	72,600	29,872	18,432	51	2,650	35,000	20,682	7,572	no	0.99	0.20
<b>Lead</b>	10	8.8	142	38.5	38.5	479	8.5	48,300	1,157	2,795	<b>yes</b>	<0.0001	<0.0001
Manganese	5	250	4,080	1,489	1,589	40	280	1,000	513	186	no	0.76	0.0076 <sup>d</sup>
<b>Mercury</b>	5	0.050	0.18	0.11	0.053	12	0.10	1.7	0.45	0.51	<b>yes</b>	0.012	0.030
Molybdenum											≥50% ND Site		
Nickel	5	23.5	51.4	34.2	12.6	40	17.3	56.8	29.1	10.0	no	0.80	0.28
Selenium											≥50% ND Site		
<b>Silver</b>	5	0.050	0.25	0.13	0.089	40	0.14	8.3	2.2	2.0	<b>yes</b>	0.0004	<0.0001
<b>Strontium</b>	5	9.3	63.6	31.0	21.2	20	36.2	90.1	63.2	15.5	<b>yes</b>	0.0054	0.0001
<b>Thallium</b>	5	0.10	0.24	0.16	0.055	12	0.11	1.3	0.47	0.36	<b>yes</b>	0.026	0.041
Tin											≥50% ND Site		
<b>Vanadium</b>	5	5.6	19.2	11.9	5.4	40	7.9	31.8	14.7	4.8	<b>yes<sup>e</sup></b>	0.15	0.098
<b>Zinc</b>	10	72.5	753	181	204	479	37.4	64,300	4,140	6,201	<b>yes</b>	<0.0001	<0.0001

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> *p*-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Assumptions of ANOVA model were not met, thus conclusions result from non-parametric *p*-value.

<sup>d</sup> ANOVA *p*-value indicates reference mean is significantly higher than site mean concentration.

<sup>e</sup> Assumptions of ANOVA model were met, thus conclusions result from parametric *p*-value.

Table 3-5. Statistical comparison of site and reference tundra soil data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	10	368	11,300	3,651	3,347	31	358	18,900	5,329	4,822	no	0.22	0.39
Antimony	10	0.11	0.28	0.17	0.062	25	0.15	25.8	6.2	6.1	<sup>c</sup>	<0.0001	<0.0001
<b>Arsenic</b>	10	0.40	6.8	2.3	1.9	31	0.30	150	17.7	26.6	<b>yes</b>	0.0002	0.0003
Barium	10	108	624	315	196	25	53.0	5,810	945	1,306	<sup>c</sup>	0.032	0.10
<b>Cadmium</b>	10	0.12	0.88	0.35	0.22	224	0.30	438	15.3	31.7	<b>yes</b>	0.0016	0.0002
Chromium	10	1.6	19.7	6.8	6.1	25	1.0	33.2	10.4	8.7	no	0.19	0.33
Cobalt	10	0.96	28.3	8.6	10.6	25	0.50	35.0	11.0	9.3	<sup>c</sup>	0.14	0.36
<b>Copper</b>	10	4.3	16.9	8.2	4.0	25	2.9	58.3	21.0	15.5	<b>yes</b>	0.0027	0.0050
Fluoride											≥50% ND Site		
Iron	10	912	45,100	12,909	13,600	31	593	181,000	26,417	35,855	no <sup>d</sup>	0.095	0.19
<b>Lead</b>	10	2.9	23.3	8.9	6.7	264	7.0	16,000	665	1,816	<b>yes</b>	<0.0001	<0.0001
Manganese	10	33.5	6,620	918	2,013	25	28.6	3,400	825	882	<sup>c</sup>	0.078	0.27
Mercury	10	0.070	0.15	0.11	0.026	12	0.10	4.2	0.71	1.2	<sup>c</sup>	0.0003	0.0015
<b>Molybdenum</b>	10	0.34	2.3	0.85	0.60	25	0.59	3.9	1.5	0.91	<b>yes</b>	0.0048	0.013
Nickel	10	4.3	36.8	16.2	10.6	25	1.6	37.5	18.7	10.7	no	0.32	0.76
Selenium											≥50% ND Site		
Silver	10	0.020	0.35	0.14	0.13	25	0.040	14.7	2.5	3.3	<sup>c</sup>	<0.0001	<0.0001
<b>Strontium</b>	10	7.3	39.6	16.1	11.3	17	4.8	150	52.2	40.2	<b>yes</b>	0.0027	0.0031
Thallium	10	0.024	0.12	0.062	0.032	12	0.014	1.6	0.45	0.50	<sup>c</sup>	0.0031	0.0074
Tin											≥50% ND Site		
Vanadium	10	1.3	24.7	9.7	7.6	25	0.70	46.5	14.5	12.4	<sup>c</sup>	0.19	0.52
<b>Zinc</b>	10	47.8	111	66.1	24.2	264	22.3	82,700	2,127	4,880	<b>yes</b>	<0.0001	<0.0001

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.

<sup>d</sup> Assumptions of ANOVA model were met, thus conclusions result from parametric p-value.

Table 3-6. Statistical comparison of site and reference stream sediment data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	<i>p</i> -values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	5	3,620	12,100	6,848	3,652	14	4,080	17,100	7,846	3,560	no	0.27	0.46
<b>Antimony</b>	5	0.030	0.050	0.036	0.0089	14	0.050	0.64	0.20	0.16	<b>yes</b>	0.0008	0.0002
<b>Arsenic</b>	5	3.5	8.1	5.1	1.8	14	3.3	11.4	7.8	2.1	<b>yes</b>	0.023	0.023
Barium	5	135	483	291	146	14	91.2	922	302	260	no	0.76	0.68
<b>Cadmium</b>	5	0.070	0.30	0.22	0.088	14	0.18	1.4	0.49	0.34	<b>yes</b>	0.017	0.030
Chromium	5	7.2	19.9	12.7	5.0	14	7.4	22.6	14.6	4.9	no	0.20	0.46
<b>Cobalt</b>	5	7.3	11.0	9.3	1.4	14	7.9	17.6	12.3	2.9	<b>yes</b>	0.013	0.031
<b>Copper</b>	5	6.0	18.5	11.3	4.6	14	9.7	28.2	15.9	4.8	<b>yes</b>	0.039	0.041
Fluoride											≥50% ND Site		
<b>Iron</b>	5	21,300	27,300	24,500	2,279	14	22,800	45,700	30,479	5,898	<b>yes</b>	0.015	0.030
<b>Lead</b>	5	5.1	9.2	7.6	1.7	14	8.2	142	31.7	44.4	<b>yes</b>	0.0013	0.032
<b>Manganese</b>	5	268	859	548	259	14	471	2,140	995	542	<b>yes</b>	0.029	0.040
Mercury											≥50% ND Site		
<b>Molybdenum</b>	5	0.28	0.52	0.37	0.094	14	0.34	2.3	0.82	0.54	<b>yes</b>	0.0054	0.014
<b>Nickel</b>	5	20.8	35.0	29.7	5.3	14	24.8	57.3	40.4	8.8	<b>yes</b>	0.015	0.020
<b>Selenium</b>	5	0.10	0.70	0.44	0.22	14	0.40	2.5	1.2	0.65	<b>yes</b>	0.0078	0.0049
<b>Silver</b>	5	0.030	0.12	0.07	0.036	14	0.050	0.42	0.19	0.12	<b>yes</b>	0.0087	0.011
<b>Strontium</b>	5	4.9	15.0	11.6	4.1	14	11.0	155	45.5	42.6	<b>yes</b>	0.057	0.033
<b>Thallium</b>	5	0.023	0.070	0.050	0.019	14	0.031	0.322	0.10	0.076	<b>yes</b>	0.048	0.079
Tin											≥50% ND Site		
Vanadium	5	10.7	24.8	18.2	5.1	14	8.8	27.1	16.7	5.9	no	0.63	0.55
<b>Zinc</b>	5	43.7	69.7	62.4	10.7	14	58.4	259	139	52.0	<b>yes</b>	0.0024	0.0004

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> *p*-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

Table 3-7. Statistical comparison of site and reference stream surface water data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	3	17.3	2,770	937	1,588	230	2.5	4,060	90.4	413	no	0.88	0.038 <sup>d</sup>
Antimony											≥50% ND Site		
Arsenic											≥50% ND Site		
Barium	3	86.1	222	159	68.5	14	12.2	266	83.6	73.9	<sup>c</sup>	0.94	0.12
Cadmium											≥50% ND Site		
Chromium											≥50% ND Site		
Cobalt	3	0.12	2.7	1.0	1.5	14	0.010	0.33	0.13	0.094	no	0.94	0.057 <sup>d</sup>
Copper	3	0.60	5.4	2.2	2.7	18	0.11	1.2	0.74	0.33	no	0.71	0.15
<b>Fluoride</b>	3	30.0	40.0	36.7	5.8	31	40.0	120	57.4	17.1	<b>yes</b>	0.0047	0.0092
Iron	3	64.2	6,710	2,295	3,823	230	2.6	10,300	318	1,078	no	0.83	0.10
Lead											≥50% ND Site		
Manganese	3	4.9	128	46.0	71.0	18	0.48	36.0	5.7	8.2	<sup>c</sup>	0.91	0.054
Mercury											≥50% ND Site		
<b>Molybdenum</b>	3	0.020	0.17	0.080	0.079	14	0.18	2.3	0.74	0.69	<b>yes</b>	0.0048	0.0015
Nickel	3	1.1	10.5	4.5	5.2	14	0.26	6.7	2.1	1.8	no	0.76	0.32
Selenium											100% ND Ref.		
Silver											≥50% ND Site		
Strontium	3	32.5	81.1	54.9	24.5	14	19.4	172	83.0	55.5	no	0.29	0.61
Thallium											≥50% ND Site		
Tin											≥50% ND Site		
Vanadium											≥50% ND Site		
Zinc											≥50% ND Site		

**Note:** Concentrations are given in µg/L unfiltered.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.

<sup>d</sup> ANOVA p-value indicates reference mean is significantly higher than site mean concentration.

Table 3-8. Statistical comparison of site and reference pond sediment data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	5	3,730	17,100	9,908	5,750	4	1,920	4,330	3,288	1,214	no	0.97	0.039 <sup>d</sup>
<b>Antimony</b>	5	0.030	0.11	0.062	0.036	4	0.19	9.0	2.4	4.4	<b>yes</b>	0.0097	0.033
Arsenic	5	2.6	13.0	6.6	4.2	4	2.6	7.5	4.7	2.1	no	0.69	0.53
Barium	5	121	772	430	257	4	281	498	372	95.3	no	0.73	0.96
<b>Cadmium</b>	5	0.27	0.66	0.39	0.16	4	0.93	101	26.2	49.9	<b>yes</b>	0.010	0.056
Chromium	5	9.6	28.0	19.4	8.6	4	9.0	13.0	10.3	1.8	no	0.97	0.078 <sup>d</sup>
Cobalt	5	1.8	21.9	10.3	7.4	4	2.7	24.1	14.2	10.7	<sup>c</sup>	0.36	0.70
Copper	5	8.0	20.7	14.8	4.8	4	6.5	45.5	23.4	16.6	no	0.27	0.51
Fluoride											≥50% ND Site		
Iron	5	17,900	43,700	27,140	11,232	4	16,000	51,900	29,300	15,682	no	0.45	0.88
Lead	5	7.5	20.3	11.6	5.0	4	9.0	1,810	484	885	<sup>c</sup>	0.070	0.10
Manganese	5	15.9	1,870	515	766	4	79.8	745	290	312	<sup>c</sup>	0.73	0.98
Mercury	5	0.030	0.070	0.054	0.018	4	0.060	1.1	0.35	0.50	<sup>c</sup>	0.042	0.080
<b>Molybdenum</b>	5	0.38	1.4	0.70	0.40	4	1.1	2.4	1.7	0.66	<b>yes</b>	0.033	0.024
Nickel	5	12.0	70.3	39.0	21.0	4	17.6	38.9	27.6	8.7	no	0.86	0.52
Selenium	5	0.50	3.1	1.2	1.1	4	0.75	3.0	1.6	0.97	no	0.14	0.35
Silver											≥50% ND Site		
<b>Strontium</b>	5	4.2	25.4	12.2	8.1	4	17.1	86.0	37.5	32.5	<b>yes</b>	0.056	0.053
Thallium	5	0.056	0.17	0.12	0.051	4	0.021	1.6	0.43	0.81	no	0.91	0.74
Tin											≥50% ND Site		
Vanadium	5	14.9	94.5	40.8	31.1	4	12.2	28.3	17.8	7.2	no	0.94	0.11
<b>Zinc</b>	5	23.4	138	76.7	41.7	4	143	21,900	5,623	10,851	<b>yes</b>	0.010	0.083

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.

<sup>d</sup> ANOVA p-value indicates reference mean is significantly higher than site mean concentration.

Table 3-9. Statistical comparison of site and reference pond surface water data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	3	14.5	170	91.9	77.8	4	11.4	177	102	73.6	no	0.57	0.91
Antimony	3	0.020	0.10	0.057	0.040	4	0.020	0.20	0.085	0.083	<sup>c</sup>	0.50	0.78
Arsenic	3	0.50	0.90	0.63	0.23	4	0.40	1.3	0.70	0.41	no	0.57	0.93
Barium	3	48.4	133	91.6	42.3	4	39.4	73.6	57.5	17.0	no	0.94	0.23
Cadmium	3	0.0050	0.060	0.038	0.029	4	0.020	0.27	0.10	0.11	<sup>c</sup>	0.18	0.33
Chromium	3	0.18	2.0	0.96	0.92	4	0.44	5.2	2.2	2.1	<sup>c</sup>	0.30	0.33
Cobalt	3	0.19	0.70	0.37	0.29	4	0.13	1.6	0.76	0.61	no	0.30	0.46
Copper	3	0.70	2.5	1.9	1.0	4	0.40	2.7	1.3	0.98	no	0.70	0.48
Fluoride	3	10.0	50.0	26.7	20.8	4	20.0	60.0	32.5	18.9	no	0.29	0.57
Iron	3	361	1,500	808	608	4	685	1,220	1,021	238	no	0.30	0.35
Lead	3	0.060	0.56	0.37	0.27	4	0.44	1.6	0.95	0.52	no	0.11	0.15
Manganese	3	4.2	71.2	32.1	34.9	4	2.9	132	53.5	58.5	<sup>c</sup>	0.57	0.82
Mercury											≥50% ND Site		
Molybdenum	3	0.020	0.22	0.097	0.11	4	0.020	0.090	0.060	0.032	no	0.57	0.84
Nickel	3	2.1	6.4	3.6	2.4	4	3.0	5.3	4.3	1.1	no	0.30	0.44
Selenium											≥50% ND Site		
Silver											≥50% ND Site		
Strontium	3	10.6	27.5	18.7	8.5	4	10.4	422	114	205	no	0.70	0.68
Thallium											≥50% ND Site		
Tin											≥50% ND Site		
Vanadium	3	0.17	2.4	1.2	1.1	4	0.24	0.64	0.37	0.18	no	0.81	0.33
Zinc	3	0.59	5.0	2.8	2.2	4	6.1	99.0	36.7	42.9	<sup>c</sup>	0.026	0.048

**Note:** Concentrations are given in µg/L unfiltered.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.



Table 3-10. Statistical comparison of site and reference lagoon sediment data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	3	7,440	14,800	11,147	3,680	8	2,450	14,300	7,574	4,548	no	0.91	0.25
<b>Antimony</b>	3	0.010	0.12	0.077	0.059	8	0.070	0.27	0.16	0.073	<b>yes<sup>c</sup></b>	0.11	0.071
<b>Arsenic</b>	3	2.6	4.9	4.0	1.2	8	5.3	17.9	7.8	4.2	<b>yes</b>	0.0093	0.042
Barium	3	164	271	226	55.5	8	54.1	350	234	97.8	no	0.38	0.87
Cadmium	13										≥50% ND Site		
Chromium	3	12.5	24.9	19.6	6.4	8	4.1	27.2	13.8	8.6	no	0.82	0.29
Cobalt	3	5.0	9.7	6.8	2.5	8	3.9	11.8	7.1	2.8	no	0.62	0.96
Copper	3	9.9	18.7	14.7	4.5	8	3.0	28.2	14.2	8.6	no	0.54	0.66
Fluoride	3										≥50% ND Site		
Iron	3	14,000	22,200	19,233	4,546	8	10,100	75,000	27,150	21,985	no	0.62	0.71
<b>Lead</b>	28	2.4	31.0	11.1	6.6	26	4.7	302	44.4	68.8	<b>yes</b>	0.0017	0.0008
<b>Manganese</b>	3	75.5	129	99.9	27.1	8	97.9	274	158	55.9	<b>yes</b>	0.041	0.069
Mercury	3	0.030	0.060	0.050	0.017	8	0.0040	0.096	0.049	0.032	no	0.38	0.63
Molybdenum	3	0.46	0.98	0.77	0.28	8	0.41	3.4	1.4	1.2	no	0.30	0.48
Nickel	3	18.7	37.0	27.2	9.2	8	12.0	39.0	24.2	10.0	no	0.76	0.58
Selenium	3	0.60	1.4	1.1	0.44	8	0.10	2.2	1.0	0.69	no	0.73	0.58
Silver	3	0.010	0.11	0.067	0.051	8	0.020	0.27	0.12	0.083	no	0.15	0.32
Strontium	3	20.9	40.0	31.9	9.9	8	10.4	108	51.8	32.6	no	0.30	0.53
Thallium	3	0.038	0.10	0.081	0.037	8	0.018	0.18	0.075	0.056	no	0.73	0.65
Tin	3										≥50% ND Site		
Vanadium	3	16.8	31.5	25.2	7.6	8	8.5	35.1	21.7	10.7	no	0.73	0.51
<b>Zinc</b>	28	16.0	371	92.7	63.8	26	36.0	1,590	242	319	<b>yes</b>	0.0033	0.0026

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Assumptions of ANOVA model were met, thus conclusions result from parametric p-value.

Table 3-11. Statistical comparison of site and reference lagoon surface water data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	3	53.5	434	182	218	8	19.7	247	81.6	77.6	no	0.91	0.34
<b>Antimony</b>	3	0.11	0.13	0.12	0.010	8	0.19	0.63	0.32	0.15	<b>yes</b>	0.0093	0.0059
Arsenic	3	52.9	98.8	76.3	23.0	8	4.5	126	56.2	48.4	no	0.76	0.32
Barium	3	144	168	156	12.0	8	112	413	233	118	no	0.13	0.35
Cadmium	5	0.050	0.26	0.15	0.10	14	0.040	0.30	0.13	0.10	no	0.65	0.66
Chromium	3	6.0	8.2	7.2	1.1	8	1.7	4.5	2.7	1.1	no	0.99	0.0017 <sup>c</sup>
Cobalt	3	3.7	5.4	4.4	0.86	8	0.45	1.4	0.90	0.35	no	0.99	0.0002 <sup>c</sup>
Copper	3	0.40	1.4	0.80	0.53	8	0.50	1.4	0.96	0.29	no	0.27	0.35
<b>Fluoride</b>	3	10.0	20.0	13.3	5.8	8	50.0	200	114	73.3	<b>yes</b>	0.0083	0.0010
Iron	3	290	693	427	230	8	200	723	445	211	no	0.62	0.97
<b>Lead</b>	5	0.095	0.85	0.29	0.32	14	0.40	2.3	1.0	0.72	<b>yes</b>	0.0091	0.0015
Manganese	3	492	801	598	176	8	13.9	277	84.5	98.3	<sup>d</sup>	0.99	0.0043 <sup>c</sup>
Mercury											≥50% ND Site		
<b>Molybdenum</b>	3	0.070	0.090	0.080	0.010	8	0.30	2.4	1.2	0.85	<b>yes</b>	0.0095	0.0009
Nickel	3	9.2	15.2	11.5	3.2	8	3.5	10.6	7.1	2.7	no	0.96	0.077 <sup>c</sup>
Selenium											100% ND Ref.		
Silver	3	0.020	0.030	0.023	0.0058	8	0.010	0.25	0.11	0.092	<sup>d</sup>	0.090	0.14
Strontium	3	991	1,470	1,157	271	8	505	1,850	1,226	546	no	0.46	0.92
Thallium											≥50% ND Site		
Tin											≥50% ND Site		
Vanadium											100% ND Ref.		
Zinc	5	17.0	30.1	21.3	5.3	14	3.1	110	25.1	32.6	no	0.83	0.33

**Note:** Concentrations are given in µg/L unfiltered, except for cadmium, lead, and zinc, which are µg/L dissolved.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> ANOVA p-value indicates reference mean is significantly higher than site mean concentration.

<sup>d</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.

Table 3-12. Statistical comparison of site and reference marine sediment data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	<i>p</i> -values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	15	1,970	8,000	5,043	1,731	18	1,990	6,070	4,700	1,164	no	0.72	0.75
Antimony											≥50% ND Site		
Arsenic	21	5.6	13.0	8.7	1.9	69	3.1	14.5	7.3	1.8	no	1.0	0.0047 <sup>c</sup>
<b>Barium</b>	21	22.0	431	207	100	69	79.5	639	239	98.0	<b>yes<sup>d</sup></b>	0.15	0.048
<b>Cadmium</b>	21	0.020	0.23	0.068	0.044	129	0.020	52.9	1.0	4.7	<b>yes</b>	<0.0001	<0.0001
<b>Chromium</b>	21	1.4	18.0	11.9	4.4	69	2.4	33.5	14.6	5.7	<b>yes</b>	0.092	0.055
Cobalt	9	4.2	8.7	7.2	1.7	18	3.2	8.9	6.8	1.3	no	0.80	0.57
<b>Copper</b>	21	3.0	10.2	6.4	1.7	69	3.7	34.8	7.8	4.4	<b>yes</b>	0.064	0.067
Fluoride	9	0.40	2.0	1.3	0.51	16	0.40	1.5	1.1	0.29	no	0.74	0.55
Iron	15	8,150	22,700	15,149	4,973	18	9,960	19,300	15,987	2,570	no	0.21	0.33
Lead	21	2.7	11.2	5.3	1.7	129	1.6	5,620	58.5	494	<sup>e</sup>	<0.0001	0.0003
Manganese	9	187	389	301	71.7	18	161	363	276	59.6	no	0.77	0.42
Mercury											≥50% ND Site		
Molybdenum	9	0.44	0.83	0.57	0.12	18	0.37	1.4	0.58	0.26	no	0.77	0.77
Nickel	15	9.8	34.8	22.5	7.2	18	11.3	33.3	24.3	6.3	no	0.21	0.43
Selenium											≥50% ND Site		
<b>Silver</b>	21	0.020	0.49	0.12	0.17	69	0.030	2.1	0.50	0.61	<b>yes</b>	0.0007	0.013
<b>Strontium</b>	9	13.0	29.0	24.0	5.5	17	24.4	33.8	28.0	2.5	<b>yes</b>	0.073	0.018
Thallium	9	0.025	0.052	0.037	0.0096	17	0.026	1.1	0.098	0.27	no	0.66	0.65
Tin											≥50% ND Site		
Vanadium	9	13.0	33.9	22.2	6.6	41	9.1	46.0	27.9	9.7	no <sup>d</sup>	0.045	0.19
<b>Zinc</b>	21	25.0	56.8	42.2	8.5	129	5.5	2,550	87.3	237	<b>yes</b>	0.0032	0.070

**Note:** Concentrations are given in mg/kg dry weight.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of alpha = 0.10.

<sup>b</sup> *p*-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> ANOVA *p*-value indicates reference mean is significantly higher than site mean concentration.

<sup>d</sup> Assumptions of ANOVA model were met, thus conclusions result from parametric *p*-value.

<sup>e</sup> Comparison not made because 90 percent confidence interval for the site mean concentration spans zero, due to small sample size and/or high variability.

Table 3-13. Statistical comparison of site and reference marine surface water data

Chemical	Reference					Site					Site > Reference? <sup>a</sup>	p-values <sup>b</sup>	
	N	Min	Max	Mean	Stdev	N	Min	Max	Mean	Stdev		non-para.	parametric
Aluminum	6	25.0	336	170	148	9	25.0	205	102	69.7	no	0.74	0.66
Antimony	6	0.20	1.7	0.70	0.62	9	0.20	1.9	0.64	0.55	no	0.52	0.97
Arsenic	6	1.1	7.5	3.8	2.9	9	1.5	6.0	3.2	1.7	no	0.57	0.91
Barium	6	9.9	38.1	23.2	14.1	9	12.1	39.4	21.3	10.6	no	0.38	0.96
Cadmium	6	2.3	4.7	3.5	1.2	9	1.6	4.6	2.9	1.3	no	0.94	0.34
Chromium											≥50% ND Site		
Cobalt	6	4.0	4.5	4.3	0.17	9	3.9	4.6	4.2	0.22	no	0.76	0.61
Copper											≥50% ND Site		
Fluoride	6	600	800	700	110	9	500	900	733	158	no	0.31	0.75
Iron	6	33.6	643	314	306	9	52.3	375	171	133	no	0.52	0.80
Lead	6	0.76	1.3	0.99	0.18	9	0.80	1.3	1.0	0.21	no	0.48	0.88
Manganese	6	10.1	25.5	17.0	7.3	9	13.1	31.9	19.2	5.9	no	0.34	0.40
Mercury											≥50% ND Site		
Molybdenum	6	8.3	10.6	9.5	0.93	9	8.4	11.0	9.9	1.1	no	0.14	0.47
Nickel											≥50% ND Site		
<b>Selenium</b>	6	0.20	0.50	0.28	0.12	9	0.20	1.0	0.49	0.24	<b>yes</b>	0.027	0.047
<b>Silver</b>	6	0.10	0.27	0.18	0.066	9	0.10	0.95	0.40	0.28	<b>yes<sup>c</sup></b>	0.082	0.11
<b>Strontium</b>	6	4,530	5,290	4,900	369	9	4,420	5,600	5,128	444	<b>yes<sup>c</sup></b>	0.088	0.33
Thallium											≥50% ND Site		
Tin											≥50% ND Site		
Vanadium											≥50% ND Site		
Zinc											≥50% ND Site		

**Note:** Concentrations are given in  $\mu\text{g/L}$  unfiltered.

Undetected values are included at one-half the detection limit. In cases where greater than or equal to 50 percent of the site values or 100 percent of the reference values were undetected, statistical analyses were not performed. Further summary information is provided in the CoPC screening tables, including detection limits and detection frequencies.

Field replicates were averaged prior to statistical analysis.

ANOVA - analysis of variance

CoPC - chemical of potential concern

ND - not detected

<sup>a</sup> Results of statistical comparison. Bold indicates chemicals for which statistical testing indicated site concentrations to be greater than reference concentrations at a significance level of  $\alpha = 0.10$ .

<sup>b</sup> p-values associated with comparison of site and reference mean concentrations.

Non-para. - Wilcoxon rank-sum one-sided test for determining if the site mean is significantly greater than the reference mean concentration.

Parametric - overall ANOVA model for determining if the site and reference mean concentrations are significantly different (higher or lower).

<sup>c</sup> Assumptions of ANOVA model were not met, thus conclusions result from non-parametric p-value.

Table 3-14. Human health screening results for surface soil

Scenario Timeframe: Current/Future  
 Medium: Soil  
 Exposure Medium: Surface Soil

Exposure Point	Chemical	Minimum Detected Concentration	Maximum Detected Concentration	Units	Location of Maximum Concentration	Detection Frequency	Range of Detection Limits	Concentration Used for Screening <sup>a</sup>	Reference Range <sup>b</sup>	Residential Screening Toxicity Value <sup>c</sup>	Frequency of Detected Values Exceeding Criteria	Frequency of Reference Values Exceeding Criteria	Non-Residential Screening Toxicity Value <sup>d</sup>	CoPC Flag (Y/N)	Rationale for Selection or Deletion
<b>All Site Surface Soil</b>															
	Aluminum	1,180	16,600	mg/kg	RF-05	51/51	--	16,600	1,640–12,400	13,688 N	2/51	0/10	255,500 N	No	REF
	Antimony	0.38 <i>J</i>	14.8	mg/kg	CAG-W29	13/40	5–26	14.8	0.17–0.6	5.5 N	1/40	0/5	102 N	Yes	ASL
	Arsenic	1.3	93.6	mg/kg	CAG-H30	54/75	10–51	93.6	4.15–35	0.8 C	54/75	10/10	77 C	No	REF
	Barium	357	7,090	mg/kg	RF-07	40/40	--	7,090	109–622	960 N	35/40	0/5	17,885 N	Yes	ASL
	Cadmium	1.0	388 <i>J</i>	mg/kg	CAG-H30	430/478	0.4–2.5	388	0.24–3.59	14 N	236/478	0/10	256 N	Yes	ASL
	Chromium	4.86	24	mg/kg	RF-05	40/40	--	24	4.94–19.3	41 N	0/40	0/5	767 N	No	REF/BSL
	Cobalt	4.21	27	mg/kg	RF-05	39/40	5–5	27	7.28–20.6	274 N	0/40	0/5	5,110 N	No	REF/BSL
	Copper	9.76	109	mg/kg	RAT5-0NA	40/40	--	109	14.3–46.5	548 N	0/40	0/5	10,220 N	No	BSL
	Fluoride	0.5 <i>J</i>	1.3 <i>J</i>	mg/kg	RF-16	9/12	0.4–0.4	1.3	0.3–0.5	821 N	0/12	0/5	15,330 N	No	BSL
	Iron	2,650	35,000	mg/kg	CAG-W29	51/51	--	35,000	5,750–72,600	4,106 N	49/51	10/10	76,650 N	No	REF
	Lead	13.5	48,300	mg/kg	1007468	467/479	8.5–12	48,300	8.75–142	400 N	279/479	0/10	1,000 N	Yes	ASL
	Manganese	280	1,000	mg/kg	170_C1	40/40	--	1,000	250–4,080	329 N	37/40	4/5	6,132 N	No	REF
	Mercury	0.1	1.69	mg/kg	RF-107	12/12	--	1.69	0.05–0.18	2.6 N	0/12	0/5	77 N	No	BSL
	Molybdenum	0.35	3.3	mg/kg	RF-07	16/40	0.9–5.1	3.3	0.27–2.8	68 N	0/40	0/5	1,278 N	No	BSL
	Nickel	17.3	56.8	mg/kg	RC-06-A	40/40	--	56.8	23.5–51.4	270 N	0/40	0/5	5,110 N	No	REF/BSL
	Selenium	0.3 <i>J</i>	3 <i>J</i>	mg/kg	RF-107	12/30	10–51	3	0.5–1	68 N	0/30	0/5	1,278 N	No	BSL
	Silver	0.14	8.3	mg/kg	RAT5-0NA	21/40	0.9–5.1	8.3	0.05–0.25	68 N	0/40	0/5	1,278 N	No	BSL
	Strontium	36.2	90.1	mg/kg	RF-16	20/20	--	90.1	9.3–63.6	8,213 N	0/20	0/5	153,300 N	No	BSL
	Thallium	0.112	1.32	mg/kg	RF-32	12/12	--	1.32	0.1–0.24	0.9 N	1/12	0/5	17 N	Yes	ASL
	Tin	3.9 <i>J</i>	6 <i>J</i>	mg/kg	RF-27	2/27	2.25–26	6	ND	8,213 N	0/27	0/5	153,300 N	No	BSL
	Vanadium	7.94	31.8	mg/kg	RF-05	40/40	--	31.8	5.62–19.2	96 N	0/40	0/5	1,789 N	No	BSL
	Zinc	37.4	64,300	mg/kg	CAG-H30	479/479	--	64,300	72.5–753	4,100 N	158/479	0/10	76,650 N	Yes	ASL

**Note:** All results reported as dry weight.

For the purposes of screening, field replicates have been averaged.

- - not applicable
- C - carcinogenic based on a cancer risk of  $1 \times 10^{-6}$
- CoPC - chemical of potential concern
- J* - estimated value
- N - noncarcinogenic based on hazard quotient of 0.1
- ND - not detected

**Rationale Codes:**

**Selection Reason:**

- ASL - above screening levels

**Deletion Reason:**

- BSL - below screening level
- REF - below or consistent with reference levels

<sup>a</sup> The maximum detected soil concentration was used for screening CoPCs.

<sup>b</sup> The reference range corresponds to road material site soil samples from areas not affected by fugitive dust.

<sup>c</sup> Residential screening toxicity values represent arctic zone soil cleanup levels (from 18 AAC 75.341, Table B1) divided by 10. Where no Table B1 value exists, screening values were calculated based on residential formulas and input parameters provided in DEC (2002).

<sup>d</sup> Non-residential screening toxicity values using industrial formulas and input parameters provided in DEC (2000).

Table 3-15. Human health screening results for drinking water ingestion in stream surface water

Scenario Timeframe: Current/Future  
 Medium: Water  
 Exposure Medium: Stream Surface Water for Drinking Water Ingestion

Exposure Point	Chemical	Minimum Detected	Maximum Detected	Location of Maximum Concentration	Detection Frequency	Range of Detection Limits	Concentration Used for Screening <sup>a</sup>	Reference Range <sup>b</sup>	Screening Toxicity Value <sup>c</sup>	Potential ARAR/TBC Value	Potential ARAR/TBC Source <sup>d</sup>	Frequency of Detected Values Exceeding	Frequency of Reference Values Exceeding	CoPC Flag (Y/N)	Rationale for Selection or Deletion	
		Concentration	Concentration									Criteria	Criteria			
<b>All Site Stream Surface Water</b>																
	Aluminum	6.45	4,060	µg/L	StrRd	133/230	2.52–10	4,060	17.3–2,770	3,650 N	50–200	MCL	2/230	0/3	No	REF
	Antimony	0.14	0.6	µg/L	NHDowRd	6/14	0.063	0.6	ND–0.08	0.6 N	6	MCL	1/14	0/3	No	BSL
	Arsenic	ND	ND	µg/L	--	0/14	0.482	ND	ND–2.2	5 C	50	MCL	ND	0/3	No	ND/BSL
	Barium	12.2	266	µg/L	NHNFUp	14/14	--	266	86.1–222	200 N	2,000	MCL	1/14	1/3	No	REF
	Cadmium	0.03	0.40	µg/L	Various	24/229	0.02–0.25	0.4	0.01–0.07	0.5 N	5	MCL	0/229	0/3	No	BSL
	Chromium	ND	ND	µg/L	--	0/18	0.4	ND	0.17–3.71	10 N	100	MCL	ND	0/3	No	ND/BSL
	Cobalt	0.03	0.33	µg/L	NHRoad	12/14	0.01	0.33	0.12–2.72	73 N	--	--	0/14	0/3	No	BSL/REF
	Copper	0.3	1.2	µg/L	OmiDowRd	16/18	0.11	1.2	0.6–5.4	130 N	1,300	MCL	0/18	0/3	No	BSL/REF
	Fluoride	40	120	µg/L	NHRoad	27/31	50	120	30–40	219,000 N	--	--	0/31	0/3	No	BSL
	Iron	6	10,300	µg/L	StrRd	186/230	2.57–25	10,300	64.2–6,710	1,095 N	300	MCL	11/230	1/3	No	REF
	Lead	0.0	7.34	µg/L	StrDowRd	84/230	0.02–0.401	7.34	0.02–1.91	1.5 N	15	MCL	5/230	1/3	Yes	ASL
	Manganese	0.56	36	µg/L	MudLkCr	18/18	--	36	4.87–128	87.6 N	50	MCL	0/18	1/3	No	BSL/REF <sup>e</sup>
	Mercury	ND	ND	µg/L	--	0/14	0.0179	ND	ND	0.2 N	2	MCL	ND	ND	No	ND/BSL
	Molybdenum	0.37	2.27	µg/L	NHDowRd	11/14	0.178	2.27	0.05–0.17	18.25 N	--	--	0/14	0/3	No	BSL
	Nickel	0.26	6.71	µg/L	NHRoad	14/14	--	6.71	1.06–10.5	10 N	100	MCL	0/14	1/3	No	BSL/REF
	Selenium	0.067	1.24	µg/L	TutMth	15/29	0.0201	1.24	ND	5 N	50	MCL	0/29	ND	No	BSL
	Silver	ND	ND	µg/L	--	0/14	0.023	ND	ND–0.03	18 N	100	MCL	ND	0/3	No	ND/BSL
	Strontium	19.4	172	µg/L	NHDowRd	14/14	--	172	32.5–81.1	2,190 N	--	--	0/14	0/3	No	BSL/REF
	Thallium	0.04	0.55	µg/L	AufRd	9/29	0.0155–0.07	0.55	ND–0.014	0.2 N	2	MCL	1/29	0/3	Yes	ASL
	Tin	1.3	5.33	µg/L	OmiNFUp	5/14	0.59	5.33	ND	2,190 N	--	--	0/14	ND	No	BSL
	Vanadium	0.67	0.93	µg/L	ARC-U	4/14	0.335	0.93	0.16–5.57	26 N	--	--	0/14	0/3	No	BSL/REF <sup>e</sup>
	Zinc	1.0	60.1	µg/L	TutDowRd	107/230	0.5–5	60	0.31–9.84	1,100 N	5,000	MCL	0/230	0/3	No	BSL

**Note:** All results reported as unfiltered.  
 For the purposes of screening, field replicates have been averaged.  
 -- - not applicable  
 ARAR - applicable or relevant and appropriate requirement  
 C - carcinogenic based on a cancer risk of 1x10<sup>-6</sup>  
 CoPC - chemical of potential concern  
 J - estimated value  
 MCL - maximum contaminant level  
 N - noncarcinogenic based on hazard quotient of 0.1  
 ND - not detected  
 TBC - to be considered

**Rationale Codes:**  
 Selection Reason:  
 ASL - above screening levels  
 Deletion Reason:  
 BSL - below screening level  
 ND - not detected in any site sample  
 REF - below or consistent with reference levels

<sup>a</sup> The maximum detected stream surface water concentration was used for screening CoPCs.

<sup>b</sup> The reference range corresponds to stream surface water samples taken from areas not affected by fugitive dust.

<sup>c</sup> Screening toxicity values represent arctic zone drinking water cleanup levels (from 18 AAC 75.345, Table C) divided by 10. Where no Table C value exists, screening values were calculated based on residential drinking water formulas and input parameters provided in DEC (2002).

<sup>d</sup> An ARAR listed as an MCL is a maximum contaminant level derived by EPA, and is considered protective of the water body for use as the sole domestic drinking water source.

<sup>e</sup> The maximum site concentration of the analyte was less than the maximum reference concentration.

Table 3-16. Human health screening results for fish consumption in stream surface water

Scenario Timeframe: Current/Future  
 Medium: Water  
 Exposure Medium: Stream Surface Water for Fish Consumption

Exposure Point	Chemical	Minimum Detected	Maximum Detected	Units	Location of Maximum Concentration	Detection Frequency	Range of Detection Limits	Concentration Used for Screening <sup>a</sup>	Reference Range <sup>b</sup>	Screening Toxicity Value <sup>c</sup>	Potential ARAR/TBC Value	Potential ARAR/TBC Source <sup>d</sup>	Frequency of Detected Values Exceeding Criteria	Frequency of Reference Values Exceeding Criteria	CoPC Flag (Y/N)	Rationale for Selection or Deletion
		Concentration	Concentration		Concentration	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency	Frequency
<b>All Site Stream Surface Water</b>																
	Aluminum	6.45	4,060	µg/L	StrRd	133/230	2.52–10	4,060	17.3–2,770	--	--	--	--	--	No	REF
	Antimony	0.14	0.6	µg/L	NHDowRd	6/14	0.063	0.6	ND–0.08	1.4 N	--	--	0/14	0/3	No	BSL
	Arsenic	ND	ND	µg/L	--	0/14	0.482	ND	ND–2.2	0.018 C	0.00982	WDOE	ND	1/3	No	ND/BSL
	Barium	12.2	266	µg/L	NHNFUp	14/14	--	266	86.1–222	--	--	--	--	--	No	REF
	Cadmium	0.03	0.40	µg/L	Various	24/229	0.02–0.25	0.4	0.01–0.07	--	5.06	WDOE	--	--	No	BWC
	Chromium	ND	ND	µg/L	--	0/18	0.4	ND	0.17–3.71	--	203	WDOE	--	--	No	ND/BWC
	Cobalt	0.03	0.33	µg/L	NHRoad	12/14	0.01	0.33	0.12–2.72	--	--	--	--	--	No	REF
	Copper	0.3	1.2	µg/L	OmiDowRd	16/18	0.11	1.2	0.6–5.4	130 N	2,660	WDOE	0/18	0/3	No	BSL/REF
	Fluoride	40	120	µg/L	NHRoad	27/31	50	120	30–40	--	--	--	--	--	No	NSC
	Iron	6	10,300	µg/L	StrRd	186/230	2.57–25	10,300	64.2–6,710	--	--	--	--	--	No	REF
	Lead	0.0	7.34	µg/L	StrDowRd	84/230	0.02–0.401	7.34	0.02–1.91	--	--	--	--	--	No	NSC
	Manganese	0.56	36	µg/L	MudLkCr	18/18	--	36	4.87–128	5 N	--	--	--	--	No	REF <sup>e</sup>
	Mercury	ND	ND	µg/L	--	0/14	0.0179	ND	ND	0.005 N	--	--	--	--	No	ND
	Molybdenum	0.37	2.27	µg/L	NHDowRd	11/14	0.178	2.27	0.05–0.17	--	--	--	--	--	No	NSC
	Nickel	0.26	6.71	µg/L	NHRoad	14/14	--	6.71	1.06–10.5	61 N	1,100	WDOE	0/14	1/3	No	BSL/REF
	Selenium	0.067	1.24	µg/L	TutMth	15/29	0.0201	1.24	ND	17 N	--	--	0/29	ND	No	BSL
	Silver	ND	ND	µg/L	--	0/14	0.023	ND	ND–0.03	--	6,480	WDOE	--	--	No	ND/BWC
	Strontium	19.4	172	µg/L	NHDowRd	14/14	--	172	32.5–81.1	--	--	--	--	--	No	REF
	Thallium	0.04	0.55	µg/L	AufRd	9/29	0.0155–0.07	0.55	ND–0.014	0.17 N	1.56	WDOE	3/29	0/3	Yes	ASL
	Tin	1.3	5.33	µg/L	OmiNFUp	5/14	0.59	5.33	ND	--	--	--	--	--	No	NSC
	Vanadium	0.67	0.93	µg/L	ARC-U	4/14	0.335	0.93	0.16–5.57	--	--	--	--	--	No	REF <sup>e</sup>
	Zinc	1.0	60.1	µg/L	TutDowRd	107/230	0.5–5	60	0.31–9.84	910 N	16,500	WDOE	0/230	0/3	No	BSL

**Note:** All results reported as unfiltered.  
 For the purposes of screening, field replicates have been averaged.  
 -- - not applicable  
 ARAR - applicable or relevant and appropriate requirement  
 AWQC - Alaska water quality criteria  
 C - carcinogenic based on a cancer risk of 1 × 10<sup>-6</sup>  
 CoPC - chemical of potential concern  
 J - estimated value  
 N - noncarcinogenic based on hazard quotient of 0.1  
 ND - not detected  
 TBC - to be considered  
 WDOE - Washington State Department of Ecology

**Rationale Codes:**  
 Selection Reason:  
 ASL - above screening levels  
 Deletion Reason:  
 BSL - below screening level  
 BWC - no AWQC available, but below WDOE surface water criteria for bioaccumulation in fish  
 ND - not detected in any site sample  
 NSC - no screening criteria  
 REF - below or consistent with reference levels

<sup>a</sup> The maximum detected stream surface water concentration was used for screening CoPCs.

<sup>b</sup> The reference range corresponds to stream surface water samples taken from areas not affected by fugitive dust.

<sup>c</sup> Screening toxicity values represent the AWQC protective for human consumption of fish/shellfish and domestic drinking water usage from the water body (DEC 2003a). The AWQC were modified, when necessary, to assume a target hazard quotient of 0.1. The arsenic screening toxicity value is a federal ambient water quality criteria (U.S. EPA 2002c) and assumes a target risk of 10<sup>-6</sup>. The ARAR represents the Washington State cleanup level for surface water and is protective of bioaccumulation into, and human consumption of, seafood (WDOE 1996).

<sup>d</sup> The ARARs represent the Washington State Department of Ecology cleanup level for surface water and are protective of bioaccumulation into, and human consumption of, fish (WDOE 1996).

<sup>e</sup> The maximum site concentration of the analyte was less than the maximum reference concentration.

Table 3-17. Human health screening results for lagoon water

Scenario Timeframe: Current/Future  
 Medium: Lagoon Water  
 Exposure Medium: Lagoon Water

Exposure Point	Chemical	Minimum Detected Concentration	Maximum Detected Concentration	Units	Location of Maximum Concentration	Detection Frequency	Range of Detection Limits	Concentration Used for Screening <sup>a</sup>	Reference Range <sup>b</sup>	Screening Toxicity Value <sup>c</sup>	Potential ARAR/TBC Value	Potential ARAR/TBC Source <sup>d</sup>	Frequency of Detected Values Exceeding Criteria	Frequency of Reference Values Exceeding Criteria	CoPC Flag (Y/N)	Rationale for Selection or Deletion
<b>Lagoon Water</b>																
	Aluminum	19.7	247	µg/L	IP-04	8/8		247	53.5–434	--	--	--	--	--	No	REF
	Antimony	0.19	0.63	µg/L	PLNL	8/8		0.63	0.11–0.13	430 N	--	--	0/8	0/3	No	BSL
	Arsenic	4.5	126	µg/L	IP-04	8/8		126	52.9–98.8	0.14 C	0.00982	WDOE	8/8	3/3	No	REF
	Barium	112	413	µg/L	PLNL	8/8		413	144–168	--	--	--	--	--	No	REF
	Cadmium	0.04 J	0.3	µg/L	NLH	11/14	0.1	0.3	ND–0.26	--	5.06	WDOE	--	--	No	REF/BWC
	Chromium	1.69	4.49	µg/L	IP-04	8/8		4.49	5.96–8.22	--	203	WDOE	--	--	No	REF/BWC
	Cobalt	0.45	1.38	µg/L	PLNL	8/8		1.38	3.7–5.35	--	--	--	--	--	No	REF
	Copper	0.5 J	1.4	µg/L	IP-03	8/8		1.4	0.4–1.4	--	2,660	WDOE	--	--	No	REF/BWC
	Fluoride	50 J	200	µg/L	IP-01,IP-02,IP-04	8/8		200	ND–20	--	--	--	--	--	No	NSC
	Iron	200	723	µg/L	PLNN	8/8		723	290–693	--	--	--	--	--	No	REF
	Lead	0.4	2.3	µg/L	PLNP	14/14		2.3	ND–0.85	--	--	--	--	--	No	NSC
	Manganese	13.9	277	µg/L	PLNN	8/8		277	492–801	10 N	--	--	--	--	No	REF
	Mercury	ND	ND	µg/L	ND	0/8		ND	ND	0.005 N	--	--	--	--	No	ND
	Molybdenum	0.3	2.41	µg/L	IP-04	8/8	0.1	2.41	0.07–0.09	--	--	--	--	--	No	NSC
	Nickel	3.5	10.6	µg/L	IP-01	8/8		10.6	9.9–15.2	460 N	1,100	WDOE	0/8	0/3	No	REF/BSL
	Selenium	0.3 J	0.6 J	µg/L	PLNN	5/8	0.4	0.6	ND	1,100 N	--	--	0/8	0/3	No	BSL
	Silver	0.01 J	0.25	µg/L	PLNL	7/8	0.1	0.25	0.02–0.03	--	6,480	WDOE	--	--	No	BWC
	Strontium	505	1,850	µg/L	PLNN	8/8		1,850	991–1,470	--	--	--	--	--	No	REF
	Thallium	0.007 J	0.07 J	µg/L	NLF,PLNL	4/8	0.026-0.06	0.07	0.006–0.009	0.63 N	1.56	WDOE	0/8	0/3	No	BSL
	Tin	23.7 J	23.7 J	µg/L	NLF	1/8	20	23.7	ND	--	--	--	--	--	No	NSC
	Vanadium	0.22	0.85 J	µg/L	IP-04	5/8	0.36-0.8	0.85	ND	--	--	--	--	--	No	NSC
	Zinc	3.09 J	110	µg/L	NLH	14/14		110	ND–30.1	6,900 N	16,500	WDOE	0/14	0/3	No	REF/BSL

**Note:** All results reported as unfiltered.

For the purposes of screening, field replicates have been averaged.

-- - not applicable

ARAR - applicable or relevant and appropriate requirement

AWQ - ambient water quality criteria

C - carcinogenic based on a cancer risk of  $1 \times 10^{-6}$

CoPC - chemical of potential concern

J - estimated value

N - noncarcinogenic based on hazard quotient of 0.1

ND - not detected

TBC - to be considered

WDO - Washington State Department of Ecology

**Rationale Codes:**

Selection Reason:

ASL - above screening levels

Deletion Reason:

BSL - below screening level

BWC - no AWQC available, but below WDOE surface water criteria for bioaccumulation in fish

ND - not detected in any site sample

NSC - no screening criteria

REF - below or consistent with reference levels

<sup>a</sup> The maximum detected lagoon surface water concentration was used for screening CoPCs.

<sup>b</sup> The reference range corresponds to lagoon surface water samples taken from areas not affected by fugitive dust.

<sup>c</sup> Screening toxicity values represent the AWQC protective for human consumption of fish/shellfish and domestic drinking water usage from the water body (DEC 2003a). The AWQC were modified, when necessary, to assume a target hazard quotient of 0.1. The arsenic screening toxicity value is a federal ambient water quality criterion (U.S. EPA 2002c) and assumes a target risk of  $10^{-6}$ . The ARAR represents the Washington State cleanup level for surface water and is protective of bioaccumulation into, and human consumption of, seafood (WDOE 1996).

<sup>d</sup> The ARARs represent the Washington State Department of Ecology cleanup level for surface water and are protective of bioaccumulation into, and human consumption of, fish (WDOE 1996).



Table 3-18. Human health screening results for marine surface water

Scenario Timeframe: Current/Future  
 Medium: Marine Water  
 Exposure Medium: Marine Surface Water

Exposure Point	Chemical	Minimum Detected	Maximum Detected	Location of Maximum Concentration	Detection Frequency	Range of Detection Limits	Concentration Used for Screening <sup>a</sup>	Reference Range <sup>b</sup>	Screening Toxicity Value <sup>c</sup>	Potential ARAR/TBC Value	Potential ARAR/TBC Source <sup>d</sup>	Frequency of Detected Values Exceeding Criteria	Frequency of Reference Values Exceeding Criteria	CoPC Flag (Y/N)	Rationale for Selection or Deletion	
		Concentration	Concentration			Units	Concentration	Concentration	Concentration	Concentration	Value	Value	Criteria	Criteria	Criteria	Criteria
<b>Marine Water</b>																
	Aluminum	43	205	µg/L	NML	8/9	25	205	ND-336	--	--	--	--	No	REF	
	Antimony	0.30 <i>J</i>	1.88 <i>J</i>	µg/L	NML	5/9	0.2-0.4	1.88	ND-1.67	430 N	--	0/9	0/6	No	REF/BSL	
	Arsenic	1.5 <i>J</i>	6.0 <i>J</i>	µg/L	NMAA	7/9	3	6	ND-7.5	0.14 C	0.00982	7/9	5/6	No	REF	
	Barium	12.1	39.4	µg/L	NMK	9/9	--	39.4	9.91-38.1	--	--	--	--	No	REF	
	Cadmium	1.6	4.6	µg/L	NML	9/9	--	4.6	2.27-4.69	--	5.06	WDOE	--	No	REF/BWC	
	Chromium	ND	ND	µg/L	--	0/9	1-2	ND	ND	--	203	WDOE	--	No	ND/BSL	
	Cobalt	3.85	4.60	µg/L	NMG	9/9	--	4.6	4.03-4.48	--	--	--	--	No	REF	
	Copper	1.0	3.6	µg/L	NML	4/9	4	3.6	ND-2.6	--	2,660	WDOE	--	No	BWC	
	Fluoride	500	900	µg/L	NMAA, NMG, NML	9/9	--	900	600-800	--	--	--	--	No	REF	
	Iron	52	375	µg/L	NMAA	9/9	--	375	33.6-643	--	--	--	--	No	REF	
	Lead	0.8 <i>J</i>	1.34	µg/L	NMAA	9/9	--	1.34	0.76-1.25	--	--	--	--	No	REF	
	Manganese	13.1	31.9	µg/L	NMK	9/9	--	31.9	10.1-25.5	10 N	--	--	--	No	REF	
	Mercury	ND	ND	µg/L	--	0/9	0.05	ND	ND	0.005 N	--	--	--	No	ND/BSL	
	Molybdenum	8.4	11	µg/L	NMAA, NMG, NML	9/9	--	11	8.26-10.6	--	--	--	--	No	REF	
	Nickel	ND	ND	µg/L	--	0/9	2-10	ND	ND	460 N	1,100	WDOE	0/9	0/6	No	ND/BSL
	Selenium	0.3 <i>J</i>	1 <i>J</i>	µg/L	NMG	8/9	0.2	1	ND-0.5	1,100 N	--	0/9	0/6	No	BSL	
	Silver	0.4	0.95	µg/L	NMAA	5/9	0.1-0.2	0.95	ND-0.27	--	6,480	WDOE	--	No	BWC	
	Strontium	4,420	5,600 <i>J</i>	µg/L	NMG, NML	9/9	--	5,600	4,530-5,290	--	--	--	--	No	NSC	
	Thallium	0.09 <i>J</i>	0.09 <i>J</i>	µg/L	NMAA	1/9	0.05-0.1	0.09	ND-0.133	0.63 N	1.56	WDOE	0/9	0/6	No	BSL/REF <sup>e</sup>
	Tin	23.3 <i>J</i>	23.3 <i>J</i>	µg/L	NMAA	1/9	3-10	23.3	ND-26.4	--	--	--	--	No	REF <sup>e</sup>	
	Vanadium	4.44 <i>J</i>	5.27 <i>J</i>	µg/L	NMK	2/9	1.4-2	5.27	ND-8.44	--	--	--	--	No	REF <sup>e</sup>	
	Zinc	ND	ND	µg/L	--	0/9	1	ND	ND	6,900 N	16,500	WDOE	0/9	0/6	No	ND/BSL

**Note:** All results reported as unfiltered.

- For the purposes of screening, field replicates have been averaged.
- - not applicable
  - ARAR - applicable or relevant and appropriate requirement
  - AWQ - ambient water quality criteria
  - C - carcinogenic based on a cancer risk of  $1 \times 10^{-6}$
  - CoPC - chemical of potential concern
  - J* - estimated value
  - N - noncarcinogenic based on hazard quotient of 0.1
  - ND - not detected
  - TBC - to be considered
  - WDO - Washington State Department of Ecology

**Rationale Codes:**

- Selection Reason:
- ASL - above screening levels
- Deletion Reason:
- BSL - below screening level
  - BWC - no AWQC available, but below WDOE surface water criteria for bioaccumulation in fish
  - ND - not detected in any site sample
  - REF - below or consistent with reference levels

<sup>a</sup> The maximum detected marine surface water concentration was used for screening CoPCs.

<sup>b</sup> The reference range corresponds to marine surface water samples taken from areas not affected by fugitive dust.

<sup>c</sup> Screening toxicity values represent the AWQC protective for human consumption of fish/shellfish and domestic drinking water usage from the water body (DEC 2003a). The AWQC were modified, when necessary, a target hazard quotient of 0.1. The arsenic screening toxicity value is a federal ambient water quality criterion (U.S. EPA 2002c) and assumes a target risk of  $10^{-6}$ . The ARAR represents the Washington State clean surface water and is protective of bioaccumulation into, and human consumption of, seafood (WDOE 1996).

<sup>d</sup> The ARARs represent the Washington State Department of Ecology cleanup level for surface water and are protective of bioaccumulation into, and human consumption of, fish (WDOE 1996).

<sup>e</sup> The maximum site concentration of the analyte was less than the maximum reference concentration.

Table 3-19. Ecological screening results for tundra soil

Analyte	Detection Frequency	Reference Concentration			Ecological Screening Benchmark			Detection Frequency Above Benchmark <sup>a</sup>				
		Minimum Detected Concentration	Maximum Detected Concentration	Detection Limits	Minimum Detected Concentration	Maximum Detected Concentration	ORNL Terrestrial Plants <sup>b</sup>	ORNL Earthworms <sup>c</sup>	ORNL Soil Microorganisms <sup>c</sup>	ORNL Terrestrial Plants	ORNL Earthworms	ORNL Soil Microorganisms
		(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Aluminum	32/32	358	18,900	-	368	11,300	50	--	600	32/32	--	29/32
Antimony	14/26	0.15	25.8	7-29	0.11	0.28	5	--	--	3/26	--	--
Arsenic	17/32	0.3	150	19-58	0.4	6.8	10	60	100	6/32	1/32	1/32
Barium	26/26	53	5,810	-	108	624	500	--	3,000	13/26	--	3/26
Cadmium	132/231	0.53	438	0.6-4.3	0.12	0.88	4	20	20	85/231	51/231	51/231
Chromium	24/26	1.03	33.2	5.8	1.57	19.7	1	0.4	10	24/26	24/26	9/26
Cobalt	25/26	0.5	35	5.8	0.96	28.3	20	--	1,000	4/26	--	0/26
Copper	26/26	2.88	58.3	-	4.34	16.9	100	50	100	0/26	3/26	0/26
Fluoride	1/13	3.8	3.8	0.7	0.4 <sup>d</sup>	0.4 <sup>d</sup>	200	--	30	0/13	--	0/13
Iron	32/32	593	181,000	-	912	45,100	--	--	200	--	--	32/32
Lead	175/271	12.1	16,000	14-110.9	2.9	23.3	50	500	900	122/271	54/271	43/271
Manganese	26/26	28.6	3,400	-	33.5	6,620	500	--	100	16/26	--	23/26
Mercury	13/13	0.1	4.16	-	0.07	0.15	0.3	0.1	30	6/13	12/13	0/13
Molybdenum	14/26	0.59	3.9	1.5-5.8	0.34	2.27	2	--	200	4/26	--	0/26
Nickel	25/26	1.58	37.5	11.5	4.33	36.8	30	200	90	6/26	0/26	0/26
Selenium	13/26	0.3	3.3	15-58	0.4	1	1	70	100	6/26	0/26	0/26
Silver	17/26	0.04	14.7	1.5-5.8	0.07	0.35	2	--	50	6/26	--	0/26
Strontium	18/18	4.8	150	-	7.3	39.6	--	--	--	--	--	--
Thallium	13/13	0.014	1.58	-	0.024	0.116	1	--	--	3/13	--	--
Tin	6/18	7.7	14	4.2-29	5	17.4	50	--	2,000	0/18	--	0/18
Vanadium	25/26	0.7	46.5	5.8	1.3	24.7	2	--	20	22/26	--	7/26
Zinc	271/271	15	82,700	-	47.8	111	50	200	100	244/271	135/271	181/271

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark
- ORNL - Oak Ridge National Laboratory

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> Efroymson et al. (1997a).

<sup>c</sup> Efroymson et al. (1997b).

<sup>d</sup> Undetected; value listed is one-half of the detection limit.

Table 3-20. Ecological screening results for stream sediment

Analyte	Detection Frequency	Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Detection Limits (mg/kg)	Reference Concentration		Ecological Screening Benchmark			Detection Frequency Above Benchmark <sup>a</sup>		
					Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Threshold Effect Concentration <sup>b</sup> (mg/kg)	Probable Effect Concentration <sup>b</sup> (mg/kg)	No Effect Concentration <sup>c</sup> (mg/kg)	Threshold Effect Concentration	Probable Effect Concentration	No Effect Concentration
Aluminum	15/15	4,080	17,100	-	3,620	12,100	--	--	73,000	--	--	0/15
Antimony	15/15	0.05	0.64	-	0.03	0.05	--	--	--	--	--	--
Arsenic	15/15	3.3	11.4	-	3.5	8.1	9.79	33	100	1/15	0/15	0/15
Barium	15/15	91.2	922	-	135	483	--	--	--	--	--	--
Cadmium	15/15	0.18	1.38	-	0.07	0.3	0.99	4.98	8	1/15	0/15	0/15
Chromium	15/15	7.35	23.3	-	7.22	19.9	43.4	111	95	0/15	0/15	0/15
Cobalt	15/15	7.9	17.6	-	7.3	11	--	--	--	--	--	--
Copper	15/15	9.66	28.2	-	5.99	18.5	31.6	149	580	0/15	0/15	0/15
Fluoride	0/15	0.95 <sup>d</sup>	1.2 <sup>d</sup>	1.9-2.3	1.2 <sup>d</sup>	1.2 <sup>d</sup>	--	--	--	--	--	--
Iron	15/15	22,800	45,700	-	21,300	27,300	--	--	290,000	--	--	0/15
Lead	15/15	8.24	142	-	5.05	9.17	35.8	128	130	2/15	2/15	1/15
Manganese	15/15	471	2140	-	268	859	--	--	4,500	--	--	0/15
Mercury	15/15	0.02	0.089	-	0.02	0.04	0.18	1.06	--	0/15	0/15	--
Molybdenum	15/15	0.34	2.32	-	0.28	0.52	--	--	--	--	--	--
Nickel	15/15	24.8	57.3	-	20.8	35	22.7	48.6	43	15/15	1/15	7/15
Selenium	15/15	0.4	2.5	-	0.4	0.7	--	--	--	--	--	--
Silver	15/15	0.05	0.42	-	0.03	0.12	--	--	--	--	--	--
Strontium	15/15	11	155	-	4.9	15	--	--	--	--	--	--
Thallium	15/15	0.031	0.322	-	0.023	0.07	--	--	--	--	--	--
Tin	4/15	4.3	7.6	1-5	2 <sup>d</sup>	2.4 <sup>d</sup>	--	--	--	--	--	--
Vanadium	15/15	8.83	28.1	-	10.7	24.8	--	--	--	--	--	--
Zinc	15/15	58.4	259	-	43.7	69.7	121	459	1,300	7/15	0/15	0/15

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> MacDonald et al. (2000).

<sup>c</sup> Ingersoll et al. (1996).

<sup>d</sup> Undetected; value listed is one-half of the detection limit.

Table 3-21. Ecological screening results for stream surface water

Analyte	Detection Frequency	Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Detection Limits (µg/L)	Reference Concentration		Ecological Screening Benchmark		Detection Frequency Above Benchmark <sup>k</sup>	
					Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Freshwater Criteria Continuous Concentration <sup>b</sup> (µg/L total recoverable)	Freshwater Criterion Maximum Concentration <sup>b</sup> (µg/L total recoverable)	Freshwater Criteria Continuous Concentration	Freshwater Criterion Maximum Concentration
Hardness	231/231	11,300	211,000	-	56,000	112,000	--	--	--	--
Aluminum	126/230	6.45	4060	5.04–62.1	17.3	2,770	87	750	29/230	3/230
Antimony	6/14	0.14	0.63	0.126	0.08	0.08	--	--	--	--
Arsenic	0/14	0.482 <sup>c</sup>	0.482 <sup>c</sup>	0.964	2.2	2.2	150	340	0/14 <sup>d</sup>	0/14 <sup>d</sup>
Barium	14/14	12.2	266	-	86.1	222	--	--	--	--
Cadmium	24/229	0.03	0.8	0.04–0.5	0.01	0.07	0.05–0.47 (0.27) <sup>e</sup>	0.23–4.6 (2.1) <sup>e</sup>	3/229	0/229
Chromium	0/18	0.396 <sup>c</sup>	0.4 <sup>c</sup>	0.791–0.8	0.17	3.71	11	16	0/18 <sup>d</sup>	0/18 <sup>d</sup>
Cobalt	11/14	0.03	0.33	0.02–0.03	0.12	2.72	--	--	--	--
Copper	16/18	0.3	1.23	0.21	0.6	5.4	1.4–18 (9.4) <sup>e</sup>	1.8–28 (14) <sup>e</sup>	0/18	0/18
Fluoride	27/31	40	120	100	30	40	--	--	--	--
Iron	186/230	6	10300	5.13–50	64.2	6,710	1,000	--	12/230	--
Lead	84/230	0.018	7.34	0.04–0.802	0.02	1.91	0.20–8.2 (3.2) <sup>e</sup>	5.1–211 (82) <sup>e</sup>	22/230	0/230
Manganese	17/18	0.56	36	0.95	4.87	128	--	--	--	--
Mercury	0/14	0.0179 <sup>c</sup>	0.0179 <sup>c</sup>	0.0358	0.05 <sup>c</sup>	0.05 <sup>c</sup>	0.91	1.6	0/14 <sup>d</sup>	0/14 <sup>d</sup>
Molybdenum	9/14	0.37	2.27	0.355–0.48	0.05	0.17	--	--	--	--
Nickel	14/14	0.26	6.71	-	1.06	10.5	8.2–98 (52) <sup>e</sup>	74–882 (470) <sup>e</sup>	0/14	0/14
Selenium	15/29	0.0666	1.24	0.0402	0.2 <sup>c</sup>	0.2 <sup>c</sup>	5.0	--	0/29	--
Silver	0/14	0.023 <sup>c</sup>	0.023 <sup>c</sup>	0.046	0.03	0.03	--	0.09–14 (3.8) <sup>e</sup>	--	0/14 <sup>d</sup>
Strontium	14/14	19.4	172	-	32.5	81.1	--	--	--	--
Thallium	4/29	0.04	0.55	0.031–0.29	0.014	0.014	--	--	--	--
Tin	5/14	1.3	5.33	1.18	10 <sup>c</sup>	10 <sup>c</sup>	--	--	--	--
Vanadium	4/14	0.67	0.93	0.669	0.16	5.57	--	--	--	--
Zinc	107/230	1	60.1	1–10	0.31	9.84	19–226 (120) <sup>e</sup>	19–226 (120) <sup>e</sup>	1/230	1/230

**Note:** Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> U.S. EPA (2002c).

<sup>c</sup> Undetected; value listed is one-half of the detection limit.

<sup>d</sup> Undetected in all samples. Undetected values expressed as one-half of the detection limit are below the screening benchmark.

<sup>e</sup> Water quality criteria for this metal is hardness-dependent. The range shown was calculated using the minimum and maximum hardness values for stream surface water. EPA provides a default water quality criteria based on a hardness of 100 mg/L CaCO<sub>3</sub>. This value is presented in parentheses.

Table 3-22. Ecological screening results for tundra pond sediment

Analyte	Detection Frequency	Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Detection Limits (mg/kg)	Reference Concentration		Ecological Screening Benchmark			Detection Frequency Above Benchmark <sup>a</sup>		
					Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Threshold Effect Concentration <sup>b</sup> (mg/kg)	Probable Effect Concentration <sup>b</sup> (mg/kg)	No Effect Concentration <sup>c</sup> (mg/kg)	Threshold Effect Concentration	Probable Effect Concentration	No Effect Concentration
Aluminum	5/5	1,920	5,270	-	3,730	17,100	--	--	73,000	--	--	0/5
Antimony	5/5	0.19	10.9	-	0.03	0.11	--	--	--	--	--	--
Arsenic	5/5	2.6	8.7	-	2.6	13	9.79	33	100	0/5	0/5	0/5
Barium	5/5	281	626	-	121	772	--	--	--	--	--	--
Cadmium	5/5	0.93	119	-	0.27	0.66	0.99	4.98	8	3/5	2/5	2/5
Chromium	5/5	8.97	15.3	-	9.57	28	43.4	111	95	0/5	0/5	0/5
Cobalt	5/5	2.66	25.9	-	1.83	21.9	--	--	--	--	--	--
Copper	5/5	6.51	53.4	-	7.99	20.7	31.6	149	580	2/5	0/5	0/5
Fluoride	3/5	2.6	4.4	2.3	1.2 <sup>d</sup>	1.2 <sup>d</sup>	--	--	--	--	--	--
Iron	5/5	16,000	51,900	-	17,900	43,700	--	--	290,000	--	--	0/5
Lead	5/5	8.96	2,180	-	7.48	20.3	35.8	128	130	3/5	2/5	2/5
Manganese	5/5	60.2	745	-	15.9	1,870	--	--	4,500	--	--	0/5
Mercury	5/5	0.06	1.31	-	0.03	0.07	0.18	1.06	--	2/5	1/5	--
Molybdenum	5/5	1.05	2.84	-	0.38	1.35	--	--	--	--	--	--
Nickel	5/5	17.6	44.2	-	12	70.3	22.7	48.6	43	4/5	0/5	1/5
Selenium	4/5	1.2	3.5	1.5	0.5	3.1	--	--	--	--	--	--
Silver	3/5	0.09	3.76	0.1–0.15	0.06	0.18	--	--	--	--	--	--
Strontium	5/5	17.1	111	-	4.2	25.4	--	--	--	--	--	--
Thallium	4/5	0.021	1.92	0.046	0.056	0.174	--	--	--	--	--	--
Tin	3/5	18.2	41.2	15.4–21.3	2.1 <sup>d</sup>	6.3 <sup>d</sup>	--	--	--	--	--	--
Vanadium	5/5	10.2	28.3	-	14.9	94.5	--	--	--	--	--	--
Zinc	5/5	143	27,000	-	23.4	138	121	459	1,300	5/5	2/5	2/5

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> MacDonald et al. (2000).

<sup>c</sup> Ingersoll et al. (1996).

<sup>d</sup> Undetected; value listed is one-half of the detection limit.

Table 3-23. Ecological screening results for tundra pond surface water

Analyte	Detection Frequency	Reference Concentration		Detection Limits (µg/L)	Ecological Screening Benchmark		Detection Frequency Above Benchmark <sup>a</sup>			
		Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)		Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Freshwater Criteria Continuous Concentration <sup>b</sup>	Freshwater Criterion Maximum Concentration <sup>b</sup>	Freshwater Criteria Continuous Concentration	Freshwater Criterion Maximum Concentration
Hardness	5/5	10,300	382,000	-	12,300	34,400	--	--	--	--
Aluminum	5/5	11.4	180	-	14.5	170	87	750	3/5	0/5
Antimony	3/5	0.03	0.2	0.04	0.02	0.1	--	--	--	--
Arsenic	5/5	0.4	1.7	-	0.5	0.9	150	340	0/5	0/5
Barium	5/5	39.4	74.5	-	48.4	133	--	--	--	--
Cadmium	5/5	0.02	0.27	-	0.05	0.06	0.05–0.73 (0.27) <sup>c</sup>	0.21–8.3 (2.1) <sup>c</sup>	1/5	0/5
Chromium	5/5	0.44	6.31	-	0.18	1.98	11	16	0/5	0/5
Cobalt	5/5	0.13	1.56	-	0.19	0.7	--	--	--	--
Copper	5/5	0.4	2.8	-	0.7	2.5	1.3–29 (9.4) <sup>c</sup>	1.6–49 (14) <sup>c</sup>	2/5	0/5
Fluoride	5/5	20	60	-	20	50	--	--	--	--
Iron	5/5	685	1,220	-	361	1,500	1,000	--	4/5	--
Lead	5/5	0.44	1.63	-	0.06	0.56	0.18–18 (3.2) <sup>c</sup>	4.5–450 (82) <sup>c</sup>	4/5	0/5
Manganese	5/5	2.87	132	-	4.22	71.2	--	--	--	--
Mercury	0/5	0.05 <sup>d</sup>	0.05 <sup>d</sup>	0.1	0.05 <sup>d</sup>	0.05 <sup>d</sup>	0.91	1.6	0/5 <sup>e</sup>	0/5 <sup>e</sup>
Molybdenum	4/5	0.05	0.09	0.03	0.05	0.22	--	--	--	--
Nickel	5/5	2.96	5.41	-	2.11	6.41	7.6–162 (52) <sup>c</sup>	69–1,458 (470) <sup>c</sup>	0/5	0/5
Selenium	0/5	0.2 <sup>d</sup>	0.2 <sup>d</sup>	0.3	0.3	0.5	5.0	--	0/5 <sup>e</sup>	--
Silver	0/5	0.005 <sup>d</sup>	0.01 <sup>d</sup>	0.01–0.02	0.04	0.04	--	0.08–38 (3.8) <sup>c</sup>	--	0/5 <sup>e</sup>
Strontium	5/5	10.4	422	-	10.6	27.5	--	--	--	--
Thallium	1/5	0.01	0.01	0.005–0.01	0.04	0.04	--	--	--	--
Tin	1/5	30	30	20	10 <sup>d</sup>	10 <sup>d</sup>	--	--	--	--
Vanadium	5/5	0.24	0.65	-	0.17	2.41	--	--	--	--
Zinc	5/5	6.08	99	-	0.59	5.01	17–373 (120) <sup>c</sup>	17–373 (120) <sup>c</sup>	1/5	1/5

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> U.S. EPA (2002c).

<sup>c</sup> Water quality criteria for this metal is hardness-dependent. The range shown was calculated using the minimum and maximum hardness values for tundra pond surface water. EPA provides a default water quality criteria based on a hardness of 100 mg/L CaCO<sub>3</sub>. This value is presented in parentheses.

<sup>d</sup> Undetected; value listed is one-half of the detection limit.

<sup>e</sup> Undetected in all samples. Undetected values expressed as one-half of the detection limit are below the screening benchmark.

Table 3-24. Ecological screening results for lagoon sediment

Analyte	Detection Frequency	Reference Concentration		Detection Limits (mg/kg)	Ecological Screening Benchmark			Detection Frequency Above Benchmark <sup>a</sup>				
		Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)		Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Effects Range Low <sup>b</sup> (mg/kg)	Effects Range Median <sup>b</sup> (mg/kg)	Marine Sediment Quality Standards <sup>c</sup> (mg/kg)	Effects Range Low	Effects Range Median	Marine Sediment Quality Standards
Aluminum	9/9	2,450	15,300	-	7,440	14,800	--	--	--	--	--	--
Antimony	9/9	0.07	0.27	-	0.1	0.12	--	--	--	--	--	--
Arsenic	9/9	4.9	17.9	-	2.6	4.9	8.2	70	57	2/9	0/9	0/9
Barium	9/9	54.1	350	-	164	271	--	--	--	--	--	--
Cadmium	14/29	0.03	8.1	0.8-5.1	0.18	0.49	1.2	9.6	5.1	6/29	0/29	1/29
Chromium	9/9	4.08	27.2	-	12.5	24.9	81	370	260	0/9	0/9	0/9
Cobalt	9/9	3.85	11.8	-	4.97	9.68	--	--	--	--	--	--
Copper	9/9	3	28.2	-	9.87	18.7	34	270	390	0/9	0/9	0/9
Fluoride	2/9	5.1	8.6	1.9-2.3	1.2 <sup>d</sup>	1.2 <sup>d</sup>	--	--	--	--	--	--
Iron	9/9	10,100	75,000	-	14,000	22,200	--	--	--	--	--	--
Lead	28/29	4.66	302	21	3.2	23	46.7	218	450	8/29	1/29	0/29
Manganese	9/9	97.9	274	-	75.5	129	--	--	--	--	--	--
Mercury	8/9	0.01	0.096	0.008	0.03	0.06	0.15	0.71	0.41	0/9	0/9	0/9
Molybdenum	9/9	0.41	3.39	-	0.46	0.98	--	--	--	--	--	--
Nickel	9/9	12	39	-	18.7	37	20.9	51.6	--	5/9	0/9	--
Selenium	8/9	0.3	2.2	0.2	0.6	1.4	--	--	--	--	--	--
Silver	9/9	0.03	0.27	-	0.08	0.11	1.0	3.7	6.1	0/9	0/9	0/9
Strontium	9/9	10.4	108	-	20.9	40	--	--	--	--	--	--
Thallium	9/9	0.018	0.184	-	0.038	0.103	--	--	--	--	--	--
Tin	1/9	6.7	6.7	1.3-5.3	4.2	5.1	--	--	--	--	--	--
Vanadium	9/9	8.5	35.1	-	16.8	31.5	--	--	--	--	--	--
Zinc	29/29	36	1,590	-	16	370.6	150	410	410	14/29	5/29	5/29

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> Long et al. (1995).

<sup>c</sup> WAC 173-204-320.

<sup>d</sup> Undetected; value listed is one-half of the detection limit.

Table 3-25. Ecological screening results for lagoon surface water

Analyte	Detection Frequency	Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Detection Limits (µg/L)	Reference Concentration		Ecological Screening Benchmark		Detection Frequency Above Benchmark <sup>a</sup>	
					Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Saltwater Criteria Continuous Concentration <sup>b</sup> (µg/L total recoverable)	Saltwater Criterion Maximum Concentration <sup>b</sup> (µg/L total recoverable)	Saltwater Criteria Continuous Concentration	Saltwater Criterion Maximum Concentration
Aluminum	9/9	19.7	247	-	53.5	434	--	--	--	--
Antimony	9/9	0.19	0.63	-	0.11	0.13	--	--	--	--
Arsenic	9/9	4.4	126	-	52.9	98.8	36	69	4/9	3/9
Barium	9/9	111	413	-	144	168	--	--	--	--
Cadmium	12/15	0.05	0.30	0.1	0.18	0.26	8.9	40	0/15	0/15
Chromium	9/9	1.63	4.49	-	5.96	8.22	50	1,100	0/9	0/9
Cobalt	9/9	0.44	1.38	-	3.7	5.35	--	--	--	--
Copper	9/9	0.5	1.4	-	0.4	1.4	3.7	5.8	0/9	0/9
Fluoride	9/9	50	200	-	20	20	--	--	--	--
Iron	9/9	200	723	-	290	693	--	--	--	--
Lead	15/15	0.4	2.3	-	0.1	0.85	8.5	220	0/15	0/15
Manganese	9/9	13.9	277	-	492	801	--	--	--	--
Mercury	0/9	0.05 <sup>c</sup>	0.05 <sup>c</sup>	0.1	0.05 <sup>c</sup>	0.05 <sup>c</sup>	1.1	2.1	0/9 <sup>d</sup>	0/9 <sup>d</sup>
Molybdenum	9/9	0.3	2.41	-	0.07	0.09	--	--	--	--
Nickel	9/9	3.42	10.6	-	9.19	15.2	8.3	75	2/9	0/9
Selenium	6/9	0.3	0.6	0.3	0.2 <sup>c</sup>	0.2 <sup>c</sup>	71	290	0/9	0/9
Silver	8/9	0.01	0.25	0.1	0.02	0.03	--	2.2	--	0/9
Strontium	9/9	503	1,850	-	991	1,470	--	--	--	--
Thallium	5/9	0.007	0.07	0.025–0.05	0.006	0.009	--	--	--	--
Tin	1/9	23.7	23.7	20	10 <sup>c</sup>	10 <sup>c</sup>	--	--	--	--
Vanadium	6/9	0.22	0.85	0.35–0.7	0.4 <sup>c</sup>	0.4 <sup>c</sup>	--	--	--	--
Zinc	15/15	3.09	110	-	17	30.1	86	95	1/15	1/15

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> U.S. EPA (2002c).

<sup>c</sup> Undetected; value listed is one-half of the detection limit.

<sup>d</sup> Undetected in all samples. Undetected values expressed as one-half of the detection limit are below the screening benchmark.



Table 3-26. Ecological screening results for marine sediment

Analyte	Detection Frequency	Reference Concentration			Ecological Screening Benchmark				Detection Frequency Above Benchmark <sup>a</sup>			
		Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Detection Limits (mg/kg)	Minimum Detected Concentration (mg/kg)	Maximum Detected Concentration (mg/kg)	Effects Range Low <sup>b</sup> (mg/kg)	Effects Range Median <sup>b</sup> (mg/kg)	Marine Sediment Quality Standards <sup>c</sup> (mg/kg)	Effects Range Low	Effects Range Median	Marine Sediment Quality Standards
Aluminum	20/20	1,990	6,070	-	1,970	8,000	--	--	--	--	--	--
Antimony	16/43	0.06	4.59	0.08–14	0.07	0.13	--	--	--	--	--	--
Arsenic	70/71	3.1	14.5	27	5.6	13	8.2	70	57	13/71	0/71	0/71
Barium	71/71	79.5	639	-	22	431	--	--	--	--	--	--
Cadmium	107/136	0.04	52.9	0.03–1.5	0.02	0.23	1.2	9.6	5.1	24/136	1/136	1/136
Chromium	71/71	2.4	33.5	-	1.4	18	81	370	260	0/71	0/71	0/71
Cobalt	20/20	3.15	8.89	-	4.2	8.71	--	--	--	--	--	--
Copper	71/71	3.7	34.8	-	3	10.2	34	270	390	1/71	0/71	0/71
Fluoride	18/18	0.4	1.5	-	0.4	2	--	--	--	--	--	--
Iron	20/20	9,960	19,300	-	8,150	22,700	--	--	--	--	--	--
Lead	136/136	1.59	5,620	-	2.7	11.2	46.7	218	450	5/136	1/136	1/136
Manganese	20/20	161	363	-	187	389	--	--	--	--	--	--
Mercury	35/69	0.01	0.58	0.01–0.2	0.01	0.02	0.15	0.71	0.41	1/69	0/69	1/69
Molybdenum	19/20	0.37	1.04	2.7	0.44	0.83	--	--	--	--	--	--
Nickel	20/20	11.3	33.6	-	9.8	34.8	20.9	51.6	--	15/20	0/20	--
Selenium	26/71	0.3	6	1–27	0.3	1.2	--	--	--	--	--	--
Silver	42/71	0.03	2.11	0.05–2.9	0.02	0.49	1.0	3.7	6.1	1/71	0/71	0/71
Strontium	19/19	24.4	33.8	-	13	29	--	--	--	--	--	--
Thallium	19/19	0.023	1.13	-	0.025	0.052	--	--	--	--	--	--
Tin	5/19	5.6	8.9	1.9–5	4.8	9.5	--	--	--	--	--	--
Vanadium	43/43	9.11	46	-	13	33.9	--	--	--	--	--	--
Zinc	136/136	5.5	2,550	-	25	56.8	150	410	410	7/136	3/136	3/136

**Note:** Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> Long et al. (1995).

<sup>c</sup> WAC 173-204-320.

Table 3-27. Ecological screening results for marine surface water

Analyte	Detection Frequency	Reference Concentration			Ecological Screening Benchmark		Detection Frequency Above Benchmark <sup>a</sup>			
		Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Detection Limits (µg/L)	Minimum Detected Concentration (µg/L unfiltered)	Maximum Detected Concentration (µg/L unfiltered)	Saltwater Criteria Continuous Concentration <sup>b</sup> (µg/L total recoverable)	Saltwater Criterion Maximum Concentration <sup>b</sup> (µg/L total recoverable)	Saltwater Criteria Continuous Concentration	Saltwater Criterion Maximum Concentration
Aluminum	9/11	44	215	50	63.9	336	--	--	--	--
Antimony	6/11	0.47	2.79	0.4–0.8	0.44	1.67	--	--	--	--
Arsenic	8/11	1.5	6.3	5	1.1	7.5	36	69	0/11	0/11
Barium	11/11	12.1	39.4	-	9.91	38.1	--	--	--	--
Cadmium	11/11	1.58	4.62	-	2.27	4.69	8.9	40	0/11	0/11
Chromium	0/11	1 <sup>c</sup>	2 <sup>c</sup>	2-3	1 <sup>c</sup>	2 <sup>c</sup>	50	1,100	0/11 <sup>d</sup>	0/11 <sup>d</sup>
Cobalt	11/11	3.85	4.6	-	4.03	4.48	--	--	--	--
Copper	5/11	1	5.9	8	1	2.6	3.7	5.8	1/11	1/11
Fluoride	11/11	500	900	-	600	800	--	--	--	--
Iron	11/11	52.3	375	-	33.6	643	--	--	--	--
Lead	11/11	0.7	1.47	-	0.76	1.25	8.5	220	0/11	0/11
Manganese	11/11	13.1	31.9	-	10.1	25.5	--	--	--	--
Mercury	0/11	0.05 <sup>c</sup>	0.05 <sup>c</sup>	0.1	0.05 <sup>c</sup>	0.05 <sup>c</sup>	1.1	2.1	0/11 <sup>d</sup>	0/11 <sup>d</sup>
Molybdenum	11/11	8.1	11.1	-	8.26	10.6	--	--	--	--
Nickel	0/11	2 <sup>c</sup>	10 <sup>c</sup>	3–20	2 <sup>c</sup>	10 <sup>c</sup>	8.3	75	0/11 <sup>e</sup>	0/11 <sup>e</sup>
Selenium	9/11	0.3	1	0.3	0.3	0.5	71	290	0/11	0/11
Silver	5/11	0.51	0.95	0.2–0.4	0.27	0.27	--	2.2	--	0/11
Strontium	11/11	4,310	5,650	-	4,530	5,290	--	--	--	--
Thallium	1/11	0.09	0.09	0.1–0.2	0.111	0.133	--	--	--	--
Tin	1/11	23.3	23.3	6-20	26.4	26.4	--	--	--	--
Vanadium	2/11	4.44	5.27	2.8–4	3.77	8.44	--	--	--	--
Zinc	0/11	1 <sup>c</sup>	1 <sup>c</sup>	2	1 <sup>c</sup>	1 <sup>c</sup>	86	95	0/11 <sup>d</sup>	0/11 <sup>d</sup>

Note: Field duplicates were screened separately.

- - detected in all samples
- - no benchmark

<sup>a</sup> Expressed as the ratio of the detected exceedances over the total analyses.

<sup>b</sup> U.S. EPA (2002c).

<sup>c</sup> Undetected; value listed is one-half of the detection limit.

<sup>d</sup> Undetected in all samples. Undetected values expressed as one-half of the detection limit are below the screening benchmark.

<sup>e</sup> Undetected in all samples. Five undetected values expressed as one-half of the detection limit exceed the benchmark.

Table 3-28. Toxicity reference values used for wildlife screening and CoPC selection

CoPC	TRVs (mg/kg-day)					
	Avian			Mammalian		
	NOAEL	LOAEL	Citation	NOAEL	LOAEL	Citation
Aluminum	120	NA	Carriere et al. (1986)	1.9	19	Ondreicka et al. (1966)
Antimony	NA	NA	NA	0.66	NA	Schroeder et al. (1970)
Arsenic <sup>a</sup>	10	40	Stanley et al. (1994)	0.13	1.3	Schroeder and Mitchener (1971)
Barium	21	42	Johnson et al. (1960)	5.1	20	Perry et al. (1983); Borzelleca et al. (1988)
Cadmium	1.5	20	White and Finley (1978)	1.0	10	Sutou et al. (1980)
Chromium	0.86	4.3	Haseltine et al. (1985) as cited in Sample et al. (1996)	3.3	69	Mackenzie et al. (1958); Gross and Heller (1946)
Cobalt	NA	NA	NA	0.5	2.0	Nation et al. (1983)
Copper	47	62	Mehring et al. (1960)	12	15	Aulerich et al. (1982)
Fluoride	7.8	32	Pattee et al. (1988)	31	53	Aulerich et al. (1987)
Iron	NA	NA	NA	NA	NA	NA
Lead	3.9	NA	Pattee (1984)	11	90	Azar et al. (1973)
Manganese	980	NA	Laskey and Edens (1985)	88	280	Laskey et al. (1982)
Mercury <sup>b</sup>	0.032	0.064	Heinz (1974, 1976a,b, 1979)	0.032	0.16	Verschuuren et al. (1976)
Molybdenum	3.5	35	Lepore and Miller (1965)	0.26	2.6	Schroeder and Mitchener (1971)
Nickel	77	110	Cain and Pafford (1981)	40	80	Ambrose et al. (1976)
Selenium	0.40	0.80	Heinz et al. (1989)	0.20	0.33	Rosenfeld and Beath (1954)
Silver	NA	NA	NA	NA	NA	NA
Strontium	NA	NA	NA	263	NA	Skoryna (1981)
Thallium	0.24	24	Hudson et al. (1984)	0.074	0.74	Formigli et al. (1986)
Tin <sup>c</sup>	6.8	17	Schlatterer et al. (1993)	23	35	Davis et al. (1987)
Vanadium	11	NA	White and Dieter (1978)	0.21	2.1	Domingo et al. (1986)
Zinc	130	NA	Stahl et al. (1990)	160	320	Schlicker and Cox (1968)

**Note:** CoPC - chemical of potential concern  
 LOAEL - lowest-observed-adverse-effect level  
 NA - not available; no suitable TRV was derived  
 NOAEL - no-observed-adverse-effect level  
 TRV - toxicity reference value

<sup>a</sup> Avian TRVs were based on exposure to arsenic as arsenate; mammalian TRVs were based on exposure to arsenic as arsenite.

<sup>b</sup> Mercury TRVs were based on exposure to methylmercury.

<sup>c</sup> Tin TRVs were based on exposure to tributyltin.

**Table 3-29. Ecological exposure assumptions for use in screening food-web models**

Representative Receptor	Body Weight (kg)	Food Ingestion Rate (kg/day (dry wt))	Soil/Sediment Ingestion Rate (kg/day (dry wt))	Area Use Factor
Tundra vole	0.029 <sup>a</sup>	0.0060 <sup>b</sup>	0.00014 <sup>c</sup>	1
Common snipe	0.081 <sup>d</sup>	0.012 <sup>e</sup>	0.0012 <sup>f</sup>	1
Red-throated loon	1.15 <sup>g</sup>	0.079 <sup>h</sup>	0.0016 <sup>i</sup>	1
River otter	6.8 <sup>j</sup>	0.19 <sup>k</sup>	0.018 <sup>l</sup>	1
Black-bellied plover	0.19 <sup>m</sup>	0.026 <sup>n</sup>	0.0075 <sup>o</sup>	1

<sup>a</sup> Minimum female body weight from Bee and Hall (1956).

<sup>b</sup> Based on Nagy et al. (1999) allometric equation for Rodentia.

<sup>c</sup> Based on 2.4 percent soil in meadow vole diet from Beyer et al. (1994).

<sup>d</sup> Minimum female body weight from Tuck (1972).

<sup>e</sup> Based on Nagy et al. (1999) allometric equation for insectivores.

<sup>f</sup> Based on 10.4 percent soil in American woodcock diet from Beyer et al. (1994).

<sup>g</sup> Minimum body weight from Dunning (1993).

<sup>h</sup> Based on Nagy et al. (1999) allometric equation for all birds.

<sup>i</sup> Based on minimum soil ingestion rate from Beyer et al. (1994).

<sup>j</sup> Minimum body weight from DFG (2002b).

<sup>k</sup> Based on Nagy et al. (1999) allometric equation for Carnivora.

<sup>l</sup> Based on 9.4 percent soil in raccoon diet from Beyer et al. (1994).

<sup>m</sup> Minimum female body weight for Alaska from Paulson (1995).

<sup>n</sup> Based on Nagy et al. (1999) allometric equation for Charadriiformes.

<sup>o</sup> Based on 29 percent sediment in black-bellied plover diet from Hui and Beyer (1998).

Table 3-30. Screening-level food-web results for tundra vole

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake (mg/day)	Body Weight Normalized Exposure		TRV (mg/kg-day)	Hazard Quotient
	Tundra Soil (mg/kg dw)	Moss (mg/kg dw)	Tundra Soil (mg/day)	Moss (mg/day)		(mg/kg-day)	(mg/kg-day)		
<b>Site</b>									
<b>Metals</b>									
Aluminum	18,900	47,900	2.73	289	291	10,000	1.9		5,300
Antimony	25.8	4.58	0.00373	0.0276	0.0313	1.08	0.66		1.6
Arsenic	150	15.7	0.0217	0.0946	0.116	4.01	0.13		31
Barium	5,810	8,800	0.840	53.0	53.9	1,857	5.1		360
Cadmium	438	48.4	0.0633	0.292	0.355	12.2	1.0		12
Chromium	33.2	32.9	0.00480	0.198	0.203	7.00	3.3		2.1
Cobalt	35	9.35	0.00506	0.0563	0.0614	2.12	0.50		4.2
Copper	58.3	40.5	0.00843	0.244	0.252	8.71	12		0.73
Fluoride	3.8	NA	0.000550	NA	0.000550	0.0189	31		0.000611
Lead	16,000	1,720	2.31	10.4	12.7	437	11		40
Manganese	3,400	842	0.492	5.07	5.57	192	88		2.2
Mercury	4.16	1.04	0.000602	0.00627	0.00687	0.237	0.032		7.4
Molybdenum	3.9	2.4	0.000564	0.0145	0.0150	0.518	0.26		2.0
Nickel	37.5	31.6	0.00542	0.190	0.196	6.75	40		0.17
Selenium	3	1.5	0.000477	0.00904	0.00952	0.328	0.20		1.6
Strontium	150	107	0.0217	0.645	0.666	23.0	260		0.088
Thallium	1.58	1.84	0.000228	0.0111	0.0113	0.390	0.074		5.3
Tin	14	3.9	0.00202	0.0235	0.0255	0.880	23		0.038
Vanadium	46.5	14.7	0.00672	0.0886	0.0953	3.29	0.21		16
Zinc	82,700	8,120	12.0	48.9	60.9	2,100	160		13
<b>Reference</b>									
<b>Metals</b>									
Aluminum	11,300	713	1.63	4.30	5.93	204	1.9		110
Antimony	0.28	0.15	0.0000405	0.000904	0.000944	0.0326	0.66		0.049
Arsenic	6.8	0.3	0.000983	0.00181	0.00279	0.0962	0.13		0.74
Barium	624	119	0.0902	0.717	0.807	27.8	5.1		5.5
Cadmium	0.88	0.38	0.000127	0.00229	0.00242	0.0833	1.0		0.083
Chromium	19.7	2.96	0.00285	0.0178	0.0207	0.713	3.3		0.22
Cobalt	28.3	2.03	0.00409	0.0122	0.0163	0.563	0.50		1.1
Copper	16.9	4.35	0.00244	0.0262	0.0287	0.988	12		0.082
Fluoride	0.4	NA	0.0000578	NA	0.0000578	0.00199	31		0.000064
Lead	23.3	9.64	0.00337	0.0581	0.0615	2.12	11		0.19
Manganese	6,620	712	0.957	4.29	5.25	181	88		2.1
Mercury	0.15	0.067	0.0000217	0.000404	0.000425	0.0147	0.032		0.46
Molybdenum	2.27	0.3	0.000328	0.00181	0.00214	0.0737	0.26		0.28
Nickel	36.8	6.34	0.00532	0.0382	0.0435	1.50	40		0.038
Selenium	1	0.1	0.000145	0.000603	0.000747	0.0258	0.20		0.13
Strontium	39.6	11	0.00573	0.0663	0.0720	2.48	260		0.0096
Thallium	0.116	0.04	0.0000168	0.000241	0.000258	0.00889	0.074		0.12
Tin	17.4	1.1	0.00252	0.00663	0.00914	0.315	23		0.014
Vanadium	24.7	1.73	0.00357	0.0104	0.0140	0.483	0.21		2.3
Zinc	111	64	0.0161	0.386	0.402	13.9	160		0.087

Note: Hazard quotients greater than 1.0 are boxed.

NA - not available

TRV - toxicity reference value

Table 3-31. Screening-level food-web results for river otter

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight Normalized Exposure	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Fish (mg/kg dw)	Sediment (mg/day)	Fish (mg/day)				
<b>Aufeis Creek</b>								
<b>Metals</b>								
Cadmium	0.31	0.17	0.00558	0.0325	0.0382	0.00562	1.0	0.0056
Lead	14.9	6	0.268	1.15	1.42	0.208	11	0.019
Selenium	2.5	7	0.0450	1.34	1.38	0.203	0.20	1.0
Zinc	136	121	2.45	23.1	25.7	3.77	160	0.024
<b>Omikviorok River</b>								
<b>Metals</b>								
Cadmium	0.59	0.14	0.0106	0.0268	0.0375	0.00551	1.0	0.0055
Lead	19	3.03	0.342	0.579	0.921	0.136	11	0.012
Selenium	0.7	3.4	0.0126	0.650	0.662	0.0974	0.20	0.49
Zinc	123	155	2.21	29.6	31.9	4.69	160	0.029
<b>Anxiety Ridge Creek</b>								
<b>Metals</b>								
Cadmium	1.38	0.39	0.0248	0.0745	0.0994	0.0146	1.0	0.015
Lead	142	2.86	2.56	0.547	3.10	0.456	11	0.041
Selenium	2.4	5.87	0.0432	1.12	1.17	0.171	0.20	0.86
Zinc	259	155 <sup>a</sup>	4.66	29.6	34.3	5.04	160	0.032

**Note:** Hazard quotients greater than 1.0 are boxed.

TRV - toxicity reference value

<sup>a</sup> No zinc tissue data available for Anxiety Ridge Creek; maximum concentration from Omikviorok River used in calculation.

Table 3-32. Screening-level food-web results for red-throated loon

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight Normalized Exposure	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Fish (mg/kg dw)	Sediment (mg/day)	Fish (mg/day)				
<b>Aufeis Creek</b>								
<b>Metals</b>								
Cadmium	0.31	0.17	0.000488	0.0134	0.0139	0.0121	1.5	0.0080
Lead	14.9	6	0.0235	0.472	0.496	0.431	3.9	0.11
Selenium	2.5	7	0.00394	0.551	0.555	0.483	0.40	1.2
Zinc	136	121	0.214	9.52	9.74	8.47	130	0.065
<b>Omikviorok River</b>								
<b>Metals</b>								
Cadmium	0.59	0.14	0.000929	0.0110	0.0120	0.0104	1.5	0.0069
Lead	19	3.03	0.0299	0.238	0.269	0.234	3.9	0.060
Selenium	0.7	3.4	0.00110	0.268	0.269	0.234	0.40	0.58
Zinc	123	155	0.194	12.2	12.4	10.8	130	0.083
<b>Anxiety Ridge Creek</b>								
<b>Metals</b>								
Cadmium	1.38	0.39	0.00217	0.0307	0.0329	0.0286	1.5	0.019
Lead	142	2.86	0.224	0.225	0.449	0.390	3.9	0.10
Selenium	2.4	5.87	0.00378	0.462	0.466	0.405	0.40	1.0
Zinc	259	155 <sup>a</sup>	0.408	12.2	12.6	11.0	130	0.084

**Note:** Hazard quotients greater than 1.0 are boxed.

TRV - toxicity reference value

<sup>a</sup> No zinc tissue data available for Anxiety Ridge Creek; maximum concentration from Omikviorok River used in calculation.

Table 3-33. Screening-level food-web results for common snipe foraging in freshwater rivers and creeks

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Invertebrates (mg/kg dw)	Sediment (mg/day)	Invertebrates (mg/day)		Normalized Exposure		
<b>New Heart Creek</b>								
<b>Metals</b>								
Aluminum	17,100	17,100	20.5	205	226	2,790	120	23
Arsenic	7.3	5.0	0.0088	0.0604	0.0692	0.85	10	0.09
Barium	293	293	0.352	3.52	3.87	47.7	21	2.3
Cadmium	0.77	6.15	0.000924	0.0738	0.0748	0.923	1.5	0.62
Chromium	15.6	7.30	0.0187	0.0876	0.106	1.31	0.86	1.5
Copper	14.2	74.6	0.0170	0.895	0.912	11.3	47	0.24
Fluoride	1.2	1.2	0.00144	0.0144	0.0158	0.196	7.8	0.025
Lead	23.8	14.4	0.0286	0.173	0.202	2.49	3.9	0.64
Manganese	939	939	1.13	11.3	12.4	153	980	0.16
Mercury	0.06	0.172	0.0000720	0.002100	0.00210	0.0264	0.032	0.82
Molybdenum	0.84	0.84	0.00101	0.0101	0.0111	0.137	3.5	0.039
Nickel	45.2	105	0.0542	1.258	1.313	16.2	77	0.21
Selenium	1.4	1.4	0.00168	0.0168	0.0185	0.228	0.4	0.57
Thallium	0.08	0.08	0.0000960	0.000960	0.00106	0.0130	0.24	0.055
Tin	7.6	7.6	0.00912	0.0912	0.100	1.24	6.8	0.18
Vanadium	13.8	13.8	0.0166	0.166	0.182	2.25	11	0.20
Zinc	206	1,551	0.247	18.6	18.9	233	130	1.8
<b>Aufeis Creek</b>								
<b>Metals</b>								
Aluminum	7,580	7,580	9.10	91.0	100	1,240	120	10
Arsenic	9.6	6.62	0.0115	0.079	0.91	1.12	10	0.11
Barium	172	172	0.206	2.06	2.27	28.0	21	1.3
Cadmium	0.31	2.48	0.000372	0.0297	0.0301	0.372	1.5	0.25
Chromium	22.1	10.3	0.0265	0.124	0.151	1.86	0.86	2.2
Copper	28.2	148	0.0338	1.777	1.810	22.4	47	0.48
Fluoride	1.2	1.2	0.00144	0.0144	0.0158	0.196	7.8	0.025
Lead	14.9	9.04	0.0179	0.109	0.126	1.56	3.9	0.40
Manganese	1,200	1,200	1.44	14.4	15.8	196	980	0.20
Mercury	0.06	0.17	0.0000720	0.00206	0.00214	0.0264	0.032	0.82
Molybdenum	1.01	1.01	0.00121	0.0121	0.0133	0.165	3.5	0.047
Nickel	107.9	108	0.0558	1.3	1.35	16.7	77	0.22
Selenium	2.5	2.5	0.00300	0.0300	0.0330	0.407	0.40	1.0
Thallium	0.115	0.115	0.000138	0.00138	0.00152	0.0187	0.24	0.079
Tin	6.4	6.4	0.00768	0.0768	0.0845	1.04	6.8	0.15
Vanadium	18.2	18.2	0.0218	0.218	0.240	2.97	11	0.27
Zinc	136	1,024	0.163	12.3	12.4	154	130	1.2



Table 3-33. (cont.)

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Invertebrates (mg/kg dw)	Sediment (mg/day)	Invertebrates (mg/day)		Normalized Exposure		
<b>Omikviorok River</b>								
<b>Metals</b>								
Aluminum	14,900	14,900	17.9	179	197	2,430	120	20
Arsenic	9.4	6.49	0.0113	0.0778	0.089	1.10	10	0.11
Barium	484	484	0.581	5.81	6.39	78.9	21	3.8
Cadmium	0.59	4.71	0.000708	0.0566	0.0573	0.7071	1.5	0.47
Chromium	23.3	10.9	0.0280	0.131	0.159	1.96	0.86	2.3
Copper	15.6	81.9	0.0187	0.983	1.002	12.36	47	0.26
Fluoride	1.2	1.2	0.00144	0.0144	0.0158	0.196	7.8	0.025
Lead	19	11.5	0.0228	0.138	0.161	1.99	3.9	0.51
Manganese	2,140	2,140	2.57	25.7	28.2	349	980	0.36
Mercury	0.06	0.172	0.0000720	0.00206	0.00214	0.0264	0.032	0.82
Molybdenum	0.66	0.66	0.000792	0.00792	0.00871	0.108	3.5	0.031
Nickel	57.3	133	0.0688	0.0688	1.60	20.5	77	0.27
Selenium	0.7	0.7	0.000840	0.00840	0.00924	0.114	0.40	0.29
Thallium	0.141	0.141	0.000169	0.00169	0.00186	0.0230	0.24	0.097
Tin	5.7	5.7	0.00684	0.0684	0.0752	0.929	6.8	0.14
Vanadium	28.1	28.1	0.0337	0.337	0.371	4.58	11	0.42
Zinc	123	926	0.148	11.1	11.3	139	130	1.1
<b>Anxiety Ridge Creek</b>								
<b>Metals</b>								
Aluminum	8,310	8,310	9.97	99.7	110	1,354	120	11
Arsenic	11.4	7.87	0.0137	0.0944	0.108	1.33	10	0.13
Barium	922	922	1.11	11.1	12.2	150	21	7.2
Cadmium	1.38	11.0	0.00166	0.132	0.134	1.654	1.5	1.1
Chromium	14.6	6.83	0.0175	0.082	0.100	1.23	0.86	1.4
Copper	20.1	106	0.0241	1.27	1.29	15.9	47	0.339
Fluoride	0.95	0.95	0.00114	0.0114	0.0125	0.155	7.8	0.020
Lead	142	86.2	0.170	1.03	1.20	14.9	3.9	3.8
Manganese	2,100	2,100	2.52	25.2	27.7	342	980	0.35
Mercury	0.089	0.255	0.000107	0.00316	0.00317	0.0391	0.032	1.2
Molybdenum	2.32	2.32	0.00278	0.0278	0.0306	0.378	3.5	0.11
Nickel	45.6	106	0.0547	1.27	1.32	16.3	77	0.21
Selenium	2.4	2.4	0.00288	0.0288	0.032	0.391	0.40	0.98
Thallium	0.322	0.322	0.000386	0.00386	0.00425	0.0525	0.24	0.22
Tin	1.1	1.1	0.00132	0.0132	0.0145	0.179	6.8	0.026
Vanadium	20.5	20.5	0.0246	0.246	0.271	3.34	11	0.30
Zinc	259	1,949	0.311	23.4	23.7	293	130	2.3

Table 3-33. (cont.)

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Invertebrates (mg/kg dw)	Sediment (mg/day)	Invertebrates (mg/day)		Normalized Exposure		
<b>Reference Creeks</b>								
<b>Metals</b>								
Aluminum	12,100	12,100	14.5	145	160	1,970	120	16
Arsenic	8.1	5.59	0.00972	0.0671	0.0768	0.948	10	0.095
Barium	483	483	0.580	5.80	6.38	78.7	21	3.7
Cadmium	0.3	2.40	0.000360	0.0288	0.0291	0.360	1.5	0.24
Chromium	19.9	9.3	0.0239	0.112	0.136	1.67	0.86	1.9
Copper	18.5	97.1	0.0222	1.17	1.19	14.7	47	0.31
Fluoride	1.2	1.2	0.00144	0.0144	0.0158	0.196	7.8	0.025
Lead	9.17	5.57	0.011	0.0668	0.0778	0.961	3.9	0.25
Manganese	859	859	1.03	10.3	11.3	140	980	0.14
Mercury	0.04	0.115	0.000480	0.00138	0.00142	0.0176	0.032	0.55
Molybdenum	0.52	0.52	0.000624	0.00624	0.00686	0.0847	3.5	0.024
Nickel	35	81.2	0.0420	0.974	1.02	12.5	77	0.16
Selenium	0.7	0.7	0.000840	0.00840	0.00924	0.114	0.40	0.29
Thallium	0.07	0.07	0.000840	0.000840	0.000924	0.0114	0.24	0.048
Tin	2.4	2.4	0.00288	0.0288	0.0317	0.391	6.8	0.058
Vanadium	24.8	24.8	0.0298	0.298	0.327	4.04	11	0.37
Zinc	69.7	525	0.0836	6.30	6.38	78.8	130	0.606

**Note:** Hazard quotients greater than 1.0 are boxed.

Invertebrate data are modeled based on maximum sediment concentrations and are not measured values.

TRV - toxicity reference value

Table 3-34. Screening-level food web results for common snipe foraging in tundra ponds

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight Normalized Exposure	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Invertebrates (mg/kg dw)	Sediment (mg/day)	Invertebrates (mg/day)				
<b>Site Tundra Ponds</b>								
<b>Metals</b>								
Aluminum	5,270	5,270	6.32	63.2	69.6	859	120	7.2
Arsenic	8.7	6.00	0.0104	0.0720	0.0825	1.02	10	0.10
Barium	626	626	0.751	7.51	8.26	102	21	4.9
Cadmium	119	951	0.143	11.4	11.6	143	1.5	95
Chromium	15.3	7.16	0.0184	0.0859	0.104	1.29	0.86	1.5
Copper	53.4	280	0.0641	3.36	3.43	42.3	47	0.90
Fluoride	4.4	4.4	0.00528	0.0528	0.0581	0.717	7.8	0.092
Lead	2,180	1,323	2.62	15.9	18.5	228	3.9	59
Manganese	745	745	0.894	8.94	9.83	121	980	0.12
Mercury	1.31	3.76	0.00157	0.0451	0.0467	0.576	0.032	18
Molybdenum	2.84	2.84	0.00341	0.0341	0.0375	0.463	3.5	0.13
Nickel	44.2	103	0.0530	1.23	1.28	15.8	77	0.21
Selenium	3.5	3.5	0.00420	0.0420	0.0462	0.570	0.40	1.4
Thallium	1.92	1.92	0.00230	0.0230	0.0253	0.313	0.24	1.3
Tin	41.2	41.2	0.0494	0.494	0.544	6.71	6.8	0.99
Vanadium	28.3	28.3	0.0340	0.340	0.374	4.61	11	0.42
Zinc	27,000	203,229	32.4	2,439	2,471	30,508	130	235
<b>Reference Ponds</b>								
<b>Metals</b>								
Aluminum	17,100	17,100	20.5	205	226	2,790	120	23
Arsenic	13	8.97	0.0156	0.108	0.123	1.52	10	0.15
Barium	772	772	0.926	9.26	10.2	126	21	6.0
Cadmium	0.66	5.27	0.000792	0.0633	0.0641	0.791	1.5	0.53
Chromium	28	13	0.0336	0.157	0.191	2.36	0.86	2.7
Copper	20.7	109	0.0248	1.30	1.33	16.4	47	0.349
Fluoride	1.2	1.2	0.00144	0.0144	0.0158	0.196	7.8	0.025
Lead	20.3	12.3	0.0244	0.148	0.172	2.13	3.9	0.55
Manganese	1,870	1,870	2.24	22.4	24.7	305	980	0.31
Mercury	0.07	0.20	0.0000840	0.00241	0.00249	0.0308	0.032	0.96
Molybdenum	1.35	1.35	0.00162	0.0162	0.0178	0.220	3.5	0.063
Nickel	163	163	0.0844	1.96	2.04	25.3	77	0.33
Selenium	3.1	3.10	0.00372	0.0372	0.0409	0.505	0.40	1.3
Thallium	0.174	0.174	0.000209	0.00209	0.00230	0.0284	0.24	0.12
Tin	6.3	6.3	0.00756	0.0756	0.0832	1.03	6.8	0.15
Vanadium	94.5	94.5	0.113	1.13	1.25	15.4	11	1.4
Zinc	138	1,039	0.166	12.5	12.6	160	130	1.2

**Note:** Hazard quotients greater than 1.0 are boxed.

Invertebrate data are modeled based on maximum sediment concentrations and are not measured values.

TRV - toxicity reference value

Table 3-35. Screening-level food-web results for black-bellied plover foraging in coastal lagoons

Analyte	Maximum Concentration		Daily Exposure		Total Daily Intake	Body Weight Normalized Exposure	TRV (mg/kg-day)	Hazard Quotient
	Sediment (mg/kg dw)	Invertebrates (mg/kg dw)	Sediment (mg/day)	Invertebrates (mg/day)				
<b>Site Coastal Lagoons</b>								
<b>Metals</b>								
Aluminum	15,300	15,300	115	398	513	2,660	120	22
Arsenic	17.9	12.4	0.135	0.321	0.456	2.36	10	0.24
Barium	350	350	2.64	9.09	11.7	60.8	21	2.9
Cadmium	8.1	64.7	0.061	1.68	1.74	9.03	1.5	6.02
Chromium	27.2	12.7	0.205	0.331	0.536	2.78	0.86	3.2
Copper	28.2	148	0.212	3.85	4.06	21.0	47	0.45
Fluoride	8.6	8.6	0.0648	0.223	0.288	1.49	7.8	0.19
Lead	302	183	2.28	4.76	7.0	36.5	3.9	9.4
Manganese	274	274	2.06	7.12	9.18	47.6	980	0.049
Mercury	0.096	0.275	0.000723	0.00715	0.00788	0.0408	0.032	1.3
Molybdenum	3.39	3.39	0.0255	0.0881	0.114	0.589	3.5	0.17
Nickel	39	90	0.294	2.35	2.64	13.7	77	0.18
Selenium	2.2	2.2	0.0166	0.0572	0.0737	0.382	0.40	0.96
Thallium	0.184	0.184	0.00139	0.00478	0.00617	0.0320	0.24	0.13
Tin	6.7	6.7	0.0505	0.174	0.225	1.16	6.8	0.17
Vanadium	35.1	35.1	0.264	0.912	1.18	6.10	11	0.55
Zinc	1,590	11,968	12.0	311	323	1,673	130	13
<b>Reference Lagoons</b>								
<b>Metals</b>								
Aluminum	14,800	14,800	112	385	496	2,570	120	21
Arsenic	4.9	3.38	0.0369	0.0879	0.125	0.647	10	0.065
Barium	271	271	2.04	7.04	9.08	47.1	21	2.2
Cadmium	0.49	3.92	0.00369	0.102	0.105	0.546	1.5	0.364
Chromium	24.9	11.7	0.188	0.303	0.490	2.54	0.86	3.0
Copper	18.7	98.2	0.141	2.55	2.69	13.9	47	0.30
Fluoride	1.2	1.2	0.00904	0.0312	0.0402	0.208	7.8	0.027
Lead	23	14	0.173	0.363	0.536	2.78	3.9	0.71
Manganese	129	129	0.972	3.35	4.32	22.4	980	0.023
Mercury	0.06	0.17	0.000452	0.00447	0.00492	0.0255	0.032	0.80
Molybdenum	0.98	0.98	0.00738	0.0255	0.0328	0.170	3.5	0.049
Nickel	37	86	0.279	2.230	2.51	13.0	77	0.169
Selenium	1.4	1.4	0.0105	0.0364	0.0469	0.243	0.40	0.61
Thallium	0.103	0.103	0.000776	0.00268	0.00345	0.0179	0.24	0.075
Tin	5.1	5.1	0.0384	0.133	0.171	0.886	6.8	0.13
Vanadium	31.5	31.5	0.237	0.818	1.06	5.47	11	0.50
Zinc	371	2,790	2.79	72.5	75.3	390	130	3.00

Note: Hazard quotients greater than 1.0 are boxed.

Invertebrate data are modeled based on maximum sediment concentrations and are not measured values.

TRV - toxicity reference value

Table 3-36. Results of screening against lowest ecological screening benchmarks

Chemical	Environment								
	Terrestrial	Streams		Ponds		Lagoons		Marine	
	Tundra Soil	Sediment	Water	Sediment	Water	Sediment	Water	Sediment	Water
Aluminum	Fail	Pass	Fail	Pass	Fail	NB	NB	NB	NB
Antimony	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Arsenic	Fail	Fail	ND <sup>a</sup>	Pass	Pass	Fail	Fail	Fail	Pass
Barium	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Cadmium	Fail	Fail	Fail	Fail	Fail	Fail	Pass	Fail	Pass
Chromium	Fail	Pass	ND <sup>a</sup>	Pass	Pass	Pass	Pass	Pass	ND <sup>a</sup>
Cobalt	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Copper	Fail	Pass	Pass	Fail	Fail	Pass	Pass	Fail	Fail
Fluoride	Pass	ND <sup>b</sup>	NB	NB	NB	NB	NB	NB	NB
Iron	Fail	Pass	Fail	Pass	Fail	NB	NB	NB	NB
Lead	Fail	Fail	Fail	Fail	Fail	Fail	Pass	Fail	Pass
Manganese	Fail	Pass	NB	Pass	NB	NB	NB	NB	NB
Mercury	Fail	Pass	ND <sup>a</sup>	Fail	ND <sup>a</sup>	Pass	ND <sup>a</sup>	Fail	ND <sup>a</sup>
Molybdenum	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Nickel	Fail	Fail	Pass	Fail	Pass	Fail	Fail	Fail	ND <sup>c</sup>
Selenium	Fail	NB	Pass	NB	ND <sup>a</sup>	NB	Pass	NB	Pass
Silver	Fail	NB	ND <sup>a</sup>	NB	ND <sup>a</sup>	Pass	Pass	Fail	Pass
Strontium	NB	NB	NB	NB	NB	NB	NB	NB	NB
Thallium	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Tin	Pass	NB	NB	NB	NB	NB	NB	NB	NB
Vanadium	Fail	NB	NB	NB	NB	NB	NB	NB	NB
Zinc	Fail	Fail	Fail	Fail	Fail	Fail	Fail	Fail	ND <sup>a</sup>

**Note:** Fail - maximum detected concentration exceeds the lowest benchmark  
 NB - no benchmark  
 ND - undetected in all samples  
 Pass - maximum detected concentration is below the lowest benchmark

<sup>a</sup> Maximum value expressed as one-half of the detection limit is below the screening benchmark.

<sup>b</sup> No benchmark.

<sup>c</sup> Maximum value expressed as one-half of the detection limit is above the screening benchmark.

Table 3-37. Results of statistical comparison with reference data

Chemical	Environment								
	Terrestrial	Streams		Ponds		Lagoons		Marine	
	Tundra Soil	Sediment	Water	Sediment	Water	Sediment	Water	Sediment	Water
Aluminum	Pass	--	Pass	--	Pass	Pass	Pass	Pass	Pass
Antimony	NA	Fail	NA	Fail	NA	Fail	Fail	NA	Pass
Arsenic	Fail	Fail	--	--	--	Fail	Pass	Pass	--
Barium	NA	Pass	NA	Pass	Pass	Pass	Pass	Fail	Pass
Cadmium	Fail	Fail	NA	Fail	NA	NA	--	Fail	--
Chromium	Pass	--	--	--	--	--	--	--	--
Cobalt	NA	Fail	Pass	NA	Pass	Pass	Pass	Pass	Pass
Copper	Fail	--	--	Pass	Pass	--	--	Fail	NA
Fluoride	--	NA	Fail	NA	Pass	NA	Fail	Pass	Pass
Iron	Pass	--	Pass	--	Pass	Pass	Pass	Pass	Pass
Lead	Fail	Fail	NA	NA	Pass	Fail	--	NA	--
Manganese	NA	--	NA	--	NA	Fail	NA	Pass	Pass
Mercury	NA	--	--	NA	--	--	--	NA	--
Molybdenum	Fail	Fail	Fail	Fail	Pass	Pass	Fail	Pass	Pass
Nickel	Pass	Fail	--	Pass	--	Pass	Pass	Pass	NA
Selenium	NA	Fail	--	Pass	--	Pass	--	NA	--
Silver	NA	Fail	--	NA	--	--	--	Fail	--
Strontium	Fail	Fail	Pass	Fail	Pass	Pass	Pass	Fail	Fail
Thallium	NA	Fail	NA	Pass	NA	Pass	NA	Pass	NA
Tin	--	NA	NA	NA	NA	NA	NA	NA	NA
Vanadium	NA	Pass	NA	Pass	Pass	Pass	NA	Pass	NA
Zinc	Fail	Fail	NA	Fail	NA	Fail	Pass	Fail	--

**Note:** -- - chemical passed earlier screening tier  
 Fail - site concentrations significantly greater than reference concentrations  
 NA - not applicable; no statistical comparison was made because of high frequency of undetected results; or the confidence interval for the site mean straddles zero as a result of small sample size or high variability  
 Pass - site concentrations not significantly greater than reference concentrations

Table 3-38. Chemicals of potential concern retained for ecological risk analysis

Chemical	Environment								
	Terrestrial	Streams		Ponds		Lagoons		Marine	
	Tundra Soil	Sediment	Water	Sediment	Water	Sediment	Water	Sediment	Water
Aluminum	--	--	--	--	--	--	--	--	--
Antimony	Fail	NB	NB	NB	NB	NB	NB	NB	--
Arsenic	Fail	Fail	--	--	--	Fail	--	--	--
Barium	Fail	--	NB	--	--	--	--	NB	--
Cadmium	Fail	Fail	Fail	Fail	Fail	Fail	--	Fail	--
Chromium	--	--	--	--	--	--	--	--	--
Cobalt	Fail	NB	--	NB	--	--	--	--	--
Copper	Fail	--	--	--	--	--	--	Fail	Fail
Fluoride	--	ND <sup>a</sup>	NB	NB	--	NB	NB	--	--
Iron	--	--	--	--	--	--	--	--	--
Lead	Fail	Fail	Fail	Fail	--	Fail	--	Fail	--
Manganese	Fail	--	NB	--	NB	NB	NB	--	--
Mercury	Fail	--	--	Fail	--	--	--	Fail	--
Molybdenum	Fail	NB	NB	NB	--	--	NB	--	--
Nickel	--	Fail	--	--	--	--	--	--	ND <sup>b</sup>
Selenium	Fail	NB	--	--	--	--	--	NB	--
Silver	Fail	NB	--	NB	--	--	--	Fail	--
Strontium	NB	NB	--	NB	--	--	--	NB	NB
Thallium	Fail	NB	NB	--	NB	--	NB	--	NB
Tin	--	NB	NB	NB	NB	NB	NB	NB	NB
Vanadium	Fail	--	NB	--	--	--	NB	--	NB
Zinc	Fail	Fail	Fail	Fail	Fail	Fail	--	Fail	--

**Note:** -- - chemical eliminated from further evaluation

Fail - chemical retained as a CoPC for the baseline ERA

NB - no benchmark; chemical retained as a CoPC for the baseline ERA

ND - undetected in all samples; chemical retained as a CoPC for the baseline ERA

<sup>a</sup> No benchmark.

<sup>b</sup> Maximum value expressed as one-half of the detection limit is above the screening benchmark.

Table 3-39. Data needs for the ecological risk assessment

Environment	Assessment Endpoint	Representative Receptor	Food Item	Data Need
Tundra	Structure and function of terrestrial plant communities	Terrestrial plant communities	NA	Tundra plant community surveys
Tundra	Structure and function of tundra soil fauna communities	Tundra soil fauna communities	NA	None. Not directly assessed; evaluated through terrestrial plant community analysis.
Tundra	Survival, growth, and reproduction of terrestrial avian herbivore populations	Willow ptarmigan	Terrestrial plants (willow and sedge)	CoPCs in terrestrial plants <sup>a</sup>
Tundra	Survival, growth, and reproduction of terrestrial mammalian herbivore populations	Tundra vole; caribou; moose	Terrestrial plants (willow, sedge, and/or lichen)	CoPCs in terrestrial plants <sup>a</sup>
Tundra	Survival, growth, and reproduction of terrestrial avian invertivore populations	Lapland longspur	Terrestrial invertebrates	CoPCs in terrestrial invertebrates <sup>a</sup>
Tundra	Survival, growth, and reproduction of terrestrial mammalian invertivore populations	Tundra shrew	Terrestrial invertebrates	CoPCs in terrestrial invertebrates <sup>a</sup>
Tundra	Survival, growth, and reproduction of terrestrial avian carnivore populations	Snowy owl	Small mammals (voles or lemmings)	CoPCs in small mammals <sup>a</sup>
Tundra	Survival, growth, and reproduction of terrestrial mammalian carnivore populations	Arctic fox	Small mammals (voles or lemmings)	CoPCs in small mammals <sup>a</sup>
Streams	Structure and function of stream aquatic and wetland plant communities	Stream aquatic and wetland plant communities	NA	CoPCs in stream aquatic/wetland plants (arsenic, cadmium, lead, nickel, and zinc)
Streams	Structure and function of stream aquatic invertebrate communities	Stream aquatic invertebrate communities	NA	Stream aquatic invertebrate community surveys
Streams	Survival, growth, and reproduction of stream avian herbivore populations	Green-winged teal	Aquatic/wetland plants (sedge)	CoPCs in stream aquatic/wetland plants <sup>a</sup>
Streams	Survival, growth, and reproduction of stream mammalian herbivore populations	Muskrat	Aquatic/wetland plants (sedge)	CoPCs in stream aquatic/wetland plants <sup>a</sup>
Streams	Survival, growth, and reproduction of stream avian invertivore populations	Common snipe	Aquatic invertebrates	CoPCs in stream invertebrates (cadmium, lead, mercury, and zinc) <sup>b</sup>
Tundra ponds	Structure and function of tundra pond aquatic and wetland plant communities	Tundra pond aquatic and wetland plant communities	NA	Tundra pond and wetland plant community surveys <sup>c</sup>



Table 3-39. (cont.)

Environment	Assessment Endpoint	Representative Receptor	Food Item	Data Need
Tundra ponds	Structure and function of tundra pond aquatic invertebrate communities	Tundra pond aquatic invertebrate communities	NA	Tundra pond aquatic invertebrate community surveys <sup>b</sup>
Tundra ponds	Survival, growth, and reproduction of tundra pond avian herbivore populations	Green-winged teal	Aquatic/wetland plants (sedge)	CoPCs in tundra pond aquatic/wetland plants <sup>a</sup>
Tundra ponds	Survival, growth, and reproduction of tundra pond mammalian herbivore populations	Muskrat	Aquatic/wetland plants (sedge)	CoPCs in tundra pond aquatic/wetland plants <sup>a</sup>
Tundra ponds	Survival, growth, and reproduction of tundra pond avian invertivore populations	Common snipe	Aquatic invertebrates	CoPCs in tundra pond aquatic invertebrates (arsenic, barium, cadmium, lead, mercury, thallium, and zinc) <sup>b</sup>
Coastal lagoons	Structure and function of coastal lagoon aquatic and wetland plant communities	Coastal lagoon aquatic and wetland plant communities	NA	Coastal lagoon aquatic and wetland plant community surveys
Coastal lagoons	Structure and function of coastal lagoon aquatic invertebrate communities	Coastal lagoon aquatic invertebrate communities	NA	Coastal lagoon aquatic invertebrate community surveys
Coastal lagoons	Structure and function of coastal lagoon fish communities	Coastal lagoon fish	NA	CoPCs in coastal lagoon fish <sup>d</sup>
Coastal lagoons	Survival, growth, and reproduction of coastal lagoon avian herbivore populations	Brant	Aquatic/wetland plants (sedge)	CoPCs in coastal lagoon aquatic/wetland plants <sup>a</sup>
Coastal lagoons	Survival, growth, and reproduction of coastal lagoon avian invertivore populations	Black-bellied plover	Aquatic invertebrates	CoPCs in coastal lagoon aquatic invertebrates (cadmium, lead and zinc)
Coastal lagoons	Survival, growth, and reproduction of coastal lagoon avian piscivore populations	Red-throated loon	Fish	CoPCs in coastal lagoon fish <sup>a</sup>

**Note:** CoPC - chemical of potential concern  
 NA - not applicable

<sup>a</sup> CoPCs for all herbivores, terrestrial invertivores, terrestrial carnivores, and lagoon piscivores are aluminum, antimony, arsenic, barium, cadmium, chromium, cobalt, lead, mercury, molybdenum, selenium, thallium, vanadium, and zinc.

<sup>b</sup> Data for terrestrial invertebrate samples collected during the Phase II field event will be used to evaluate this assessment endpoint.

<sup>c</sup> Data for terrestrial plant community surveys collected during the Phase II field event will be used to evaluate this assessment endpoint.

<sup>d</sup> CoPCs for coastal lagoon fish are arsenic, cadmium, lead, and zinc.

Table 4-1. Overview of Phase II data

Assessment Endpoint	Station/Transect	Distance (m)	Small Mammals	Ptarmigan	Terrestrial Invertebrate Tissue	Vegetation					Salmon-berries	Sour-dock	Vegetation Plots	Tundra Soil	Aquatic Invertebrates		Sediment		Water Quality Parameters
						Willow	Birch	Sedge	Lichen						Tissue	Community Analysis	Chemistry	Toxicity Test	
									Peltigera	Cladina									
Terrestrial	TT5	10			X	X		X	X			X	X						
		20	X										X						
		100	a		X	X		X	X			X	X						
		1,000	X		X	X	X	X	X			X	X						
		2,000	X		X		X	X		X		X	X						
	TT2	10			X	X		X	X					X					
		20	a											X					
		100	X		X	X		X	X					X					
		1,000	X		X	X		X	X					X					
	TT8	10					X		X	X				X	X				
		20	a												X				
		50												X <sup>o</sup>	X				
		100	a				X		X	X				X	X				
		150												X <sup>b</sup>	X				
		200												X <sup>b</sup>	X				
		250												X <sup>b</sup>	X				
		300												X <sup>b</sup>	X				
		350												X <sup>b</sup>	X				
		400												X <sup>b</sup>	X				
		450												X <sup>b</sup>	X				
		500												X <sup>b</sup>	X				
		550												X <sup>b</sup>	X				
		600												X <sup>b</sup>	X				
		650												X <sup>b</sup>	X				
		700												X <sup>b</sup>	X				
		750												X <sup>b</sup>	X				
	800												X <sup>b</sup>	X					
	900												X <sup>o</sup>	X					
	1,000						X		X	X	X		X	X					
	TT3	10				X	X		X	X				X	X				
		20	X												X				
		100	X			X	X	X	X	X				X	X				
		1,000	X			X		X	X	X	X			X	X				
	TT6	10				X	X		X	X	X			X	X				
		20	c												X				
		100				X	X		X	X	X			X	X				
		1,000				X	X		X	X	X			X	X				
		2,000					X		X	X	X				X				
	TT7	10					X		X		X				X				
		1,000					X		X		X				X				
		2,000					X		X		X				X				

Table 4-1. (cont.)

Assessment Endpoint	Station/ Transect	Distance (m)	Small Mammals	Ptarmigan	Terrestrial Invertebrate Tissue	Vegetation						Salmon-berries	Sour-dock	Vegetation Plots	Tundra Soil	Aquatic Invertebrates		Sediment		Water Quality Parameters
						Willow	Birch	Sedge	Lichen		Tissue					Analysis	Chemistry	Toxicity Test		
									Peltigera	Cladina										
	TS-REF-5		X		X	X		X	X	X			X	X						
	TS-REF-7					X	X	X	X	X			X	X						
	TS-REF-11					X	X	X	X	X			X	X						
	TS-REF-12												X	X						
	Near the DMTS road																			
	Terrestrial reference area																			
	Site A																			
	Site B																			
	Site C																			
<b>Streams</b>																				
	AC-R							X						X	X	X	X		X	
	OR-R							X	X					X	X	X	X		X	
	ARC-R							X	X					X	X	X	X		X	
	ST-REF-3							X	X					X	X	X	X		X	
	ST-REF-5							X	X					X	X	X	X		X	
	ST-REF-6							X	X					X	X	X	X		X	
<b>Tundra Ponds</b>																				
	TP1	100						X						X					X	
		1,000						X						X					X	
	TP3							X						X					X	
	TP4							X						X					X	
	TP-REF-2							X						X					X	
	TP-REF-3							X						X					X	
	TP-REF-5							X						X					X	
<b>Coastal Lagoons</b>																				
	PLNL							X						X	X	X <sup>e</sup>	X	X	X	
	NLK							X						X	X	X <sup>e</sup>	X	X	X	
	NLF													X	X	X <sup>e</sup>	X	X	X	
	CL-REF-1							X						X	X	X <sup>e</sup>	X	X	X	
	CL-REF-2							X						X	X <sup>f</sup>	X <sup>e</sup>	X	X	X	
	CL-REF-3													X		X <sup>e</sup>	X	X	X	
<b>Marine (Pre-shipment - June 2004)</b>																				
	NMA																		X	
	NMB																		X	
	NMC																		X	
	NMD																		X	
	NME																		X	
	NMF																		X	
	NMG																		X	
	NMH																		X	
	NMJ																		X	

Table 4-1. (cont.)

Assessment Endpoint	Station/ Transect	Distance (m)	Small Mammals	Ptarmigan	Terrestrial Invertebrate Tissue	Vegetation					Salmon- berries	Sour- dock	Vegetation Plots	Tundra Soil	Aquatic Invertebrates		Sediment		Water Quality Parameters	
						Willow	Birch	Sedge	Lichen						Community	Chemistry	Toxicity Test			
									Peltigera	Cladina										
	NMK																X			
	NML																	X		
	NMM																	X		
	NMN																	X		
	NMO																	X		
	NMP																	X		
	NMQ																	X		
	NMS																	X		
	NMT																	X		
	NMU																	X		
	NMV																	X		
	NMW																	X		
	NMX																	X		
	NMY																	X		
	NMZ																	X		
	NMAA																	X		
	NMGZ																	X		
	NM-REF-1																	X		
	NM-REF-2																	X		
	NM-REF-3																	X		

<sup>a</sup> Grid set; no small mammals collected.

<sup>b</sup> Single microplot for vegetation community.

<sup>c</sup> No small mammal grid set; not correct habitat (too open).

<sup>d</sup> Five washed and five unwashed samples were collected at each site.

<sup>e</sup> Samples archived.

<sup>f</sup> Aquatic invertebrate tissue sample collected from area encompassed by both Stations CL-REF-2 and CL-REF-3.

Table 4-2. Phase II data collection matrix

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
<b>Terrestrial</b>							
<b>Small Mammals (presented in ascending order from port facility to mine along the DMTS)</b>							
TT5	-- Port transect 20 m north of road 100 m north of road 1,000 m north of road 2,000 m north of road	4	3	5; 2 brown lemmings and 3 tundra voles None 3 northern red backed voles 1 masked shrew	Tissue chemistry; whole body; each individual mammal equals one sample	List 1 <sup>a</sup>	Grid set; no small mammals were collected
TT2	-- DMTS road transect 20 m north of road 100 m north of road 1,000 m north of road	3	2	None 3 northern red backed voles 1 northern red backed vole			Grid set; no small mammals were collected
TT8	-- DMTS road transect 20 m north of road 100 m north of road	0	0	None None			Stations were added in the field. Grid set; no small mammals were collected Grid set; no small mammals were collected
TT3	-- DMTS road transect 20 m north of road 100 m north of road 1,000 m north of road	3	3	1 masked shrew 2; 1 masked shrew and 1 tundra shrew 3; 2 northern red backed voles and 1 masked shrew			
TT6	-- DMTS road transect 20 m north of road 100 m north of road 1,000 m north of road	3	0	None None None			No grid set; not correct habitat (too open) Grid set; no small mammals were collected Grid set; no small mammals were collected
TT7	-- Solid waste permit boundary transect 10 m downwind of boundary 1,000 m downwind of mine	2	0	None None			No grid set; not correct habitat (rock face) No grid set; not correct habitat (rock face)
TS-REF-5	-- Terrestrial reference area	1	1	4; 3 masked shrews and 1 northern red backed vole			
<b>Ptarmigan</b>	Near the DMTS road Terrestrial reference area	NA NA	NA NA	5 individual birds 3 individual birds	Tissue chemistry; breast muscle tissue (skin on), liver, and kidneys from each bird analyzed separately	List 2: Antimony, barium, cadmium, lead, thallium, and zinc	
<b>Soil Invertebrate Tissue (presented in ascending order from port facility to mine along the DMTS)</b>							
TT5	-- Port transect 10 m north of road 100 m north of road 1,000 m north of road 2,000 m north of road	4	4	1 spiders-only composite and 1 multi-species composite 1 crane flies-only composite and 1 multi-species composite 2 spiders-only composites and 1 multi-species composite 1 multi-species composite	Tissue chemistry; whole body; composite tissue sample of all soil invertebrates collected at a given station	List 1 <sup>a</sup>	

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
TT2	-- DMTS road transect 10 m north of road 100 m north of road 1,000 m north of road	3	3	1 multi-species composite 1 multi-species composite 1 multi-species composite			
TT3	-- DMTS road transect 10 m north of road 100 m north of road 1,000 m north of road	0	3	1 multi-species composite 1 multi-species composite 1 multi-species composite			Stations were added in the field.
TT6	-- DMTS road transect 10 m north of road 100 m north of road 1,000 m north of road	0	3	1 multi-species composite 1 multi-species composite 1 multi-species composite			Stations were added in the field.
TS-REF-5	-- Terrestrial reference area	1	1	1 multi-species composite			
<b>Vegetation Tissue (presented in ascending order from port facility to mine along the DMTS)</b>							
TT5	-- Port transect 10 m north of road 100 m north of road 1,000 m north of road 2,000 m north of road	4	4	3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 4; willow, birch, sedge, and lichen (Peltigera) (1 composite sample per species) 3; birch, sedge, and lichen (Cladina) (1 composite sample per species)	Tissue chemistry; unwashed willow or birch leaves (debris removed in field), unwashed sedge blades (minimum 3 plants per station), unwashed lichen (debris removed in field with minimum 3 plants per station)	List 1 <sup>a</sup>	Willow and birch leaves were collected at this station. No willow leaves were collected at this station. Birch leaves were collected at this station.
TT2	-- DMTS road transect 10 m north of road 1,000 m north of road 1 km north of road	3	3	3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 3; willow, sedge, and lichen (Peltigera) (1 composite sample per species)			
TT8	-- DMTS road transect 10 m north of road 100 m north of road 1,000 m north of road	3	3	3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 3; willow, sedge, and lichen (Peltigera) (1 composite sample per species) 4; willow, sedge, and lichen (both Peltigera and Cladina) (1 composite sample per species)			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
TT3	-- DMTS road transect 10 m north of road	3	3	3; willow, sedge, and lichen ( <i>Peltigera</i> ) (1 composite sample per species)			
	100 m north of road			4; willow, birch, sedge, and lichen ( <i>Peltigera</i> ) (1 composite sample per species)		Willow and birch leaves were collected at this station.	
	1,000 m north of road			4; birch, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)		No willow leaves were collected at this station. Birch leaves were collected at this station.	
TT6	-- Port transect 10 m north of road	4	4	4; willow, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			
	100 m north of road			4; willow, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			
	1,000 m north of road			4; willow, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			
	2,000 m north of road			4; willow, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			
TT7	-- Solid waste permit boundary transect 10 m downwind of boundary	3	3	3; willow, sedge, and lichen ( <i>Cladina</i> ) (1 composite sample per species)			
	1,000 m downwind of mine			3; willow, sedge, and lichen ( <i>Cladina</i> ) (1 composite sample per species)			
	2,000 m downwind of mine			3; willow, sedge, and lichen ( <i>Cladina</i> ) (1 composite sample per species)			
TS-REF-5	-- Terrestrial reference area	1	1	4; willow, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			
TS-REF-7	-- Terrestrial reference area	1	1	5; willow, birch, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			Willow and birch leaves were collected at this station.
TS-REF-11	-- Terrestrial reference area	1	1	5; willow, birch, sedge, and lichen ( <i>Peltigera</i> and <i>Cladina</i> ) (1 composite sample per species)			Willow and birch leaves were collected at this station.
<b>Berries</b>							
Site A	-- just north of the port ambient air boundary at Ipiavik Lagoon	1	1	10; 5 washed and 5 unwashed salmonberry samples	Tissue chemistry; for all washed samples any debris was removed in field	Antimony, barium, cadmium, lead, thallium, and zinc	
Site B	-- north of the port facility but closer to Kivalina	1	1	10; 5 washed and 5 unwashed salmonberry samples			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
Site C	-- reference area north of Kivalina	1	1	10; 5 washed and 5 unwashed salmonberry samples			
Site D	-- south of Site A on Ipiavik Lagoon but closer to the port facility	0	1	10; 5 washed and 5 unwashed salmonberry samples			
<b>Sourdock</b>							
Site A	-- just north of the port ambient air boundary at Ipiavik Lagoon	1	1	10; 5 washed and 5 unwashed sourdock samples	Tissue chemistry; for all washed samples any debris was removed in field; minimum 3 sourdock plants per station	Antimony, barium, cadmium, lead, thallium, and zinc	
Site B	-- north of the port facility but closer to Kivalina	1	1	10; 5 washed and 5 unwashed sourdock samples			
Site C	-- reference area north of Kivalina	1	1	10; 5 washed and 5 unwashed sourdock samples			
Site D	-- south of Site A on Ipiavik Lagoon but closer to the port facility	0	1	10; 5 washed and 5 unwashed sourdock samples			
<b>Vegetation Plots (presented in ascending order from port facility to mine along the DMTS)</b>					Community analysis	--	
TT5	-- Port transect	4	4				
	10 m north of road			--			
	100 m north of road			--			
	1,000 m north of road			--			
	2,000 m north of road			--			
TT8	-- DMTS road transect	3	19				
	10 m north of road			--			Station was added in the field.
	50 m north of road			--			Single microplot; station was added in the field.
	100 m north of road			--			
	150 m north of road			--			Single microplot; station was added in the field.
	200 m north of road			--			Single microplot; station was added in the field.
	250 m north of road			--			Single microplot; station was added in the field.
	300 m north of road			--			Single microplot; station was added in the field.
	350 m north of road			--			Single microplot; station was added in the field.
	400 m north of road			--			Single microplot; station was added in the field.
	450 m north of road			--			Single microplot; station was added in the field.
	500 m north of road			--			Single microplot; station was added in the field.
	550 m north of road			--			Single microplot; station was added in the field.
	600 m north of road			--			Single microplot; station was added in the field.
	650 m north of road			--			Single microplot; station was added in the field.
	700 m north of road			--			Single microplot; station was added in the field.
	750 m north of road			--			Single microplot; station was added in the field.
	800 m north of road			--			Single microplot; station was added in the field.
	900 m north of road			--			Single microplot; station was added in the field.
	1,000 m north of road			--			Single microplot; station was added in the field.
TT3	-- DMTS road transect	3	3				
	10 m north of road			--			
	100 m north of road			--			
	1,000 m north of road			--			
TT6	-- Port transect	3	3				
	10 m north of road			--			
	100 m north of road			--			
	1,000 m north of road			--			



Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
TS-REF-5	-- Terrestrial reference area	1	1	--			
TS-REF-7	-- Terrestrial reference area	1	1	--			
TS-REF-11	-- Terrestrial reference area	1	1	--			Reference area station location was modified to better match vegetation community at site stations.
TS-REF-12	-- Terrestrial reference area	0	1	--			Station was added in the field.
<b>Tundra Soil (presented in ascending order from port facility to mine along the DMTS)</b>					Chemistry; 0–2 cm	List 3 <sup>b</sup> and pH	
TT5	-- Port transect	5	5				
	10 m north of road			1			
	20 m north of road			1			
	100 m north of road			1			
	1,000 m north of road			1			
	2,000 m north of road			1			
TT2	-- DMTS road transect	4	4				
	10 m north of road			1			
	20 m north of road			1			
	100 m north of road			1			
	1,000 m north of road			1			
TT8	-- DMTS road transect	3	19				
	10 m north of road			1			
	50 m north of road			1			Station was added in the field.
	100 m north of road			1			
	150 m north of road			1			Station was added in the field.
	200 m north of road			1			Station was added in the field.
	250 m north of road			1			Station was added in the field.
	300 m north of road			1			Station was added in the field.
	350 m north of road			1			Station was added in the field.
	400 m north of road			1			Station was added in the field.
	450 m north of road			1			Station was added in the field.
	500 m north of road			1			Station was added in the field.
	550 m north of road			1			Station was added in the field.
	600 m north of road			1			Station was added in the field.
	650 m north of road			1			Station was added in the field.
	700 m north of road			1			Station was added in the field.
	750 m north of road			1			Station was added in the field.
	800 m north of road			1			Station was added in the field.
	900 m north of road			1			Station was added in the field.
	1,000 m north of road			1			Station was added in the field.
TT3	-- DMTS road transect	6	4				
	10 m north of road			1			
	20 m north of road			1			
	100 m north of road			1			
	1,000 m north of road			1			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
TT6	-- Port transect	5	4				
	10 m north of road			1			
	20 m north of road			0			No tundra soil sample was collected at 20 m.
	100 m north of road			1			
	1,000 m north of road			1			
	2,000 m north of road			1			
TT7	-- Solid waste permit boundary transect	3	3				
	10 m downwind of boundary			1			
	1,000 m downwind of mine			1			
	2,000 m downwind of mine			1			
TS-REF-5	-- Terrestrial reference area	1	1	1			
TS-REF-7	-- Terrestrial reference area	1	1	1			
TS-REF-11	-- Terrestrial reference area	1	1	1			Reference area station location was modified to better match vegetation community at site stations.
TS-REF-12	-- Terrestrial reference area	0	1	1			Station was added in the field.
<b>Streams</b>							
<b>Aquatic Invertebrate Tissue (presented in ascending order from port facility to mine along the DMTS)</b>					Tissue chemistry	Cadmium, lead, mercury, and zinc	Stations were added in the field.
AC-R	Aufeis Creek	0	1	1 multi-species composite			
OR-R	Omikviorok River	0	1	1 multi-species composite			
ARC-R	Anxiety Ridge Creek	0	1	1 multi-species composite			
ST-REF-3	-- Freshwater aquatic reference area	0	1	1 multi-species composite			
ST-REF-6	-- Freshwater aquatic reference area	0	1	1 multi-species composite			
<b>Aquatic Invertebrate Community (presented in ascending order from port facility to mine along the DMTS)</b>					Community analysis	--	
AC-R	Aufeis Creek	1	1	5 replicates per station			
OR-R	Omikviorok River	1	1	5 replicates per station			
ARC-R	Anxiety Ridge Creek	1	1	5 replicates per station			
ST-REF-3	-- Freshwater aquatic reference area	1	1	5 replicates per station			
ST-REF-5	-- Freshwater aquatic reference area	0	1	5 replicates per station			
ST-REF-6	-- Freshwater aquatic reference area	0	1	5 replicates per station			
<b>Vegetation Tissue (presented in ascending order from port facility to mine along the DMTS)</b>					Tissue chemistry; unwashed willow leaves (debris removed in field), sedge plant (rinsed roots [no sediment] and unwashed blades with minimum 3 plants per station)	List 1 <sup>a</sup>	
AC-R	Aufeis Creek	1	1	1; willow (1 composite sample)			Sedge was not collected at this station.
OR-R	Omikviorok River	1	1	2; willow and sedge (1 composite sample per species)			
ARC-R	Anxiety Ridge Creek	1	1	2; willow and sedge (1 composite sample per species)			
ST-REF-3	-- Freshwater aquatic reference area	1	1	2; willow and sedge (1 composite sample per species)			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
ST-REF-5	-- Freshwater aquatic reference area	1	1	2; willow and sedge (1 composite sample per species)			
ST-REF-6	-- Freshwater aquatic reference area	1	1	2; willow and sedge (1 composite sample per species)			
<b>Tundra Soil (presented in ascending order from port facility to mine along the DMTS)</b>					Chemistry; 0–2 cm	List 3 <sup>b</sup> and pH	
AC-R	Aufeis Creek	1	1	1			
OR-R	Omikviorok River	1	1	1			
ARC-R	Anxiety Ridge Creek	1	1	1			
ST-REF-3	-- Freshwater aquatic reference area	1	1	1			
ST-REF-5	-- Freshwater aquatic reference area	1	1	1			
ST-REF-6	-- Freshwater aquatic reference area	1	1	1			
<b>Stream Sediment (presented in ascending order from port facility to mine along the DMTS)</b>					Chemistry; 0–2 cm	List 3 <sup>b</sup> and pH	Stations were added in the field (associated with the stream aquatic invertebrate tissue samples).
AC-R	Aufeis Creek	0	1	1			
OR-R	Omikviorok River	0	1	1			
ARC-R	Anxiety Ridge Creek	0	1	1			
ST-REF-3	-- Freshwater aquatic reference area	0	1	1			
ST-REF-6	-- Freshwater aquatic reference area	0	1	1			
<b>Stream Water (presented in ascending order from port facility to mine along the DMTS)</b>					Field measurements	Water quality parameters <sup>c</sup>	
AC-R	Aufeis Creek	1	1	1			
OR-R	Omikviorok River	1	1	1			
ARC-R	Anxiety Ridge Creek	1	1	1			
ST-REF-3	-- Freshwater aquatic reference area	1	1	1			
ST-REF-5	-- Freshwater aquatic reference area	1	1	1			
ST-REF-6	-- Freshwater aquatic reference area	1	1	1			
<b>Tundra Ponds</b>							
<b>Vegetation Tissue (presented in ascending order from port facility to mine along the DMTS)</b>					Tissue chemistry;	List 1 <sup>a</sup>	
TP1	-- Port transect						
	100 m north of road	1	1	1			
	1 km north of road	1	1	1			
	--DMTS road						
	TP3	1	1	1			
	TP4	1	1	1			
TP-REF-2	-- Freshwater aquatic reference area	1	1	1			
TP-REF-3	-- Freshwater aquatic reference area	1	1	1			
TP-REF-5	-- Freshwater aquatic reference area	1	1	1			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
<b>Tundra Soil (presented in ascending order from port facility to mine along the DMTS)</b>					Chemistry; 0–2 cm	List 3 <sup>b</sup>	
TP1	-- Port transect						
	100 m north of road	1	1	1			
	1 km north of road	1	1	1			
	--DMTS road						
	TP3	1	1	1			
	TP4	1	1	1			
TP-REF-2	-- Freshwater aquatic reference area	1	1	1			
TP-REF-3	-- Freshwater aquatic reference area	1	1	1			
TP-REF-5	-- Freshwater aquatic reference area	1	1	1			
<b>Tundra Pond Water (presented in ascending order from port facility to mine along the DMTS)</b>					Field measurements	Water quality parameters <sup>c</sup>	
TP1	-- Port transect						
	100 m north of road	1	1	1			
	1 km north of road	1	1	1			
	--DMTS road						
	TP3	1	1	1			
	TP4	1	1	1			
TP-REF-2	-- Freshwater aquatic reference area	1	1	1			
TP-REF-3	-- Freshwater aquatic reference area	1	1	1			
TP-REF-5	-- Freshwater aquatic reference area	1	1	1			
<b>Coastal Lagoons</b>							
<b>Aquatic Invertebrate Tissue (presented in ascending order from port facility to the north))</b>					Tissue chemistry; composite sample of all invertebrates collected at a station	List 4: Cadmium, lead, and zinc	
PLNL	Port Lagoon North (inland shore)	1	1	1 multi-species composite			
NLK	North Lagoon (inland shore)	1	1	1 multi-species composite			
NLF	North Lagoon (seaward shore)	1	1	1 multi-species composite			
CL-REF-1	-- Reference lagoon	1	1	1 multi-species composite			
CL-REF-2/3	-- Control lagoon	0	1	1 multi-species composite			Station was added in the field.
<b>Aquatic Invertebrate Community (presented in ascending order from port facility to the north)</b>					Community analysis	--	
PLNL	Port Lagoon North (inland shore)	1	1	5 replicates per station			
NLK	North Lagoon (inland shore)	1	1	5 replicates per station			
NLF	North Lagoon (seaward shore)	1	1	5 replicates per station			
CL-REF-1	-- Reference lagoon	1	1	5 replicates per station			
CL-REF-2	-- Control lagoon (inland shore)	1	1	5 replicates per station			
CL-REF-3	-- Control lagoon (seaward shore)	1	1	5 replicates per station			
<b>Fish</b>							
	2 site lagoons	2	0	0			All 3 coastal lagoons were seined and trapped; no fish were collected.
	Reference lagoon TBD	1	0	0			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
<b>Vegetation Tissue (presented in ascending order from port facility to the north)</b>					Tissue chemistry; sedge plant (rinsed roots [no sediment] and unwashed blades with minimum 3 plants per station); entire plant will be sampled	List 1 <sup>a</sup>	
PLNL	Port Lagoon North (inland shore)	1	1	1			
NLK	North Lagoon (inland shore)	1	1	1			
NLF	North Lagoon (seaward shore)	1	0	0			No sedge was present at Station NLF.
CL-REF-1	-- Reference lagoon	1	1				
CL-REF-2	-- Control lagoon (inland shore)	1	1	1			
CL-REF-3	-- Control lagoon (northern shore)	1	0	0			
<b>Vegetation Plots (presented in ascending order from port facility to the north)</b>					Community analysis	--	
PLNL	Port Lagoon North (inland shore)	1	1	--			
NLK	North Lagoon (inland shore)	1	1	--			
NLF	North Lagoon (seaward shore)	1	0	--			No vegetation plots were surveyed at Station NLF; sand dune environment
CL-REF-1	-- Reference lagoon	1	1	--			
CL-REF-2	-- Control lagoon (inland shore)	1	1	--			
CL-REF-3	-- Control lagoon (northern shore)	1	0	--			No vegetation plots were surveyed at Station CL-REF-3
<b>Tundra Soil (presented in ascending order from port facility to the north)</b>					Chemistry; 0–2 cm	List 3 <sup>b</sup> and pH	
PLNL	Port Lagoon North (inland shore)	1	1	1			
NLK	North Lagoon (inland shore)	1	1	1			
NLF	North Lagoon (seaward shore)	1	1	1			
CL-REF-1	-- Reference lagoon	1	1	1			
CL-REF-2	-- Control lagoon (inland shore)	1	1	1			
CL-REF-3	-- Control lagoon (northern shore)	1	1	1			
<b>Lagoon Sediment (presented in ascending order from port facility to mine along the DMTS)</b>					Chemistry and toxicity test; 0–2 cm	List 5: Arsenic, cadmium, lead, zinc; List 6: Grain size and total solids; <i>Hyaella</i> survival and growth	
PLNL	Port Lagoon North (inland shore)	1	1	1			
NLK	North Lagoon (inland shore)	1	1	1			
NLF	North Lagoon (seaward shore)	1	1	1			
CL-REF-1	-- Reference lagoon	1	1	1			
CL-REF-2	-- Control lagoon (inland shore)	1	1	1			
CL-REF-3	-- Control lagoon (seaward shore)	1	1	1			
<b>Lagoon Water (presented in ascending order from port facility to the north)</b>					Field measurements	Water quality parameters <sup>c</sup>	
PLNL	Port Lagoon North (inland shore)	1	1	1			
NLK	North Lagoon (inland shore)	1	1	1			
NLF	North Lagoon (seaward shore)	1	1	1			
CL-REF-1	-- Reference lagoon	1	1	1			
CL-REF-2	-- Control lagoon (inland shore)	1	1	1			
CL-REF-3	-- Control lagoon (seaward shore)	1	1	1			

Table 4-2. (cont.)

Sample Type	Description	No. of Proposed Stations	No. of Stations Sampled	No. of Field Samples	Kind of Sample	Analytes	Comments
<b>Marine</b>							
Surface sediment	19 stations around the port	19	19	38 (two events) <sup>d</sup>	Chemistry and toxicity test; 0–2 cm	List 4: Cadmium, lead, zinc	
Surface sediment	7 port stations - NMD, NMGZ, NML, NMM, NMN, NMO, NMAA	7	7	14 (two events) <sup>d</sup>		List 7: Cadmium, copper, lead, mercury, silver, zinc; List 6: Grain size, total solids; <i>Hyalella</i> survival and growth <sup>e</sup>	
Surface sediment	3 reference stations to the southeast	3	3	6 (two events) <sup>d</sup>			

**Note:** DMTS - DeLong Mountain Regional Transportation System  
 NA - not applicable  
 TBD - to be determined

<sup>a</sup> List 1: Aluminum, antimony, arsenic, barium, cadmium, chromium, cobalt, lead, mercury, molybdenum, selenium, thallium, vanadium, and zinc.

<sup>b</sup> List 3: Antimony, arsenic, barium, cadmium, cobalt, copper, lead, manganese, mercury, molybdenum, selenium, silver, thallium, vanadium, and zinc.

<sup>c</sup> Water quality measurements were taken in the field (i.e., pH, dissolved oxygen, temperature, conductivity, and salinity).

<sup>d</sup> Marine sediment sampling events conducted in June and September 2004.

<sup>e</sup> The criteria described in the sampling and analysis plan (Exponent 2004b) were not met, so no sediment toxicity testing was conducted on marine sediments.

Table 5-1. Summary of exposure point concentrations for environmental media

	N	#ND	%ND	Min.	Max.	Mean	Std.Dev.	Distribution Tests			UCL		EPC
								normal	gamma	lognormal	Method	UCL	
<b>Stream Surface water (µg/L)</b>													
Lead	229	145	63%	0.018	7.3	0.33	0.75	--	--	--	--	--	0.33 Mean
Thallium	28	24	86%	0.016	0.55	0.055	0.11	No	No	No	Chebyshev NP	0.14	0.14 UCL
<b>Soil Subareas (mg/kg)</b>													
<b>Port Soil (mg/kg)</b>													
Antimony	23	18	78%	0.93	26.0	9.6	8.8	No	No	No	Chebyshev NP	17.5	17.5 UCL
Barium	23	0	0%	357	2,110	1,304	383	Yes	Yes	No	Student's-t	1,441	1,441 UCL
Cadmium	428	41	10%	0.40	388	27.6	39.2	No	No	No	Chebyshev NP <sup>a</sup>	39.4	39.4 UCL
Lead	433	12	3%	8.5	48,300	1,255	2,921	--	--	--	--	--	1,255 Mean
Thallium	4	0	0%	0.29	0.78	0.53	0.21		n < 10		--	--	0.78 Max
Zinc	433	0	0%	37.4	64,300	4,494	6,415	No	No	No	Chebyshev NP <sup>a</sup>	6,419	6,419 UCL
<b>Road Soil (mg/kg)</b>													
Antimony	12	6	50%	0.38	5.5	2.9	2.4	No	No	No	Chebyshev NP <sup>b</sup>	9.8	5.5 Max
Barium	12	0	0%	650	6,290	2,216	1,870	No	Yes	Yes	Approx. gamma	3,373	3,373 UCL
Cadmium	32	2	6%	0.50	29.3	4.0	5.5	No	No	No	Chebyshev NP	8.3	8.3 UCL
Lead	32	0	0%	13.5	2,440	198	423	--	--	--	--	--	198 Mean
Thallium	6	0	0%	0.11	0.46	0.22	0.13		n < 10		--	--	0.46 Max
Zinc	32	0	0%	102	4,840	731	952	No	No	Yes	H-statistic	962	962 UCL
<b>DMTS Area-weighted Soil (mg/kg)<sup>c</sup></b>													
Antimony	--	--	--	--	--	--	--	--	--	--			6.5 UCL/Max
Barium	--	--	--	--	--	--	--	--	--	--			3,219 UCL
Cadmium	--	--	--	--	--	--	--	--	--	--			10.8 UCL
Lead	--	--	--	--	--	--	--	--	--	--			282 Mean
Thallium	--	--	--	--	--	--	--	--	--	--			0.49 Max
Zinc	--	--	--	--	--	--	--	--	--	--			1,399 UCL

**Note:** All UCL calculations were done using ProUCL 3.0. UCL methods are recommendations per EPA guidance (U.S. EPA 2002b). Undetected sample results included based on one-half of the detection limit.

- - not applicable
- DMTS - DeLong Mountain Regional Transportation System
- EPC - exposure point concentration
- Min. - minimum result
- Max. - maximum result
- N - number of results
- ND - not detected
- NP - nonparametric
- Std.Dev. - standard deviation
- UCL - upper confidence limit

<sup>a</sup> 97.5% UCL was used to obtain 95% coverage level, per ProUCL recommendation.

<sup>b</sup> 99% UCL was used to obtain 95% coverage level, per ProUCL recommendation.

<sup>c</sup> A DMTS area-weighted soil concentration was derived for each metal assuming that the port area soil samples represent an area of 26 hectares and that the road area soil samples represent an area of 312 hectares (see Figure 5-2). The total assumed DMTS site area is (26 + 312) 338 hectares; therefore, the port soil mean was adjusted by 0.08 (26/338) and the road soil mean was adjusted by 0.92 (312/338): DMTS Area-weighted Soil = (Port Area EPC x 0.08) + (Road Area EPC x 0.92).

Table 5-2. Summary of exposure point concentrations for subsistence foods

	N	#ND	%ND	Min.	Max.	Mean	Std.Dev.	Distribution Tests			UCL Method	UCL	EPC
								normal	gamma	lognormal			
<b>Caribou (mg/kg wet)<sup>a</sup></b>													
<b>Caribou Tissue-Specific Data</b>													
<b>Kidney</b>													
Barium	--	--	--	--	--	--	--	--	--	--	--	--	3.2 <sup>b</sup>
Cadmium	11	0	0%	1.3	9.9	4.7	2.9	yes	yes	yes	Student's-t	6.3	6.3 UCL
Lead	11	0	0%	0.35	5.8	2.0	1.8	--	--	--	--	--	2.0 Mean
Zinc	11	0	0%	10.0	53.8	22.1	11.8	no	yes	yes	Approx. Gamma	29.1	29.1 UCL
<b>Liver</b>													
Barium	--	--	--	--	--	--	--	--	--	--	--	--	2.7 <sup>b</sup>
Cadmium	11	0	0%	0.36	3.3	1.4	0.96	yes	yes	yes	Student's-t	1.9	1.9 UCL
Lead	11	0	0%	0.72	5.6	2.6	1.7	--	--	--	--	--	2.6 Mean
Zinc	11	0	0%	20.3	120	39.1	28.0	no	yes	no	Approx. Gamma	54.1	54.1 UCL
<b>Muscle</b>													
Barium	--	--	--	--	--	--	--	--	--	--	--	--	1.2 <sup>b</sup>
Cadmium	11	3	27%	0.0050	0.080	0.041	0.025	yes	yes	no	Student's-t	0.055	0.055 UCL
Lead	11	0	0%	0.020	0.26	0.11	0.086	--	--	--	--	--	0.11 Mean
Zinc	11	0	0%	20.1	69.0	29.1	13.8	no	no	no	Modified-t NP	36.6	36.6 UCL
<b>Edible Tissue Weighted Average<sup>c</sup></b>													
Barium	--	--	--	--	--	--	--	--	--	--	--	--	1.3 <sup>b</sup>
Cadmium	33	--	--	--	--	--	--	--	--	--	--	--	0.22 UCL
Lead	33	--	--	--	--	--	--	--	--	--	--	--	0.19 Mean
Zinc	33	--	--	--	--	--	--	--	--	--	--	--	36.8 UCL
<b>Fish (mg/kg wet)</b>													
Lead	151	83	55%	0.0015	0.091	0.010	0.016	--	--	--	--	--	0.010 Mean
Thallium	--	--	--	--	--	--	--	--	--	--	--	--	0.0026 <sup>d</sup>
<b>Ptarmigan (mg/kg wet)</b>													
<b>Ptarmigan Tissue-Specific Data</b>													
<b>Breast</b>													
Barium	5	0	0%	0.040	0.48	0.19	0.17			n<10	--	--	0.48 Max
Cadmium	5	0	0%	0.16	0.48	0.31	0.12			n<10	--	--	0.48 Max
Lead	5	0	0%	0.011	0.045	0.025	0.013			n<10	--	--	0.025 Mean
Zinc	5	0	0%	6.3	10.2	8.6	1.5			n<10	--	--	10.2 Max
<b>Kidney</b>													
Barium	5	0	0%	0.38	3.8	1.2	1.5			n<10	--	--	3.8 Max
Cadmium	5	0	0%	52.6	108.1	80.9	26.2			n<10	--	--	108 Max
Lead	5	0	0%	0.44	2.7	1.3	0.9			n<10	--	--	1.3 Mean
Zinc	5	0	0%	41.0	67.1	54.5	9.7			n<10	--	--	67.1 Max



Table 5-2. (cont.)

	N	#ND	%ND	Min.	Max.	Mean	Std.Dev.	Distribution Tests			UCL	UCL	EPC
								normal	gamma	lognormal	Method		
<b>Liver</b>													
Barium	5	0	0%	0.12	0.53	0.29	0.16		n<10		--	--	0.53 Max
Cadmium	5	0	0%	7.8	22.5	15.2	6.8		n<10		--	--	22.5 Max
Lead	5	0	0%	0.11	0.97	0.38	0.34		n<10		--	--	0.38 Mean
Zinc	5	0	0%	28.2	64.8	41.8	14.1		n<10		--	--	64.8 Max
<b>Edible tissue weighted average<sup>e</sup></b>													
Barium	15	--	--	--	--	--	--	--	--	--	--	--	0.52 Max
Cadmium	15	--	--	--	--	--	--	--	--	--	--	--	3.5 Max
Lead	15	--	--	--	--	--	--	--	--	--	--	--	0.07 Mean
Zinc	15	--	--	--	--	--	--	--	--	--	--	--	15.7 Max
<b>Salmonberry (mg/kg wet)</b>													
Barium	6	0	0%	0.022	0.078	0.052	0.019		n<10		--	--	0.078 Max
Cadmium	23	0	0%	0.0069	0.21	0.037	0.040	no	yes	yes	Approx. Gamma	0.049	0.049 UCL
Lead	23	0	0%	0.00060	1.8	0.13	0.38	--	--	--	--	--	0.13 Mean
Zinc	23	0	0%	1.9	9.2	4.1	1.8	no	yes	yes	Approx. Gamma	4.7	4.7 UCL
<b>Sourdock (mg/kg wet)</b>													
Antimony	6	0	0%	0.0037	0.012	0.0084	0.0034		n<10		--	--	0.012 Max
Barium	6	0	0%	0.76	10.6	3.4	3.7		n<10		--	--	10.6 Max
Cadmium	12	0	0%	0.0032	0.021	0.010	0.0053	yes	yes	yes	Student's-t	0.013	0.013 UCL
Lead	12	0	0%	0.047	0.42	0.21	0.11	--	--	--	--	--	0.21 Mean
Thallium	6	4	67%	0.00012	0.00049	0.00020	0.00015		n<10		--	--	0.00049 Max
Zinc	12	0	0%	0.00012	7.4	4.6	1.5	yes	yes	yes	Student's-t	5.4	5.4 UCL

**Note:** All UCL calculations were done using ProUCL 3.0. UCL methods are recommendations per EPA guidance (U.S. EPA 2002b).

Undetected sample results included based on one-half of the detection limit.

- - not applicable
- CoPC - chemical of potential concern
- EPC - exposure point concentration
- Min. - minimum result
- Max. - maximum result
- N - number of results
- ND - not detected
- NP - non parametric
- Std.Dev. - standard deviation
- UCL - upper confidence limit
- EPA - U.S. Environmental Protection Agency

<sup>a</sup> Caribou tissue samples were not analyzed for antimony, barium, and thallium. Ptarmigan tissue EPCs were used to predict the caribou barium concentration. Antimony was never detected in ptarmigan, and thallium was only rarely detected and at concentrations near or below reference concentrations. Therefore, antimony and thallium were not included as caribou or ptarmigan CoPCs (see Section 5.2.1.2.1.2).

<sup>b</sup> This calculated EPC value used for barium is based on the relationship between barium and lead in the corresponding ptarmigan tissue (see Section 5.2.1.2.1.2).

<sup>c</sup> The EPC concentration for the edible caribou tissue weighted average was calculated using a mass-weighted calculation. Kidney and liver tissue each contributed 2 percent and muscle tissue contributed 96 percent of the concentration (ADPH 2001).

<sup>d</sup> This calculated EPC value used for thallium is based on the relationship between thallium and lead in stream surface water.

<sup>e</sup> The EPC concentration for the edible ptarmigan tissue weighted average was calculated using a mass-weighted calculation. Muscle tissue contributed 90 percent, kidney tissue contributed 1 percent, and liver tissue contributed 9 percent of the concentration (Kalas et al. 1995; Remington and Braun 1988) (Section 5.2.1.2.2).

Table 5-3. Calculation of predicted fish thallium exposure point concentration

	Max.	Mean	UCL	EPC	Ratio of Thallium Mean to Lead Mean in Surface Water	
<b>Stream Surface Water (<math>\mu\text{g/L}</math>)</b>						
Lead	7.3	0.33	0.55	0.33 Mean		
Thallium	0.55	0.055	0.14	0.14 UCL	0.17	(0.055/0.33)
	Max.	Mean	UCL	EPC	Calculation of Thallium EPC from Lead UCL in Fish	
<b>Fish (mg/kg wet)</b>						
Lead	0.091	0.010	0.016	0.010 Mean		
Thallium	--	--		0.0026 <sup>a</sup>	0.0026	(0.016*0.17)

**Note:** EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> The fish thallium EPC is calculated by multiplying the 95%UCL for lead in fish by the ratio of the mean thallium to mean lead concentrations in surface water.

Table 5-4. Calculation of predicted caribou barium exposure point concentrations for kidney, liver, and muscle tissue

					Ratios of Ptarmigan Mean Barium Value to Means for:		
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Kidney Tissue</b>							
<b>Ptarmigan (mg/kg wet)</b>							
Barium	3.8	1.2			0.015	0.96	0.023
Cadmium	108	80.9			(1.2/80.9)	(1.2/1.3)	(1.2/54.5)
Lead	2.7	1.3					
Zinc	67.1	54.5					
Calculation of Barium EPC from Caribou EPCs through Application of Ratios for Other Metals							
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Caribou (mg/kg wet)</b>							
Barium	--	--		3.2 <sup>a</sup>	0.10	3.2	0.66
Cadmium	9.9	4.7	6.3	6.3 UCL	(6.3*0.015)	(3.4*0.96)	(29.1*0.023)
Lead	5.82	1.97	3.4	2.0 Mean			
Zinc	53.8	22.1	29.1	29.1 UCL			
Ratios of Ptarmigan Mean Barium Value to Means for:							
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Liver Tissue</b>							
<b>Ptarmigan (mg/kg wet)</b>							
Barium	0.53	0.29			0.019	0.77	0.007
Cadmium	22.5	15.2			(0.29/15.2)	(0.29/0.38)	(0.29/41.8)
Lead	0.97	0.38					
Zinc	64.8	41.8					
Calculation of Barium EPC from Caribou EPCs through Application of Ratios for Other Metals							
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Caribou (mg/kg wet)</b>							
Barium	--	--		2.7 <sup>a</sup>	0.038	2.7	0.38
Cadmium	3.32	1.42	1.9	1.9 UCL	(1.9*0.019)	(3.5*0.77)	(54.1*0.007)
Lead	5.6	2.6	3.5	2.6 Mean			
Zinc	120	39.1	54.1	54.1 UCL			
Ratios of Ptarmigan Mean Barium Value to Means for:							
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Muscle Tissue</b>							
<b>Ptarmigan (mg/kg wet)</b>							
Barium	0.48	0.19			0.62	7.67	0.022
Cadmium	0.48	0.31			(0.19/0.31)	(0.19/0.025)	(0.19/8.6)
Lead	0.045	0.025					
Zinc	10.2	8.6					
Calculation of Barium EPC from Caribou EPCs through Application of Ratios for Other Metals							
	Max.	Mean	UCL	EPC	Cadmium	Lead	Zinc
<b>Caribou (mg/kg wet)</b>							
Barium	--	--		1.2 <sup>a</sup>	0.034	1.2	0.80
Cadmium	0.080	0.041	0.055	0.055 UCL	(0.055*0.62)	(0.16*7.67)	(36.6*0.022)
Lead	0.26	0.11	0.16	0.11 Mean			
Zinc	69.0	29.1	36.6	36.6 UCL			

Note: EPC - exposure point concentration

UCL - upper confidence limit

<sup>a</sup> The predicted caribou barium EPCs were calculated by:

- 1) Calculating ratios of mean barium to mean cadmium, lead, and zinc in each of the ptarmigan tissues (i.e., kidney, liver, and muscle). For all tissues the ratio of barium to lead gave the highest ratio.
- 2) Multiplying the barium to lead ratio for each tissue by the 95%UCL for lead in the corresponding caribou tissue.

Table 5-5. Ptarmigan tissue weight calculations

Tissue	Weight (g-wet weight)	Fraction of Total	Basis	Source
Kidney	3	0.01	Twice the highest value for one kidney reported for willow ptarmigan (range was 1.2–1.5 g).	Kalas et al. (1995)
Liver	26.5	0.09	Average liver weight for adult male and female sage grouse.	Remington and Braun (1988)
Muscle	257	0.91	Average weight for adult male and female sage grouse pectoralis and supracoracoideus muscles.	Remington and Braun (1988)
<b>Total</b>	<b>283.5</b>			

Table 5-6. EPA IEUBK lead model exposure parameters and input values

Parameter	Input Value(s)	Source
<b>Air</b>		
Outdoor air lead concentration ( $\mu\text{g}/\text{m}^3$ )	0.100	EPA default
Indoor air lead concentration (percent of outdoor air)	30%	EPA default
Time spent outdoors (hours/day)	1, 2, 3, 4, 4, 4, 4 <sup>a</sup>	EPA default
Ventilation rates ( $\text{m}^3/\text{day}$ )	2, 3, 5, 5, 5, 7, 7 <sup>a</sup>	EPA default
Lung absorption (percentage)	32	EPA default
<b>Diet</b>		
Diet intake ( $\mu\text{g}/\text{day}$ )	3.16, 2.60, 2.87, 2.74, 2.61, 2.74, 2.99 <sup>a</sup>	Update to EPA default
Alternative diet values	Not used	EPA default
Alternate source, subsistence food ( $\mu\text{g}/\text{day}$ )	1.6	Site data, see Table 5-8
Bioavailability of lead in food (percent)	50	EPA default
<b>Drinking Water</b>		
Lead concentration in drinking water ( $\mu\text{g}/\text{L}$ )	0.33	Site data
Drinking water intake (L/day)	0.20, 0.50, 0.52, 0.53, 0.55, 0.58, 0.59 <sup>a</sup>	EPA default
Alternative water values	Not used	EPA default
Bioavailability of lead in drinking water (percent)	50	EPA default
<b>Soil/Dust</b>		
Soil lead levels (ppm; $\mu\text{g}/\text{g}$ )	282	Site data <sup>b</sup>
Indoor dust lead levels (percent of soil levels)	70%	EPA default
Ingestion weighting factor (percent soil/percent dust)	45/55	EPA default
Amount of soil/dust ingested daily (g/day)	0.008, 0.012, 0.012, 0.012 0.009, 0.008, 0.008 <sup>a</sup>	EPA default <sup>c</sup>
Bioavailability of lead in soil and dust (percent)	30, 9.7	EPA default and site-specific <sup>d</sup>
<b>Other</b>		
Alternate source, subsistence food ( $\mu\text{g}/\text{day}$ )	1.6	Site data, see Table 5-8
Bioavailability of lead from subsistence foods (percent)	50	EPA default
Maternal contribution method	Infant model	EPA default
Maternal blood lead at birth of child ( $\mu\text{g}/\text{dL}$ )	2.5	EPA default
Geometric standard deviation	1.6	EPA default

**Note:** EPA - U.S. Environmental Protection Agency  
IEUBK - integrated exposure uptake/biokinetic

<sup>a</sup> Value varies by age group. Values listed are for the following ages, respectively: 0–1, 1–2, 2–3, 3–4, 4–5, 5–6, 6–7.

<sup>b</sup> The soil concentration is area weighted to account for the difference in concentrations in the port area and the road area.

<sup>c</sup> The soil ingestion rates were derived by multiplying the EPA model default age-specific soil ingestion rates (0.085, 0.135, 0.135, 0.135, 0.100, 0.090, and 0.085 g/day) by the site fractional intake of 0.09.

<sup>d</sup> The EPA default for the IEUBK lead model is 30 percent. The site-specific value is 9.7 percent (see Table 5-7), based on data from the lead bioavailability study conducted by the National Toxicology Program and reported by the Alaska Division of Public Health (ADPH 2001; Arnold and Middaugh 2001; Arnold et al. 2003).

Table 5-7. Bioavailability of lead in Red Dog ore concentrate

Lead Concentration in Amended Food (mg/kg) <sup>a</sup>	Blood Lead ( $\mu\text{g/dL}$ )		Relative Bioavailability	Child Absolute Bioavailability <sup>b</sup>	Adult Absolute Bioavailability <sup>b</sup>
	Lead Acetate	Red Dog Concentrate			
0		5.05	--	--	--
10	16	4.32	27.0%	13.5%	5.4%
30	31.8	5.65	17.8%	8.9%	3.6%
100	84.8	11.5	13.6%	6.8%	2.7%
Average	--	--	19.4%	9.7%	3.9%

**Source:** ADPH (2001); Arnold and Middaugh (2001); Arnold et al. (2003)

**Note:** -- - not applicable

<sup>a</sup> Animals were fed a diet amended with either Red Dog ore concentrate or soluble lead acetate so that the animals' food had the specific lead concentrations listed.

<sup>b</sup> Absolute bioavailability is calculated by multiplying the relative bioavailability of Red Dog concentrate by the absolute bioavailability of lead acetate. The absolute bioavailability of lead acetate was assumed to be 50 percent for children and 20 percent for adults, per U.S. EPA (U.S. EPA 1994, 1996c) guidance.

**Table 5-8. Calculation of subsistence food lead intake for EPA IEUBK child lead model**

Scenario Timeframe: Current/Future  
 Exposure Medium: Food  
 Exposure Point: Subsistence Food  
 Receptor Population: Subsistence User  
 Receptor Age: Young child

Exposure Route	Food		EPC Value	EPC Units	Daily Food Intake <sup>a</sup>	Daily Food Intake Units	Average Daily Intake	Average Daily Intake Units
<b>Ingestion</b>								
	Caribou	Lead	195	µg/kg	7.6E-3	kg/day	1.5	µg/day
	Fish	Lead	10.2	µg/kg	5.6E-3	kg/day	0.06	µg/day
	Ptarmigan	Lead	69.3	µg/kg	9.0E-5	kg/day	0.006	µg/day
	Salmonberry	Lead	133	µg/kg	3.8E-4	kg/day	0.05	µg/day
	Sourdock	Lead	211	µg/kg	6.3E-5	kg/day	0.01	µg/day
<b>Total</b>							1.6	µg/day

**Note:** -- - not applicable  
 EPA - U.S. Environmental Protection Agency  
 IEUBK - integrated exposure uptake biokinetic  
 EPC - exposure point concentration

<sup>a</sup> Derivation of intake rates presented in Table 5-11. See text for chronic daily intake algorithm.

**Table 5-9. Exposure assumptions used to calculate risk for non-lead metals for adults in the subsistence use scenario**

Scenario Timeframe: Current/Future  
 Receptor Population: Subsistence Use  
 Receptor Age: Adult

Exposure Medium and Route	Parameter Code	Parameter Definition	Units	Value	Rationale/Reference	Intake Equation/Model Name
<b>Soil Ingestion</b>						
	C <sub>S</sub>	Chemical concentration in soil	mg/kg	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_S \times CF \times IR_S \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/mg	0.000001	--	
	IR <sub>S</sub>	Ingestion rate	mg soil/day	100	DEC (2002)	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	200	DEC (2002)	
	ED	Exposure duration	years	30	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	10,950	DEC (2002)	
<b>Water Ingestion</b>						
	C <sub>W</sub>	Chemical concentration in surface water	μg/L	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_W \times CF \times IR_W \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	mg/μg	0.001	--	
	IR <sub>W</sub>	Ingestion rate	L/day	2	DEC (2002)	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	30	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	10,950	DEC (2002)	
<b>Food Ingestion</b>						
	C <sub>F</sub>	Chemical concentration in food <sup>b</sup>	mg/kg-wet wt.	see Table 5-2	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_F \times CR_F \times CF \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/g	0.001	--	
	CR <sub>F</sub>	Ingestion rate <sup>b</sup>	g/day	see Table 5-11	DFG (2001a)	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	30	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	10,950	DEC (2002)	

**Note:** -- - not applicable  
 RME - reasonable maximum exposure

<sup>a</sup> Based on a calculation of the fraction of the assumed subsistence use area on the site divided by the total subsistence use areas for Kivalina and Noatak (see Figures 5-2 and 5-3 and Section 5.2.1.1).

<sup>b</sup> A separate calculation is done for each food item.



**Table 5-10. Exposure assumptions used to calculate risk for non-lead metals for children in the subsistence use scenario**

Scenario Timeframe: Current/Future
Receptor Population: Subsistence Use
Receptor Age: Child

Exposure Medium and Route	Parameter Code	Parameter Definition	Units	Value	Rationale/Reference	Intake Equation/Model Name
<b>Soil Ingestion</b>						
	C <sub>S</sub>	Chemical concentration in soil	mg/kg	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_S \times CF \times IR_S \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/mg	0.000001	--	
	IR <sub>S</sub>	Ingestion rate	mg soil/day	200	DEC (2002)	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	200	DEC (2002)	
	ED	Exposure duration	years	6	DEC (2002)	
	BW	Body weight	kg	15	DEC (2002)	
	AT	Averaging time	days	2,190	DEC (2002)	
<b>Water Ingestion</b>						
	C <sub>W</sub>	Chemical concentration in surface water	µg/L	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_W \times CF \times IR_W \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	mg/µg	0.001	--	
	IR <sub>W</sub>	Ingestion rate	L/day	1	?	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	6	DEC (2002)	
	BW	Body weight	kg	15	DEC (2002)	
	AT	Averaging time	days	2,190	DEC (2002)	
<b>Food Ingestion</b>						
	C <sub>F</sub>	Chemical concentration in food <sup>b</sup>	mg/kg-wet wt.	see Table 5-2	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_F \times CR_F \times CF \times FI \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/g	0.001	--	
	CR <sub>F</sub>	Ingestion rate <sup>b</sup>	g/day	see Table 5-11	DFG (2001a)	
	FI	Fractional intake from site	unitless	0.09	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	6	DEC (2002)	
	BW	Body weight	kg	15	DEC (2002)	
	AT	Averaging time	days	2,190	DEC (2002)	

**Note:** -- - not applicable  
RME - reasonable maximum exposure

<sup>a</sup> Based on a calculation of the fraction of the assumed subsistence use area on the site divided by the total subsistence use areas for Kivalina and Noatak (see Figures 5-2 and 5-3 and Section 5.2.1.1).

<sup>b</sup> A separate calculation is done for each food item.

Table 5-11. Estimated subsistence food consumption rates

	Mean per Capita Consumption (g/day)			Caloric Intake Weighted Mean per Capita Consumption (g/day)	
	Kivalina	Noatak	Average of two villages	Adult	Child
<b>Land Mammals</b>	212.1	305.8	259.0	168	84
Caribou <sup>a</sup>	177.5	300.6	239.1	155	78
Moose	70.0	36.9	53.4	35	17
<b>Migratory Birds</b>	10.6	9.9	10.3	6.7	3.3
<b>Game Birds</b>	3.1	3.1	3.1	2.0	1.0
Ptarmigan <sup>a</sup>	3.1	3.1	3.1	2.0	1.0
<b>All Fish</b>	314.8	248.7	281.7	183	91
Salmon	29.2	216.1	122.6	80	40
Non-salmon fish <sup>a</sup>	296.4	85.0	190.7	124	62
Char	252.3	57.7	155.0	101	50
White fish	28.2	36.0	32.1	21	10
Cod	24.8	1.1	12.9	8.4	4.2
<b>Marine Invertebrates</b>	1.8	3.8	2.8	1.8	0.9
Clams	0.0	1.3	0.6	0.4	0.2
Crabs	0.8	6.4	3.6	2.3	1.2
Shrimp	1.6	0.0	0.8	0.5	0.3
<b>Marine Mammals</b>	415.1	106.0	260.6	169	85
Seal	251.8	101.6	176.7	115	57
Walrus	101.1	52.9	77.0	50	25
Whale	89.8	20.2	55.0	36	18
<b>Vegetation</b>	18.3	7.5	12.9	8.4	4.2
Berries <sup>a</sup>	17.5	8.2	12.9	8.4	4.2
Plants/greens/mushrooms <sup>a</sup>	1.5	2.5	2.0	1.3	0.7
Sum of Main Categories	976	685	830	539	270
Total kcal/day (@5.1 kcal/g)	4,977	3,492	4,234	2,750	1,375
Caloric Intake Weighting Factor	--	--	--	0.65	0.32

**Note:** Data from Community Profile Database (DFG 2001a). Kivalina data are from 1992. Noatak data are from 1994.

The sum of consumption rates for individual food items, or for sub-categories within a category, does not equal the consumption rate for the entire category in the database. For example, the sum of salmon and non-salmon fish consumption does not equal all fish consumption. This could be an artifact of the statistical methods used to derive consumption rates for entire categories based on data for individual items.

-- - not applicable

EPC - exposure point concentration

<sup>a</sup>Consumption rates for ptarmigan and non-salmon fish were used to derive risk estimates using EPCs for those foods. Consumption of land mammals was evaluated using EPCs for caribou. Consumption of all berries was evaluated using EPCs for salmonberries. Consumption of all plants, greens, and mushrooms was evaluated based on EPCs for sourdock.

**Table 5-12. Daily dietary intake of Alaska native adults**

	Males		Females	
	grams	kcal	grams	kcal
Protein	127	508	90	360
Fat	117	1,053	81	729
Carbohydrates	282	1,128	214	856
Total Energy <sup>a</sup>	526	2,689	385	1,945
Average kcal/g		5.1		5.1

**Source:** Nobmann et al. (1992)

**Note:** kcal - kilocalories; commonly called calories. Caloric intake was calculated by multiplying the intake in grams from Nobmann et al. (1992) by the number of kcal/g in each energy source: protein, 4 kcal/g; fat, 9 kcal/g; carbohydrate, 4 kcal/g

<sup>a</sup> The total energy estimates differ slightly from the values reported by Nobmann et al. (1992) (i.e., 2,750 kcal for males and 1,950 kcal for females), likely because of the standard rounding used for the specific energy content of protein, fat, and carbohydrates. The values calculated here are used solely for the purpose of calculating the average caloric density of the diet.

**Table 5-13. Adult lead model exposure parameters**

Scenario Timeframe: Current/Future  
 Receptor Population: Worker/Subsistence User  
 Receptor Age: Adult

Parameter Code	Parameter Definition	Units	Input Parameters	Rationale
C <sub>S</sub>	Soil lead concentration average	μg/g or ppm	282	see Table 5-1
R <sub>fetal/maternal</sub>	Fetal/maternal PbB ratio	--	0.9	EPA default
BKSF	Biokinetic slope factor	μg/dL per μg/day	0.4	EPA default
GSD <sub>i</sub>	Geometric standard deviation PbB	--	2.1	U.S. EPA (2002a)
PbB <sub>0</sub>	Baseline PbB	μg/dL	1.53	U.S. EPA (2002a)
IR <sub>S_W</sub>	Soil ingestion rate while at work (including soil and dust)	g/day	0.050	DEC (2004a)
IR <sub>S_S</sub>	Soil ingestion rate during subsistence activities (including soil and dust)	g/day	0.100	DEC (2004a)
AF <sub>S</sub>	Absorption fraction	--	0.12 and 0.039	EPA default, site specific
EF <sub>S</sub>	Exposure frequency	days/year	200	DEC (2002)
FI <sub>S_W</sub>	Fractional intake for soil ingestion while at work	--	0.67	Site specific
FI <sub>S_S</sub>	Fractional intake for soil ingestion during subsistence activities	--	0.03	Site specific
IR <sub>F</sub>	Daily lead intake from subsistence foods (see Table 5-8)	g/day	3.2	see Table 5-14
AF <sub>F</sub>	Absorption fraction for food	--	0.20	U.S. EPA (1994, 1996c)
EF <sub>F</sub>	Exposure frequency for food	days/year	182.5	Site specific
AT	Averaging time	days/year	365	365

**Note:** -- - not applicable  
 EPA - U.S. Environmental Protection Agency  
 PbB - blood lead

**Table 5-14. Calculator of subsistence food lead intake for adult lead model**

Scenario Timeframe: Current/Future  
 Exposure Medium: Food  
 Exposure Point: Subsistence Food  
 Receptor Population: Subsistence User  
 Receptor Age: Adult

Exposure Route	Food		EPC Value	EPC Units	Daily Food Intake	Daily Food Intake Units	Average Daily Intake	Average Daily Intake Units
<b>Ingestion</b>								
	Caribou	Lead	195	µg/kg	1.5E-2	kg/day	2.9	µg/day
	Fish	Lead	10.2	µg/kg	1.1E-2	kg/day	0.11	µg/day
	Ptarmigan	Lead	69.3	µg/kg	1.8E-4	kg/day	0.012	µg/day
	Salmonberry	Lead	133	µg/kg	7.6E-4	kg/day	0.10	µg/day
	Sourdock	Lead	211	µg/kg	1.2E-4	kg/day	0.02	µg/day
<b>Total</b>							3.2	µg/day

**Note:** EPC - exposure point concentration

**Table 5-15. Exposure assumptions used to calculate risk for non-lead metals for adults in the combined worker/ subsistence user scenario**

Scenario Timeframe: Current/Future  
 Receptor Population: Combined Worker/Subsistence Use  
 Receptor Age: Adult

Exposure Medium and Route	Parameter Code	Parameter Definition	Units	Value	Rationale/Reference	Intake Equation/Model Name
<b>Soil Ingestion</b>						
	C <sub>s</sub>	Chemical concentration in soil	mg/kg	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_s \times CF \times ((IR_{s_w} \times FI_{s_w}) + (IR_{s_s} \times FI_{s_s})) \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/mg	0.000001	--	
	IR <sub>s_w</sub>	Ingestion rate	mg soil/day	50	DEC (2004a)	
	IR <sub>s_s</sub>	Ingestion rate	mg soil/day	100	DEC (2004a)	
	FI <sub>s_w</sub>	Fractional intake from site	unitless	0.67	Area calculated <sup>a</sup>	
	FI <sub>s_s</sub>	Fractional intake from site	unitless	0.03	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	200	DEC (2002)	
	ED	Exposure duration	years	25	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	9,125	DEC (2002)	
<b>Water Ingestion</b>						
	C <sub>w</sub>	Chemical concentration in surface water	µg/L	see Table 5-1	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_w \times CF \times IR_w \times FI_{wf} \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	mg/µg	0.001	--	
	IR <sub>w</sub>	Ingestion rate	L/day	2	DEC (2002)	
	FI <sub>ww</sub>	Fractional intake from site	unitless	0.045	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	25	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	9,125	DEC (2002)	
<b>Food Ingestion</b>						
	C <sub>f</sub>	Chemical concentration in food <sup>b</sup>	mg/kg-wet wt.	see Table 5-2	--	Chronic Daily Intake (CDI) (mg/kg-day) = $C_f \times IR_f \times CF \times FI_{wf} \times EF \times ED / (BW \times AT)$
	CF	Conversion factor	kg/g	0.001	--	
	IR <sub>f</sub>	Ingestion rate <sup>b</sup>	g/day	see Table 5-11	DFG (2001a)	
	FI <sub>wf</sub>	Fractional intake from site	unitless	0.045	Area calculated <sup>a</sup>	
	EF	Exposure frequency	days/year	365	DEC (2002)	
	ED	Exposure duration	years	25	DEC (2002)	
	BW	Body weight	kg	70	DEC (2002)	
	AT	Averaging time	days	9,125	DEC (2002)	

**Note:** -- - not applicable

RME - reasonable maximum exposure

<sup>a</sup> Based on a calculation of the fraction of the total subsistence use area comprised of the site, combined with the relative amount of time individuals spend at work vs. off work (see Section 5.2.3.2).

<sup>b</sup> A separate calculation is done for each food item.

Table 5-16. Noncancer toxicity data—oral reference doses

Chemical of Concern	Oral Chronic RfD (mg/kg-day)	Primary Target Organ or System	Uncertainty Factor	Source	Date RfD Accessed
<b>Inorganics</b>					
Antimony	0.0004	Longevity; metabolic	1,000	IRIS	1/15/05
Barium	0.07	Hypertension; kidney <sup>a</sup>	3	IRIS	1/15/05
Cadmium (food and soil)	0.001	Kidney	10	IRIS	1/15/05
Cadmium (water)	0.0005	Kidney	10	IRIS	1/15/05
Lead	NA	NA	NA	NA	1/15/05
Thallium	0.00008	Liver enzymes	3,000	IRIS	1/15/05
Zinc	0.3	Iron and copper status	3	IRIS	1/15/05

**Note:** IRIS - Integrated Risk Information System

NA - not applicable

RfD - reference dose

<sup>a</sup> No adverse effects were observed in the studies on which the RfD is based.

Table 5-17. Results for adult lead model

Scenario Timeframe: Current  
 Exposure Medium: Surface soil and foods  
 Exposure Point: DMTS surface soil and subsistence foods  
 Receptor Population: Combined worker/subsistence user  
 Receptor Age: Adult

Exposure Variable	Description of Exposure Variable	Units	Default Bioavailability	Site-Specific Bioavailability
$C_s$	Soil lead concentration average	$\mu\text{g/g}$ or ppm	282	282
$R_{\text{fetal/maternal}}$	Fetal/maternal PbB ratio	--	0.9	0.9
BKSF	Biokinetic slope factor	$\mu\text{g/dL}$ per $\mu\text{g/day}$	0.4	0.4
$GSD_i$	Geometric standard deviation PbB	--	2.1	2.1
$PbB_0$	Baseline PbB	$\mu\text{g/dL}$	1.53	1.53
$IR_{S\_W}$	Soil ingestion rate while at work (including soil and dust)	g/day	0.050	0.050
$IR_{S\_S}$	Soil ingestion rate during subsistence activities (including soil and dust)	g/day	0.100	0.100
$AF_s$	Absorption fraction	--	0.12	0.039
$EF_s$	Exposure frequency	days/year	200	200
$FI_{S\_W}$	Fractional intake for soil ingestion while at work	--	0.67	0.67
$FI_{S\_S}$	Fractional intake for soil ingestion during subsistence activities	--	0.03	0.03
$IR_f$	Daily lead intake from subsistence foods (see Table 5-8)	g/day	3.2	3.2
$AF_f$	Absorption fraction for food	--	0.20	0.20
$EF_f$	Exposure frequency for food	days/year	182.5	182.5
$AT_s$	Averaging time (same for soil and dust)	days/year	365	365
$PbB_{\text{adult}}$	PbB of adult worker, geometric mean	$\mu\text{g/dL}$	1.9	1.7
$PbB_{\text{fetal}}$	PbB among fetuses of adult workers, geometric mean	$\mu\text{g/dL}$	1.7	1.6
$PbB_{\text{fetal}, 0.95}$	95th percentile PbB among fetuses of adult workers	$\mu\text{g/dL}$	5.9	5.4
$PbB_t$	Target PbB level of concern (e.g., 10 $\mu\text{g/dL}$ )	$\mu\text{g/dL}$	10.0	10.0
$P(PbB_{\text{fetal}} > PbB_t)$	Probability that fetal PbB > $PbB_t$ , assuming lognormal distribution	%	0.9%	0.7%

**Note:**  $PbB_{\text{adult}} = PbB_0 + (BKSF \times ((C_s \times ((IR_{S\_W} \times FI_{S\_W}) + (IR_{S\_S} \times FI_{S\_S})) \times EF_s \times AF_s) + (IR_f \times EF_f \times AF_f))) / AT_s$

$PbB_{\text{fetal}, 0.95} = PbB_{\text{adult}} \times (GSD_i^{1.645} \times R)$

DMTS - DeLong Mountain Regional Transportation System

PbB - blood lead



**Table 5-18. Noncancer hazards for adult subsistence soil ingestion**

Scenario Timeframe: Current/Future  
 Exposure Medium: Surface Soil  
 Exposure Point: DMTS Area Weighted Surface Soil  
 Receptor Population: Subsistence User  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	6.46	mg/kg	4.6E-7	mg/kg-day	4.0E-4	mg/kg-day	0.001
	Barium	3,219	mg/kg	2.3E-4	mg/kg-day	7.0E-2	mg/kg-day	0.003
	Cadmium	10.8	mg/kg	7.6E-7	mg/kg-day	1.0E-3	mg/kg-day	0.0008
	Thallium	0.49	mg/kg	3.4E-8	mg/kg-day	8.0E-5	mg/kg-day	0.0004
	Zinc	1,399	mg/kg	9.9E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0003
<b>Total Hazard Index for All CoPCs</b>								0.006

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-19. Noncancer hazards for child subsistence soil ingestion**

Scenario Timeframe: Current/Future
Exposure Medium: Surface Soil
Exposure Point: DMTS Area Weighted Surface Soil
Receptor Population: Subsistence User
Receptor Age: Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	6.46	mg/kg	4.2E-6	mg/kg-day	4.0E-4	mg/kg-day	0.01
	Barium	3,219	mg/kg	2.1E-3	mg/kg-day	7.0E-2	mg/kg-day	0.03
	Cadmium	10.8	mg/kg	7.1E-6	mg/kg-day	1.0E-3	mg/kg-day	0.007
	Thallium	0.49	mg/kg	3.2E-7	mg/kg-day	8.0E-5	mg/kg-day	0.004
	Zinc	1,399	mg/kg	9.2E-4	mg/kg-day	3.0E-1	mg/kg-day	0.003
<b>Total Hazard Index for All CoPCs</b>								0.05

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-20. Noncancer hazards for adult subsistence surface water ingestion**

Scenario Timeframe: Current/Future  
 Exposure Medium: Surface Water  
 Exposure Point: Site Stream Surface Water  
 Receptor Population: Subsistence User  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>	Thallium	0.14	µg/L	2.2E-7	mg/kg-day	8.0E-5	mg/kg-day	0.003
<b>Total Hazard Index for All CoPCs</b>								0.003

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-21. Noncancer hazards for child subsistence surface water ingestion**

Scenario Timeframe: Current/Future  
 Exposure Medium: Surface Water  
 Exposure Point: Site Stream Surface Water  
 Receptor Population: Subsistence User  
 Receptor Age: Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Thallium	0.14	µg/L	8.5E-7	mg/kg-day	8.0E-5	mg/kg-day	0.01
<b>Total Hazard Index for All CoPCs</b>								<b>0.01</b>

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-22. Noncancer hazards for adult subsistence caribou consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Caribou
Exposure Point: Site Caribou
Receptor Population: Subsistence User
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	1.3	mg/kg	2.7E-4	mg/kg-day	7.0E-2	mg/kg-day	0.004
	Cadmium	0.22	mg/kg	4.7E-5	mg/kg-day	1.0E-3	mg/kg-day	0.05
	Zinc	36.8	mg/kg	8.0E-3	mg/kg-day	3.0E-1	mg/kg-day	0.03
<b>Total Hazard Index for All CoPCs</b>								<b>0.08</b>

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-23. Noncancer hazards for child subsistence caribou consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Caribou  
 Exposure Point: Site Caribou  
 Receptor Population: Subsistence User  
 Receptor Age: Young Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	1.3	mg/kg	6.4E-4	mg/kg-day	7.0E-2	mg/kg-day	0.01
	Cadmium	0.22	mg/kg	1.1E-4	mg/kg-day	1.0E-3	mg/kg-day	0.1
	Zinc	36.8	mg/kg	1.9E-2	mg/kg-day	3.0E-1	mg/kg-day	0.06
<b>Total Hazard Index for All CoPCs</b>								0.2

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-24. Noncancer hazards for adult subsistence fish consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Fish
Exposure Point: Site Fish
Receptor Population: Subsistence User
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>	Thallium	0.0026	mg/kg	4.2E-7	mg/kg-day	8.0E-5	mg/kg-day	0.005
<b>Total Hazard Index for All CoPCs</b>								0.005

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-25. Noncancer hazards for child subsistence fish consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Fish
Exposure Point: Site Fish
Receptor Population: Subsistence User
Receptor Age: Young Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>	Thallium	0.0026	mg/kg	9.7E-7	mg/kg-day	8.0E-5	mg/kg-day	0.01
<b>Total Hazard Index for All CoPCs</b>								0.01

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).



**Table 5-26. Noncancer hazards for adult subsistence ptarmigan consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Ptarmigan  
 Exposure Point: Site Ptarmigan  
 Receptor Population: Subsistence User  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.52	mg/kg	1.3E-6	mg/kg-day	7.0E-2	mg/kg-day	0.00002
	Cadmium	3.5	mg/kg	9.1E-6	mg/kg-day	1.0E-3	mg/kg-day	0.009
	Zinc	15.7	mg/kg	4.0E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0001
<b>Total Hazard Index for All CoPCs</b>								0.009

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-27. Noncancer hazards for child subsistence ptarmigan consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Ptarmigan  
 Exposure Point: Site Ptarmigan  
 Receptor Population: Subsistence User  
 Receptor Age: Young Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.52	mg/kg	3.1E-6	mg/kg-day	7.0E-2	mg/kg-day	0.00004
	Cadmium	3.5	mg/kg	2.1E-5	mg/kg-day	1.0E-3	mg/kg-day	0.02
	Zinc	15.7	mg/kg	9.4E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0003
<b>Total Hazard Index for All CoPCs</b>								0.02

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-28. Noncancer hazards for adult subsistence berry consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Berries
Exposure Point: Site Salmonberries
Receptor Population: Subsistence User
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.078	mg/kg	8.4E-7	mg/kg-day	7.0E-2	mg/kg-day	0.00001
	Cadmium	0.049	mg/kg	5.2E-7	mg/kg-day	1.0E-3	mg/kg-day	0.001
	Zinc	4.7	mg/kg	5.1E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0002
<b>Total Hazard Index for All CoPCs</b>								0.001

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-29. Noncancer hazards for child subsistence berry consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Berries
Exposure Point: Site Salmonberries
Receptor Population: Subsistence User
Receptor Age: Young Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.078	mg/kg	2.0E-6	mg/kg-day	7.0E-2	mg/kg-day	0.00003
	Cadmium	0.049	mg/kg	1.2E-6	mg/kg-day	1.0E-3	mg/kg-day	0.001
	Zinc	4.7	mg/kg	1.2E-4	mg/kg-day	3.0E-1	mg/kg-day	0.0004
<b>Total Hazard Index for All CoPCs</b>								0.002

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-30. Noncancer hazards for adult subsistence sourdock consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Sourdock  
 Exposure Point: Site Sourdock  
 Receptor Population: Subsistence User  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	0.012	mg/kg	2.1E-8	mg/kg-day	4.0E-4	mg/kg-day	0.00005
	Barium	10.6	mg/kg	1.8E-5	mg/kg-day	7.0E-2	mg/kg-day	0.0003
	Cadmium	0.013	mg/kg	2.2E-8	mg/kg-day	1.0E-3	mg/kg-day	0.00002
	Thallium	0.00049	mg/kg	8.2E-10	mg/kg-day	8.0E-5	mg/kg-day	0.00001
	Zinc	5.4	mg/kg	9.0E-6	mg/kg-day	3.0E-1	mg/kg-day	0.00003
<b>Total Hazard Index for All CoPCs</b>								0.0004

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-31. Noncancer hazards for child subsistence sourdock consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Sourdock
Exposure Point: Site Sourdock
Receptor Population: Subsistence User
Receptor Age: Young Child

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	0.012	mg/kg	5.2E-8	mg/kg-day	4.0E-4	mg/kg-day	0.0001
	Barium	10.6	mg/kg	4.5E-5	mg/kg-day	7.0E-2	mg/kg-day	0.001
	Cadmium	0.013	mg/kg	5.5E-8	mg/kg-day	1.0E-3	mg/kg-day	0.00005
	Thallium	0.00049	mg/kg	2.1E-9	mg/kg-day	8.0E-5	mg/kg-day	0.00003
	Zinc	5.4	mg/kg	2.3E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0001
<b>Total Hazard Index for All CoPCs</b>								0.001

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-32. Noncancer hazards for adult DMTS worker/subsistence user soil ingestion**

Scenario Timeframe: Current/Future  
 Exposure Medium: Surface Soil  
 Exposure Point: DMTS Area Weighted Surface Soil  
 Receptor Population: Worker/Subsistence  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	6.5	mg/kg	1.8E-6	mg/kg-day	4.0E-4	mg/kg-day	0.005
	Barium	3,219	mg/kg	9.2E-4	mg/kg-day	7.0E-2	mg/kg-day	0.01
	Cadmium	10.8	mg/kg	3.1E-6	mg/kg-day	1.0E-3	mg/kg-day	0.003
	Thallium	0.49	mg/kg	1.4E-7	mg/kg-day	8.0E-5	mg/kg-day	0.002
	Zinc	1,399	mg/kg	4.0E-4	mg/kg-day	3.0E-1	mg/kg-day	0.001
<b>Total Hazard Index for All CoPCs</b>								0.02

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-33. Noncancer hazards for adult DMTS worker/subsistence user surface water ingestion**

Scenario Timeframe: Current/Future  
 Exposure Medium: Stream Surface Water  
 Exposure Point: Site Stream Surface Water  
 Receptor Population: Worker/Subsistence  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>	Thallium	0.14	µg/L	1.8E-7	mg/kg-day	8.0E-5	mg/kg-day	0.002
<b>Total Hazard Index for All CoPCs</b>								0.002

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).



**Table 5-34. Noncancer hazards for adult DMTS worker/subsistence user caribou consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Caribou
Exposure Point: Site Caribou
Receptor Population: Worker/Subsistence
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	1.3	mg/kg	1.4E-4	mg/kg-day	7.0E-2	mg/kg-day	0.002
	Cadmium	0.22	mg/kg	2.3E-5	mg/kg-day	1.0E-3	mg/kg-day	0.02
	Zinc	36.8	mg/kg	4.0E-3	mg/kg-day	3.0E-1	mg/kg-day	0.01
<b>Total Hazard Index for All CoPCs</b>								0.04

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-35. Noncancer hazards for adult DMTS worker/subsistence user fish consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Fish
Exposure Point: Site Fish
Receptor Population: Worker/Subsistence
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>	Thallium	0.0026	mg/kg	2.1E-7	mg/kg-day	8.0E-5	mg/kg-day	0.003
<b>Total Hazard Index for All CoPCs</b>								0.003

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-36. Noncancer hazards for adult DMTS worker/subsistence user ptarmigan consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Ptarmigan  
 Exposure Point: Site Ptarmigan  
 Receptor Population: Worker/Subsistence  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.52	mg/kg	6.6E-7	mg/kg-day	7.0E-2	mg/kg-day	0.00001
	Cadmium	3.5	mg/kg	4.5E-6	mg/kg-day	1.0E-3	mg/kg-day	0.005
	Zinc	15.7	mg/kg	2.0E-5	mg/kg-day	3.0E-1	mg/kg-day	0.00007
<b>Total Hazard Index for All CoPCs</b>								0.005

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-37. Noncancer hazards for adult DMTS worker/subsistence user berry consumption**

Scenario Timeframe: Current/Future  
 Exposure Medium: Berries  
 Exposure Point: Site Salmonberries  
 Receptor Population: Worker/Subsistence  
 Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Barium	0.078	mg/kg	4.2E-7	mg/kg-day	7.0E-2	mg/kg-day	0.00001
	Cadmium	0.049	mg/kg	2.6E-7	mg/kg-day	1.0E-3	mg/kg-day	0.000
	Zinc	4.7	mg/kg	2.5E-5	mg/kg-day	3.0E-1	mg/kg-day	0.0001
<b>Total Hazard Index for All CoPCs</b>								0.0004

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

**Table 5-38. Noncancer hazards for adult DMTS worker/subsistence user sourdock consumption**

Scenario Timeframe: Current/Future
Exposure Medium: Sourdock
Exposure Point: Site Sourdock
Receptor Population: Worker/Subsistence
Receptor Age: Adult

Exposure Route	CoPC	EPC Value <sup>a</sup>	EPC Units	Intake	Intake Units	Reference Dose <sup>b</sup>	Reference Dose Units	Hazard Quotient
<b>Ingestion</b>								
	Antimony	0.012	mg/kg	1.0E-8	mg/kg-day	4.0E-4	mg/kg-day	0.00003
	Barium	10.6	mg/kg	8.8E-6	mg/kg-day	7.0E-2	mg/kg-day	0.0001
	Cadmium	0.013	mg/kg	1.1E-8	mg/kg-day	1.0E-3	mg/kg-day	0.00001
	Thallium	0.00049	mg/kg	4.1E-10	mg/kg-day	8.0E-5	mg/kg-day	0.00001
	Zinc	5.39	mg/kg	4.5E-6	mg/kg-day	3.0E-1	mg/kg-day	0.00001
<b>Total Hazard Index for All CoPCs</b>								0.0002

**Note:** CoPC - chemical of potential concern  
 DMTS - DeLong Mountain Regional Transportation System  
 EPA - U.S. Environmental Protection Agency  
 EPC - exposure point concentration  
 UCL - upper confidence limit

<sup>a</sup> Values for all chemicals reflect the lower of either the 95th percentile UCL on the mean or the maximum concentration.

<sup>b</sup> Toxicity values obtained from the EPA Integrated Risk Information System (IRIS) (January 2005).

Table 5-39. Summary of total hazard indices for reasonable maximum exposure scenarios

Receptor/Exposure Pathway	Adult		Young Child		Chemicals Accounting for 90 percent of Hazard Indices for each Pathway
	Hazard Index	% Contribution by Pathway	Hazard Index	% Contribution by Pathway	
<b>Subsistence User—Current/Future</b>					
Surface soil ingestion	0.006	6%	0.05	19%	Barium, antimony, cadmium, thallium
Water ingestion	0.003	3%	0.01	4%	Thallium
Caribou consumption	0.08	76%	0.2	64%	Cadmium, zinc
Fish consumption	0.005	5%	0.01	4%	Thallium
Ptarmigan consumption	0.009	9.1%	0.02	7.6%	Cadmium
Berry consumption	0.001	0.7%	0.002	0.6%	Cadmium, zinc
Sourdock consumption	0.0004	0.4%	0.001	0.3%	Barium, antimony, zinc
<b>Total for Subsistence User</b>	<b>0.1</b>	<b>100%</b>	<b>0.3</b>	<b>100%</b>	
<b>Worker—Current/Future</b>					
Surface soil ingestion	0.02	33%			Barium, antimony, cadmium, thallium
Water ingestion	0.002	3%			Thallium
Caribou consumption	0.04	53%			Cadmium, zinc
Fish consumption	0.003	4%			Thallium
Ptarmigan consumption	0.005	6.4%			Cadmium
Berry consumption	0.0004	0.5%			Cadmium, zinc
Sourdock consumption	0.0002	0.3%			Barium, antimony, zinc
<b>Total for DMTS Worker</b>	<b>0.07</b>	<b>100%</b>			

**Note:** DMTS - DeLong Mountain Regional Transportation System

**Table 6-1. Refined assessment endpoints, representative receptors, and measurement endpoints**

Environment	Assessment Endpoint	Representative Receptor <sup>a</sup>	Measurement Endpoint
Tundra	Structure and function of terrestrial plant communities	Terrestrial plant communities	Plant abundance, diversity, biomass, percent cover
Tundra	Structure and function of tundra soil fauna communities	Tundra soil fauna communities	Not directly assessed, evaluated through terrestrial plant community analysis
Tundra	Survival, growth, and reproduction of terrestrial avian herbivore populations	Willow ptarmigan	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to avian TRVs
Tundra	Survival, growth, and reproduction of terrestrial mammalian herbivore populations	Tundra vole; caribou; moose	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to mammalian TRVs
Tundra	Survival, growth, and reproduction of terrestrial avian invertivore populations	Lapland longspur	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to avian TRVs
Tundra	Survival, growth, and reproduction of terrestrial mammalian invertivore populations	Tundra shrew	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to mammalian TRVs
Tundra	Survival, growth, and reproduction of terrestrial avian carnivore populations	Snowy owl	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to avian TRVs
Tundra	Survival, growth, and reproduction of terrestrial mammalian carnivore populations	Arctic fox	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, soil, and surface water) relative to mammalian TRVs
Streams	Structure and function of stream aquatic and wetland plant communities	Stream aquatic and wetland plant communities	Plant abundance, diversity, biomass, percent cover
Streams	Structure and function of stream aquatic invertebrate communities	Stream aquatic invertebrate communities	Abundance and diversity of stream aquatic invertebrates
Streams	Survival, growth, and reproduction of stream avian herbivore populations	Green-winged teal	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to avian TRVs
Streams	Survival, growth, and reproduction of stream mammalian herbivore populations	Muskrat; moose	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to mammalian TRVs

8601997.001 4400 0105 SS18  
\\oswego2\data\docs\8601997.001 4400\dms\_fa\_la.doc

Table 6-1. (cont.)

Environment	Assessment Endpoint	Representative Receptor <sup>a</sup>	Measurement Endpoint
Streams	Survival, growth, and reproduction of stream avian invertivore populations	Common snipe	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to avian TRVs
Tundra ponds	Structure and function of tundra pond aquatic and wetland plant communities	Tundra pond aquatic and wetland plant communities	Plant abundance, diversity, biomass, percent cover
Tundra ponds	Structure and function of tundra pond aquatic invertebrate communities	Tundra pond aquatic invertebrate communities	Abundance and diversity of tundra pond aquatic invertebrates
Tundra ponds	Survival, growth, and reproduction of tundra pond avian herbivore populations	Green-winged teal	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to avian TRVs
Tundra ponds	Survival, growth, and reproduction of tundra pond mammalian herbivore populations	Muskrat	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to mammalian TRVs
Tundra ponds	Survival, growth, and reproduction of tundra pond avian invertivore populations	Common snipe <sup>b</sup>	Range of modeled total dietary exposures (based on measured CoPC concentrations in food, sediment, and surface water) relative to avian TRVs
Coastal lagoons	Structure and function of coastal lagoon aquatic and wetland plant communities	Coastal lagoon aquatic and wetland plant communities	Plant abundance, diversity, biomass, percent cover
Coastal lagoons	Structure and function of coastal lagoon aquatic invertebrate communities	Coastal lagoon aquatic invertebrate communities	Abundance and diversity of coastal lagoon aquatic invertebrates
Coastal lagoons	Survival, growth, and reproduction of coastal lagoon avian herbivore populations	Brant	Range of modeled total dietary exposures (based on measured CoPC concentrations in food and sediment) relative to avian TRVs
Coastal lagoons	Survival, growth, and reproduction of coastal lagoon avian invertivore populations	Black-bellied plover	Range of modeled total dietary exposures (based on measured CoPC concentrations in food and sediment) relative to avian TRVs

**Note:** CoPC - chemical of potential concern  
TRV - toxicity reference value

<sup>a</sup> Receptors to be evaluated in the risk assessment.

<sup>b</sup> Evaluated as a terrestrial receptor.



**Table 6-2. Cover classes for quantifying percent cover of plant species**

Cover Class	Range (percent)	Midpoint (percent)
+	Trace	0
1	<5	2.5
2	5–25	15
3	25–50	37.5
4	50–75	62.5
5	75–95	85
6	95–100	97.5

**Source:** Modified from Daubenmire (1959).

Table 6-3. Summary of *p*-values for site to reference comparison by vegetation community type

	Coastal		Tundra and Coastal				Hillslope	Lagoon
	Plain	Tundra	All	10 m <sup>a</sup>	100 m	1,000 m <sup>b</sup>		
Forb cover	0.429	0.371	0.930	0.108	0.543	0.237	0.655	0.439
Graminoid cover	0.157	0.505	1.000	0.480	0.248	1.000	0.655	0.439
Deciduous shrub cover	0.157	<b>0.096</b>	0.499	0.724	0.564	0.480	0.180	--
Evergreen shrub cover	0.157	0.739	0.271	<b>0.034</b>	1.000	0.858	0.180	--
Moss cover	0.480	<b>0.096</b>	<b>0.091</b>	0.289	<b>0.083</b>	0.157	0.180	0.439
Moss frequency	0.617	0.564	0.419	0.387	0.221	--	--	0.317
Lichen cover	0.157	<b>0.044</b>	<b>0.011</b>	<b>0.028</b>	<b>0.083</b>	<b>0.034</b>	0.180	0.317
Lichen frequency	0.277	0.153	<b>0.060</b>	<b>0.022</b>	<b>0.053</b>	0.387	0.157	0.317
Unvegetated cover	0.429	0.252	0.629	0.142	0.739	0.248	1.000	0.439
Diversity	0.480	0.739	0.866	0.480	1.000	0.724	0.655	0.439
Evenness	1.000	0.505	0.499	<b>0.077</b>	<b>0.083</b>	<b>0.034</b>	0.655	0.121
Richness	1.000	0.211	0.300	0.472	0.197	<b>0.028</b>	0.655	1.000
Antimony	<b>0.079</b>	<b>0.023</b>	<b>0.006</b>	<b>0.017</b>	<b>0.042</b>	<b>0.017</b>	<b>0.090</b>	<b>0.061</b>
Arsenic	<b>0.079</b>	0.251	<b>0.064</b>	<b>0.016</b>	0.282	0.238	0.327	<b>0.061</b>
Barium	0.234	0.159	0.118	<b>0.016</b>	<b>0.042</b>	0.760	<b>0.090</b>	<b>0.061</b>
Cadmium	<b>0.079</b>	<b>0.023</b>	<b>0.006</b>	<b>0.017</b>	<b>0.042</b>	<b>0.017</b>	<b>0.090</b>	<b>0.061</b>
Cobalt	<b>0.079</b>	0.909	0.433	0.144	0.718	0.638	0.673	0.500
Copper	<b>0.079</b>	0.369	0.118	<b>0.017</b>	0.124	0.638	<b>0.090</b>	<b>0.061</b>
Lead	<b>0.079</b>	<b>0.091</b>	<b>0.032</b>	<b>0.017</b>	<b>0.042</b>	0.240	<b>0.090</b>	<b>0.061</b>
Manganese	0.240	0.631	0.306	<b>0.039</b>	0.282	0.856	0.910	0.500
Mercury	<b>0.079</b>	<b>0.048</b>	<b>0.046</b>	<b>0.039</b>	0.124	0.144	<b>0.090</b>	<b>0.061</b>
Molybdenum	<b>0.079</b>	0.748	0.400	0.240	0.718	0.429	<b>0.090</b>	0.500
Selenium	<b>0.079</b>	0.120	<b>0.026</b>	<b>0.017</b>	0.124	0.106	<b>0.090</b>	<b>0.061</b>
Silver	<b>0.079</b>	<b>0.091</b>	<b>0.017</b>	<b>0.017</b>	<b>0.042</b>	0.106	<b>0.090</b>	<b>0.061</b>
Thallium	0.500	0.253	0.306	<b>0.017</b>	<b>0.042</b>	0.240	0.327	0.781
Vanadium	<b>0.079</b>	0.253	0.155	<b>0.079</b>	0.282	0.362	0.327	0.500
Zinc	0.240	<b>0.023</b>	<b>0.021</b>	<b>0.017</b>	<b>0.042</b>	0.144	<b>0.090</b>	<b>0.061</b>
pH	0.147	0.129	<b>0.032</b>	<b>0.032</b>	<b>0.083</b>	0.172	0.180	0.121
Total solids	1.000	0.505	0.612	<b>0.077</b>	1.000	0.480	0.655	0.439

**Note:** Metals compared using one-sided test for higher concentrations at site.

**Bold** values indicate a significant difference from reference,  $p < 0.10$ .

Wilcoxon test was used for comparisons between site and reference groups.

-- - comparison was not possible because there was no variability or no relevant species were observed

<sup>a</sup> Also includes Station TT5-0100, located approximately 85 m from the nearest dust source.

<sup>b</sup> Also includes Stations TT5-1000 and TT5-2000, located approximately 450 m and 1,430 m from the nearest dust source, respectively.

Table 6-4. Summary of parameter relationships with distance from DMTS road

	All Transects				Coastal and Tundra Transects only			
	Correlation		Linear Regression		Correlation		Linear Regression	
	Estimate	P-value	P-value	R-square	Estimate	P-value	P-value	R-square
Forb cover	<b>-0.494</b>	<b>0.085</b>	0.503	4.2%	<b>-0.710</b>	<b>0.032</b>	0.163	22.8%
Graminoid cover	0.186	0.526	0.427	5.8%	0.241	0.481	0.545	4.7%
Deciduous shrub cover	-0.211	0.458	0.343	8.2%	-0.321	0.326	0.206	19.2%
Evergreen shrub cover	<b>0.589</b>	<b>0.042</b>	<b>0.040</b>	<b>33.1%</b>	<b>0.741</b>	<b>0.027</b>	<b>0.010</b>	<b>58.6%</b>
Moss cover	0.285	0.329	0.354	7.9%	<b>0.593</b>	<b>0.078</b>	<b>0.051</b>	<b>39.7%</b>
Moss frequency	0.263	0.367	0.337	8.4%	0.310	0.361	0.321	12.3%
Lichen cover	<b>0.595</b>	<b>0.040</b>	0.144	18.3%	<b>0.994</b>	<b>0.003</b>	<b>&lt;0.001</b>	<b>77.0%</b>
Lichen frequency	<b>0.717</b>	<b>0.013</b>	<b>0.003</b>	<b>57.9%</b>	<b>0.911</b>	<b>0.007</b>	<b>&lt;0.001</b>	<b>95.5%</b>
Unvegetated cover	<b>-0.716</b>	<b>0.013</b>	<b>0.005</b>	<b>53.1%</b>	<b>-0.803</b>	<b>0.015</b>	<b>0.001</b>	<b>74.9%</b>
Diversity	0.307	0.292	0.140	18.7%	0.358	0.291	0.179	21.4%
Evenness	<b>0.631</b>	<b>0.029</b>	<b>0.013</b>	<b>44.3%</b>	<b>0.908</b>	<b>0.007</b>	<b>&lt;0.001</b>	<b>84.8%</b>
Richness	<b>-0.551</b>	<b>0.055</b>	0.667	1.7%	<b>-0.892</b>	<b>0.007</b>	<b>0.052</b>	<b>39.3%</b>
Antimony	<b>-0.654</b>	<b>0.023</b>	<b>0.017</b>	<b>41.9%</b>	<b>-0.809</b>	<b>0.015</b>	<b>0.021</b>	<b>50.8%</b>
Arsenic	<b>-0.897</b>	<b>0.002</b>	<b>&lt;0.001</b>	<b>72.9%</b>	<b>-0.950</b>	<b>0.004</b>	<b>&lt;0.001</b>	<b>84.9%</b>
Barium	<b>-0.684</b>	<b>0.017</b>	<b>0.013</b>	<b>44.7%</b>	<b>-0.836</b>	<b>0.012</b>	<b>0.008</b>	<b>60.5%</b>
Cadmium	<b>-0.673</b>	<b>0.019</b>	<b>0.018</b>	<b>41.1%</b>	<b>-0.833</b>	<b>0.012</b>	<b>0.013</b>	<b>55.7%</b>
Cobalt	<b>-0.823</b>	<b>0.004</b>	<b>0.003</b>	<b>57.9%</b>	<b>-0.803</b>	<b>0.015</b>	<b>0.017</b>	<b>52.7%</b>
Copper	<b>-0.769</b>	<b>0.007</b>	<b>0.002</b>	<b>58.4%</b>	<b>-0.914</b>	<b>0.006</b>	<b>&lt;0.001</b>	<b>78.2%</b>
Lead	<b>-0.817</b>	<b>0.005</b>	<b>0.004</b>	<b>53.9%</b>	<b>-0.840</b>	<b>0.011</b>	<b>0.006</b>	<b>63.4%</b>
Manganese	<b>-0.707</b>	<b>0.014</b>	<b>0.017</b>	<b>41.6%</b>	<b>-0.698</b>	<b>0.035</b>	<b>0.037</b>	<b>44.0%</b>
Mercury	-0.256	0.371	0.205	14.2%	-0.278	0.394	0.241	16.7%
Molybdenum	-0.042	0.876	0.880	0.22%	-0.084	0.788	0.750	1.3%
Selenium	<b>-0.504</b>	<b>0.079</b>	<b>0.054</b>	<b>29.6%</b>	<b>-0.678</b>	<b>0.040</b>	<b>0.034</b>	<b>44.9%</b>
Silver	<b>-0.685</b>	<b>0.017</b>	<b>0.010</b>	<b>46.4%</b>	<b>-0.895</b>	<b>0.007</b>	<b>0.006</b>	<b>64.0%</b>
Thallium	<b>-0.662</b>	<b>0.021</b>	<b>0.007</b>	<b>49.4%</b>	<b>-0.914</b>	<b>0.006</b>	<b>&lt;0.001</b>	<b>78.5%</b>
Vanadium	<b>-0.696</b>	<b>0.015</b>	<b>0.006</b>	<b>51.1%</b>	<b>-0.945</b>	<b>0.004</b>	<b>0.001</b>	<b>75.7%</b>
Zinc	<b>-0.817</b>	<b>0.005</b>	<b>0.008</b>	<b>49.1%</b>	<b>-0.796</b>	<b>0.016</b>	<b>0.011</b>	<b>57.9%</b>
pH	<b>-0.808</b>	<b>0.005</b>	<b>0.001</b>	<b>63.2%</b>	<b>-0.878</b>	<b>0.008</b>	<b>&lt;0.001</b>	<b>84.2%</b>
Total Solids	<b>-0.919</b>	<b>0.001</b>	<b>&lt;0.001</b>	<b>83.0%</b>	<b>-0.945</b>	<b>0.004</b>	<b>&lt;0.001</b>	<b>81.7%</b>

**Note:** Spearman rank non-parametric correlation was used.

Linear regression models related  $\log_{10}$  distance to each variable.  $\log_{10}$  transform of metals concentrations and total solids was used to better meet method assumptions of equal variability and normality.

**Bold** indicates significant relationship with distance ( $p < 0.10$ ).

Table 6-5. Summary of correlations among vegetation variables and soil parameters

	Antimony	Arsenic	Barium	Cadmium	Cobalt	Copper	Lead	Manganese	Mercury	Molybdenum	Selenium	Silver	Thallium	Vanadium	Zinc	pH	Total Solids
Distance	-0.650	-0.900	-0.680	-0.670	-0.820	-0.770	-0.820	-0.710			-0.500	-0.680	-0.660	-0.700	-0.820	-0.810	-0.920
Forb cover		0.600	0.390	0.430		0.660	0.510				0.660	0.540	0.530	0.600	0.400	0.510	
Graminoid cover	-0.440		-0.410	-0.410				-0.390	-0.400			-0.480	-0.520				
Deciduous shrub cover			0.510					0.640									
Evergreen shrub cover		-0.850		-0.470	-0.470	-0.720	-0.570			-0.370	-0.570	-0.530	-0.520	-0.730	-0.410	-0.430	
Moss cover																	
Moss frequency	-0.460			-0.370			-0.410					-0.460			-0.410		
Lichen cover	-0.430	-0.420		-0.570		-0.390	-0.570				-0.440	-0.430			-0.590	-0.500	
Lichen frequency		-0.710		-0.510	-0.510	-0.590	-0.670				-0.610	-0.540	-0.470	-0.680	-0.620	-0.700	-0.400
Unvegetated cover		0.440				0.380								0.520	0.540	0.600	
Diversity		-0.620			-0.370									-0.570			
Evenness		-0.750	-0.490	-0.390	-0.530	-0.780	-0.540			-0.460	-0.450	-0.520	-0.640	-0.710	-0.460	-0.430	-0.430
Richness			0.540				0.670					0.420	0.550				0.480
Antimony	1		0.700	0.920		0.680	0.830	0.390	0.790		0.700	0.910	0.790		0.860	0.690	
Arsenic		1	0.420	0.450	0.690	0.840	0.630			0.440	0.690	0.560	0.590	0.890	0.490	0.550	0.420
Barium	0.700	0.420	1	0.660		0.730	0.740	0.700			0.580	0.740	0.850	0.400	0.770	0.820	0.580
Cadmium	0.920	0.450	0.660	1		0.770	0.890	0.410	0.760		0.750	0.940	0.870		0.860	0.750	0.410
Cobalt		0.690			1	0.640	0.440				0.450	0.430	0.430	0.580			0.490
Copper	0.680	0.840	0.730	0.770	0.640	1	0.840	0.450	0.470	0.460	0.810	0.850	0.900	0.700	0.730	0.750	0.490
Lead	0.830	0.630	0.740	0.890	0.440	0.840	1	0.530	0.580		0.750	0.900	0.870	0.450	0.930	0.800	0.530
Manganese	0.390		0.700	0.410		0.450	0.530	1				0.470	0.620		0.490	0.560	0.480
Mercury	0.790			0.760		0.470	0.580		1		0.470	0.670	0.530		0.580	0.410	
Molybdenum		0.440				0.460				1	0.410			0.510			
Selenium	0.700	0.690	0.580	0.750	0.450	0.810	0.750		0.470	0.410	1	0.790	0.710	0.670	0.710	0.770	
Silver	0.910	0.560	0.740	0.940	0.430	0.850	0.900	0.470	0.670		0.790	1	0.920	0.430	0.830	0.790	0.470
Thallium	0.790	0.590	0.850	0.870	0.430	0.900	0.870	0.620	0.530		0.710	0.920	1	0.480	0.790	0.800	0.600
Vanadium		0.890	0.400		0.580	0.700	0.450			0.510	0.670	0.430	0.480	1		0.550	
Zinc	0.860	0.490	0.770	0.860		0.730	0.930	0.490	0.580		0.710	0.830	0.790		1	0.790	0.490
pH	0.690	0.550	0.820	0.750		0.750	0.800	0.560	0.410		0.770	0.790	0.800	0.550	0.790	1	0.460
Total solids		0.420	0.580	0.410	0.490	0.490	0.530	0.480				0.470	0.600		0.490	0.460	1

**Note:** Only significant ( $p < 0.10$ ) correlation estimates are reported.  
Spearman rank non-parametric correlation was used to estimate correlations.

Table 6-6. Summary of correlations among vegetation variables

	Distance	Diversity	Evenness	Richness	Deciduous shrub cover	Evergreen shrub cover	Forb cover	Graminoid cover	Lichen cover	Lichen frequency	Moss cover	Moss frequency	Unvegetated cover
Distance	1		0.630	-0.550		0.590	-0.490		0.600	0.720			-0.720
Diversity		1	0.450	0.580	0.370	0.460				0.470	0.470		
Evenness	0.630	0.450	1			0.650	-0.600			0.520			-0.450
Richness	-0.550	0.580		1	0.650			-0.630			0.460		
Deciduous shrub cover		0.370		0.650	1			-0.690	0.420				
Evergreen shrub cover	0.590	0.460	0.650			1	-0.660		0.400	0.650			
Forb cover	-0.490		-0.600			-0.660	1			-0.470			
Graminoid cover				-0.630	-0.690			1					
Lichen cover	0.600				0.420	0.400			1	0.750	0.720	0.390	-0.380
Lichen frequency	0.720	0.470	0.520			0.650	-0.470		0.750	1	0.550	0.400	-0.630
Moss cover		0.470		0.460					0.720	0.550	1	0.570	
Moss frequency									0.390	0.400	0.570	1	
Unvegetated cover	-0.720		-0.450						-0.380	-0.630			1
Antimony	-0.650							-0.440	-0.430			-0.460	
Arsenic	-0.900	-0.620	-0.750			-0.850	0.600		-0.420	-0.710			0.440
Barium	-0.680		-0.490	0.540	0.510		0.390	-0.410					
Cadmium	-0.670		-0.390			-0.470	0.430	-0.410	-0.570	-0.510		-0.370	
Cobalt	-0.820	-0.370	-0.530			-0.470				-0.510			
Copper	-0.770		-0.780			-0.720	0.660		-0.390	-0.590			0.380
Lead	-0.820		-0.540			-0.570	0.510		-0.570	-0.670		-0.410	
Manganese	-0.710			0.670	0.640			-0.390					
Mercury								-0.400					
Molybdenum			-0.460			-0.370							
Selenium	-0.500		-0.450			-0.570	0.660		-0.440	-0.610			
Silver	-0.680		-0.520	0.420		-0.530	0.540	-0.480	-0.430	-0.540		-0.460	
Thallium	-0.660		-0.640	0.550		-0.520	0.530	-0.520		-0.470			
Vanadium	-0.700	-0.570	-0.710			-0.730	0.600			-0.680			0.520
Zinc	-0.820		-0.460			-0.410	0.400		-0.590	-0.620		-0.410	
pH	-0.810		-0.430			-0.430	0.510		-0.500	-0.700			0.540
Total solids	-0.920		-0.430	0.480						-0.400			0.600

**Note:** Only significant ( $p < 0.10$ ) correlation estimates are reported.

Spearman rank non-parametric correlation was used to estimate correlations.

Table 6-7. Summary of correlations among vegetation variables and soil parameters at coastal plain and tundra communities

	Antimony	Arsenic	Barium	Cadmium	Cobalt	Copper	Lead	Manganese	Mercury	Molybdenum	Selenium	Silver	Thallium	Vanadium	Zinc	pH	Total Solids
Distance	-0.809	-0.950	-0.836	-0.833	-0.803	-0.914	-0.840	-0.698			-0.678	-0.895	-0.914	-0.945	-0.796	-0.878	-0.945
Forb cover		0.618	0.617	0.502	0.576	0.714	0.634	0.694			0.548	0.541	0.634	0.551	0.528	0.497	
Graminoid cover					-0.484												
Deciduous shrub cover									-0.500								
Evergreen shrub cover	-0.580	-0.764		-0.759	-0.649	-0.724	-0.732				-0.487	-0.748	-0.790	-0.547	-0.558	-0.475	-0.707
Moss cover				-0.478			-0.500					-0.531	-0.500			-0.491	-0.555
Moss frequency																	
Lichen cover	-0.895	-0.778	-0.741	-0.912		-0.773	-0.834	-0.575	-0.503		-0.753	-0.877	-0.873	-0.657	-0.807	-0.912	-0.602
Lichen frequency	-0.866	-0.790	-0.836	-0.832		-0.768	-0.922	-0.733	-0.486		-0.717	-0.892	-0.861	-0.692	-0.910	-0.905	-0.750
Unvegetated cover	0.480	0.489	0.611			0.493	0.524	0.527				0.512	0.493	0.527	0.540	0.605	0.662
Diversity														-0.571			-0.544
Evenness		-0.664	-0.699		-0.604	-0.676	-0.588	-0.643				-0.534	-0.577	-0.681	-0.549		-0.808
Richness						0.575		0.533									0.525
Antimony	1.000	0.821	0.680	0.934	0.484	0.802	0.874	0.588	0.632		0.893	0.955	0.901	0.621	0.852	0.826	0.615
Arsenic	0.821	1.000	0.796	0.840	0.810	0.953	0.848	0.664			0.852	0.919	0.942	0.887	0.741	0.711	0.807
Barium	0.680	0.796	1.000	0.613	0.525	0.784	0.746	0.845			0.695	0.719	0.751	0.869	0.790	0.846	0.669
Cadmium	0.934	0.840	0.613	1.000	0.582	0.830	0.874	0.522	0.654		0.854	0.944	0.929	0.610	0.780	0.803	0.604
Cobalt	0.484	0.810	0.525	0.582	1.000	0.868	0.527	0.549			0.672	0.624	0.703	0.736			0.599
Copper	0.802	0.953	0.784	0.830	0.868	1.000	0.808	0.742			0.876	0.880	0.934	0.841	0.676	0.756	0.725
Lead	0.874	0.848	0.746	0.874	0.527	0.808	1.000	0.742			0.736	0.930	0.940	0.648	0.874	0.776	0.720
Manganese	0.588	0.664	0.845	0.522	0.549	0.742	0.742	1.000			0.620	0.605	0.698	0.632	0.676	0.731	
Mercury	0.632			0.654					1.000		0.554	0.580	0.489				
Molybdenum										1.000							
Selenium	0.893	0.852	0.695	0.854	0.672	0.876	0.736	0.620	0.554		1.000	0.872	0.843	0.733	0.705	0.750	0.567
Silver	0.955	0.919	0.719	0.944	0.624	0.880	0.930	0.605	0.580		0.872	1.000	0.968	0.740	0.825	0.795	0.787
Thallium	0.901	0.942	0.751	0.929	0.703	0.934	0.940	0.698	0.489		0.843	0.968	1.000	0.764	0.775	0.795	0.747
Vanadium	0.621	0.887	0.869	0.610	0.736	0.841	0.648	0.632			0.733	0.740	0.764	1.000	0.571	0.700	0.786
Zinc	0.852	0.741	0.790	0.780		0.676	0.874	0.676			0.705	0.825	0.775	0.571	1.000	0.773	0.621
pH	0.826	0.711	0.846	0.803		0.756	0.776	0.731			0.750	0.795	0.795	0.700	0.773	1.000	0.536
Total solids	0.615	0.807	0.669	0.604	0.599	0.725	0.720				0.567	0.787	0.747	0.786	0.621	0.536	1.000

Note: Only significant ( $p < 0.10$ ) correlation estimates are reported.  
Spearman rank non-parametric correlation was used.

Table 6-8. Summary of correlations among vegetation variables at coastal plain and tundra communities

	Distance	Diversity	Evenness	Richness	Deciduous shrub cover	Evergreen shrub cover	Forb cover	Graminoid cover	Lichen cover	Lichen frequency	Moss cover	Moss frequency	Unvegetated cover
Distance	1.000		0.908	-0.892		0.741	-0.710		0.994	0.911	0.593		-0.803
Diversity		1.000						0.544					-0.593
Evenness	0.908		1.000	-0.769		0.553	-0.726			0.631			-0.700
Richness	-0.892		-0.769	1.000		-0.489	0.699						
Deciduous shrub cover					1.000								
Evergreen shrub cover	0.741		0.553	-0.489		1.000	-0.551	0.586	0.645	0.630			
Forb cover	-0.710		-0.726	0.699		-0.551	1.000		-0.502	-0.549			
Graminoid cover		0.544				0.586		1.000					
Lichen cover	0.994					0.645	-0.502		1.000	0.889	0.624		-0.615
Lichen frequency	0.911		0.631			0.630	-0.549		0.889	1.000	0.561		-0.697
Moss cover	0.593								0.624	0.561	1.000	0.513	-0.606
Moss frequency											0.513	1.000	
Unvegetated cover	-0.803	-0.593	-0.700						-0.615	-0.697	-0.606		1.000
Antimony	-0.809					-0.580			-0.895	-0.866			0.480
Arsenic	-0.950		-0.664			-0.764	0.618		-0.778	-0.790			0.489
Barium	-0.836		-0.699				0.617		-0.741	-0.836			0.611
Cadmium	-0.833					-0.759	0.502		-0.912	-0.832	-0.478		
Cobalt	-0.803		-0.604			-0.649	0.576	-0.484					
Copper	-0.914		-0.676	0.575		-0.724	0.714		-0.773	-0.768			0.493
Lead	-0.840		-0.588			-0.732	0.634		-0.834	-0.922	-0.500		0.524
Manganese	-0.698		-0.643	0.533			0.694		-0.575	-0.733			0.527
Mercury					-0.500				-0.503	-0.486			
Molybdenum													
Selenium	-0.678					-0.487	0.548		-0.753	-0.717			
Silver	-0.895		-0.534			-0.748	0.541		-0.877	-0.892	-0.531		0.512
Thallium	-0.914		-0.577			-0.790	0.634		-0.873	-0.861	-0.500		0.493
Vanadium	-0.945	-0.571	-0.681			-0.547	0.551		-0.657	-0.692			0.527
Zinc	-0.796		-0.549			-0.558	0.528		-0.807	-0.910			0.540
pH	-0.878					-0.475	0.497		-0.912	-0.905	-0.491		0.605
Total Solids	-0.945	-0.544	-0.808	0.525		-0.707			-0.602	-0.750	-0.555		0.662

**Note:** Only significant ( $p < 0.10$ ) correlation estimates are reported.  
Spearman rank non-parametric correlation was used.

**Table 6-9. Summary of correlations between PCA factors and distance or soil characteristics**

	Factor 1		Factor 2		Factor 3	
	Correlation	<i>P</i> -value	Correlation	<i>P</i> -value	Correlation	<i>P</i> -value
Distance	0.286	0.2875	-0.480	0.0712	-0.182	0.4927
<b>Soil Parameters</b>						
Antimony	-0.101	0.6484	<b>0.486</b>	<b>0.0301</b>	<b>0.453</b>	<b>0.0430</b>
Arsenic	<b>-0.565</b>	<b>0.0115</b>	<b>0.578</b>	<b>0.0099</b>	-0.198	0.3739
Barium	0.108	0.6297	<b>0.645</b>	<b>0.0040</b>	0.120	0.5950
Cadmium	-0.223	0.3164	<b>0.569</b>	<b>0.0111</b>	<b>0.394</b>	<b>0.0789</b>
Cobalt	-0.327	0.1425	<b>0.430</b>	<b>0.0549</b>	0.104	0.6443
Copper	-0.271	0.2237	<b>0.801</b>	<b>0.0003</b>	-0.026	0.9052
Lead	-0.332	0.1363	<b>0.636</b>	<b>0.0045</b>	0.274	0.2215
Manganese	0.245	0.2736	<b>0.482</b>	<b>0.0314</b>	0.138	0.5400
Mercury	-0.067	0.7624	0.283	0.2061	<b>0.447</b>	<b>0.0461</b>
Molybdenum	-0.209	0.3494	0.244	0.2758	-0.361	0.1055
Selenium	<b>-0.405</b>	<b>0.0695</b>	<b>0.495</b>	<b>0.0270</b>	0.087	0.7005
Silver	-0.174	0.4346	<b>0.695</b>	<b>0.0019</b>	0.301	0.1796
Thallium	-0.030	0.8914	<b>0.773</b>	<b>0.0006</b>	0.084	0.7079
Vanadium	<b>-0.456</b>	<b>0.0412</b>	<b>0.534</b>	<b>0.0171</b>	-0.256	0.2514
Zinc	-0.284	0.2024	<b>0.532</b>	<b>0.0174</b>	0.321	0.1522
pH	-0.271	0.2251	<b>0.508</b>	<b>0.0233</b>	0.269	0.2307
Total solids	-0.021	0.9213	<b>0.400</b>	<b>0.0740</b>	-0.042	0.8480

**Note:** Bold values indicate significant correlation at  $p < 0.10$ .

Spearman rank non-parametric correlation was used to estimate correlations.

PCA - principal component analysis



Table 6-10. Average percent cover and frequency results at coastal plain<sup>a</sup> stations

Species	Common Name	Site								Reference	
		TT50010		TT50100		TT51000		TT52000		TS-REF-12	
		C	F	C	F	C	F	C	F	C	F
<b>Forbs</b>											
<i>Anemone narcissiflora</i>	Anemone	0.25	10	--	--	--	--	--	--	--	--
<i>Androsace</i> sp.	Primrose	--	10	--	--	--	--	--	--	--	--
<i>Pedicularis capitata</i>	Lousewort	--	10	--	--	--	--	--	--	--	--
<i>Petasites frigidus</i> or <i>hyperboreus</i>	Sweet coltsfoot	4.75	100	7.25	100	--	--	--	--	--	--
<i>Polemonium acutiflorum</i>	Jacob's ladder	0.25	50	1.25	90	--	--	--	--	--	--
<i>Polygonum viviparum</i>	Alpine meadow bistort	--	20	--	--	--	--	--	--	--	--
<i>Saussurea angustifolia</i>	Saussurea	--	10	--	--	--	--	--	--	--	--
<i>Stellaria laeta</i>	Chickweed	--	30	0.75	60	--	--	--	--	--	--
<i>Valeriana capitata</i>	Valerian	--	20	1.75	20	--	--	--	--	--	--
<b>Graminoids</b>											
<i>Arctagrostis latifolia</i> var. <i>arundinaceae</i>	Polar grass	0.25	20	0.5	60	--	--	--	--	--	--
<i>Carex aquatilis</i>	Carex	0.25	10	1.75	30	--	--	1.25	70	1	50
<i>Caryx bigelowii</i>	Bigelow's sedge	0.25	10	0.25	20	1	40	--	20	0.75	30
<i>Calamagrostis holmii</i>	Bluejoint grass	--	--	--	--	--	--	--	--	0.25	10
<i>Calamagrostis</i> sp.	Bluejoint grass	--	--	0.25	10	--	10	--	--	--	--
<i>Eriophorum angustifolium subarcticum</i>	Cottongrass	3.25	40	5.25	60	3.5	40	0.25	10	2.5	60
<i>Eriophorum vaginatum</i>	Cottongrass	8.25	80	8	90	13.5	100	20.5	100	18.3	100
<i>Hierchloe alpina</i>	Holy grass	--	10	--	10	--	--	--	--	--	--
<i>Luzula multiflora multiflora</i>	Wood rush	0.25	10	--	--	--	--	--	--	--	--
<i>Luzula wahlenbergii</i>	Wood rush	--	--	--	--	--	--	--	--	1	40
<i>Poa lanata</i>	Bluegrass	5.25	70	3.75	100	--	--	--	--	--	10
<b>Deciduous Shrubs</b>											
<i>Betula nana exilis</i>	Dwarf birch	9.25	40	23	60	14.3	90	12.3	60	3	20
<i>Rubus chamaemorus</i>	Salmonberry	0.75	60	7.5	100	1.5	80	6	100	13.5	100
<i>Salix arctica</i>	Arctic willow	--	10	--	--	--	--	--	--	--	--
<i>Salix planifolia pulchra</i>	Diamondleaf willow	21.5	70	0.25	10	--	10	--	--	--	--
<i>Salix polaris</i>	Polar willow	0.25	10	--	--	--	--	--	--	--	--
<i>Vaccinium uliginosum alpinum</i>	Alpine blueberry	--	--	--	--	8.25	90	14.75	100	3.75	50
<b>Evergreen Shrubs</b>											
<i>Empitrum nigrum hermaphroditum</i>	Crowberry	--	10	--	--	2.5	60	4.75	50	1.5	10
<i>Ledum palustre decumbens</i>	Labrador tea	1	40	--	--	12.3	100	14.8	100	21.8	100
<i>Vaccinium vitis-idaea minus</i>	Lingonberry	0.25	20	--	--	13.3	100	12.3	100	13.8	100

Table 6-10. (cont.)

Species	Common Name	Site								Reference	
		TT50010		TT50100		TT51000		TT52000		TS-REF-12	
		C	F	C	F	C	F	C	F	C	F
<b>Other</b>											
Broadleaf litter	Broadleaf litter	18.3	90	13.5	100	2.25	100	10	100	17	100
Dry blades	Dry blades	37.3	100	46.5	100	45.3	100	38.3	100	38	100
Lichen	Lichen	--	--	0.25	40	2.75	100	8.25	90	15.8	100
Moss	Moss	4.25	90	62	100	34.5	100	39.8	100	45	100
Water	Water	0.5	20	--	--	--	--	--	--	--	--
<b>Unvegetated</b>											
Bare ground	Bare ground	2.25	90	--	--	--	10	--	--	0.5	20
Road gravel	Road gravel	4	70	--	--	--	--	--	--	--	--

**Note:** -- - not identified in any 1-m<sup>2</sup> microplot  
 C - average 1-m<sup>2</sup> microplot cover percentage  
 F - percent frequency in ten 1-m<sup>2</sup> microplots

<sup>a</sup> Coastal plain mesic tussock tundra community.

Table 6-11. Average percent cover and frequency results at tundra<sup>a</sup> stations

Species	Common Name	Site												Reference			
		TT30010		TT80010		TT30100		TT80100		TT31000		TT81000		TS-REF-5		TS-REF-7	
		C	F	C	F	C	F	C	F	C	F	C	F	C	F	C	F
<b>Forbs</b>																	
<i>Arnica lessingii lessingii</i>	Arnica	0.25	10	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<i>Equisetum arvense</i>	Horsetail	--	10	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<i>Petasites frigidus or hyperboreus</i>	Sweet coltsfoot	--	--	6.5	100	--	--	--	10	--	--	--	--	--	--	1.5	20
<i>Pedicularis labradorica</i>	Lousewort	--	--	0.25	10	--	--	0.25	10	0.25	10	--	--	0.25	10	0.5	30
<i>Stellaria laeta</i>	Chickweed	--	--	--	10	--	--	--	--	--	--	--	--	--	--	--	--
<b>Graminoids</b>																	
<i>Arctagrostis latifolia var. latifolia</i>	Polar grass	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.25	20
<i>Carex aquatilis</i>	Carex	2.5	50	--	--	2.5	60	0.25	10	--	--	2.25	40	--	--	--	--
<i>Caryx bigelowii</i>	Bigelow's sedge	--	--	15.5	90	--	--	14.3	90	--	50	3.75	50	--	10	1.75	70
<i>Carex rotundata</i>	Sedge	--	--	--	--	0.25	10	--	--	--	--	--	--	--	--	--	--
<i>Eriophorum angustifolium subarcticum</i>	Cottongrass	0.25	10	--	--	2	30	--	--	0.25	10	0.25	10	0.25	10	0.25	10
<i>Eriophorum vaginatum</i>	Cottongrass	15.8	100	5.25	70	20.5	100	12.8	80	14.8	100	24.3	100	12.3	100	24.5	100
<i>Luzula multiflora multiflora</i>	Wood rush	--	--	0.25	10	--	--	--	--	--	--	--	--	--	--	--	--
<b>Deciduous Shrubs</b>																	
<i>Betula nana exilis</i>	Dwarf birch	14.5	100	35.5	100	16.8	100	31	100	11	100	8.75	100	5.25	70	16.8	100
<i>Rubus chamaemorus</i>	Salmonberry	22.8	100	1	50	11.8	80	3.75	50	4.75	100	2.75	80	28.5	100	15.3	100
<i>Salix ovalifolia</i>	Ovaleaf willow	--	--	--	--	3.75	10	--	--	--	--	--	--	--	--	--	--
<i>Salix planifolia pulchra</i>	Diamondleaf willow	--	--	8	30	--	--	0.25	10	--	--	--	--	0.25	10	1.75	20
<i>Vaccinium uliginosum alpinum</i>	Alpine blueberry	28.8	100	1	40	26.3	100	3	20	28.8	100	20	90	37.8	100	26.5	70
<b>Evergreen Shrubs</b>																	
<i>Andromeda polifolia</i>	Bog rosemary	0.75	70	--	--	2	70	--	--	--	--	--	--	0.5	50	0.25	30
<i>Empitrum nigrum hermaphroditum</i>	Crowberry	4.25	90	0.75	50	2.5	80	5.25	20	3.75	70	5	50	8.5	100	8.5	90
<i>Ledum palustre decumbens</i>	Labrador tea	1.75	100	8.75	100	13.5	100	24	100	11.3	100	15.8	100	15.5	100	16	100
<i>Vaccinium vitis-idaea minus</i>	Lingonberry	0.75	90	1.75	100	2	40	10	100	15.8	100	18	100	4.25	100	7	100
<b>Other</b>																	
Broadleaf litter	Broadleaf litter	13.3	100	8.5	100	21	80	10	100	14.5	90	3.5	100	45.3	100	12.3	100
Dry blades	Dry blades	21.8	100	55	100	32.3	100	40	100	35.8	100	40.3	100	17	100	31.3	100
Lichen	Lichen	--	--	--	--	2.25	60	0.5	50	4.75	100	5	100	21.8	100	9.75	100
Moss	Moss	26.3	100	14.3	100	34.3	90	40.5	100	37.5	100	48.8	100	45.5	100	52.3	100
Water	Water	1.5	10	0.25	10	18	50	3.5	40	--	--	4.75	40	0.5	30	0.75	40
<b>Unvegetated</b>																	
Bare ground	Bare ground	4	20	2	30	--	--	0.5	20	--	--	--	--	--	--	--	--
Road gravel	Road gravel	2.25	70	3.75	70	--	--	--	--	--	--	--	--	--	--	--	--
Rock	Rock	0.5	20	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Note: -- - not identified in any 1-m<sup>2</sup> microplot  
 C - average 1-m<sup>2</sup> microplot cover percentage  
 F - percent frequency in ten 1-m<sup>2</sup> microplots

<sup>a</sup> Foothills mesic tussock tundra community.

Table 6-12. Average percent cover and frequency results at hillslope<sup>a</sup> stations

Species	Common Name	Site						Reference	
		TT60010		TT60100		TT61000		TS-REF-11	
		C	F	C	F	C	F	C	F
<b>Forbs</b>									
<i>Anemone</i> sp.	Anemone	--	20	--	--	--	20	--	--
<i>Artemisia arctica arctica</i>	Wormwood	--	--	--	--	0.25	10	0.25	30
<i>Arnica lessingii lessingii</i>	Arnica	--	--	--	--	--	--	0.25	20
<i>Equisetum arvense</i>	Horsetail	--	--	0.25	30	3.5	40	2.75	80
<i>Equisetum</i> sp.	Horsetail	--	--	--	10	--	--	--	--
Unknown forb	Unknown forb	--	10	--	--	--	--	--	10
Unknown forb	Unknown forb	--	--	--	10	--	--	--	--
Unknown forb	Unknown forb	0.25	10	--	30	--	--	--	--
Unknown forb	Unknown forb	--	--	--	--	0.25	20	--	--
Unknown forb	Unknown forb	--	--	--	--	--	40	--	--
Unknown forb, LEG Family	Unknown forb	--	--	--	--	0.25	20	--	--
<i>Lupinus arcticus</i>	Lupine	--	--	--	--	0.25	10	--	--
<i>Lycopodium clavatum monostachyon</i>	Club moss	--	--	--	--	--	--	0.25	10
<i>Minuartia arctica</i>	Sandwort	0.25	10	--	--	0.5	20	--	--
<i>Papaver macounni</i>	Alaska poppy	--	--	--	--	0.25	10	--	--
<i>Petasites frigidus or hyperboreus</i>	Sweet coltsfoot	1.75	80	0.75	80	--	10	2	50
<i>Pedicularis labradorica</i>	Lousewort	0.75	40	--	--	0.75	30	--	--
<i>Polemonium acutiflorum</i>	Jacob's ladder	--	--	--	20	--	--	--	--
<i>Polygonum bistorta plumosum</i>	Bistort	--	20	--	--	0.75	50	--	10
<i>Potentilla fruiticosa</i>	Shrubby cinquefoil	--	--	--	--	--	10	--	--
<i>Polygonum viviparum</i>	Alpine meadow bistort	--	--	--	--	0.75	30	--	--
<i>Pyrola grandiflora</i>	Wintergreen	0.5	20	--	10	0.25	20	--	--
<i>Ranunculus lapponicus</i>	Buttercup	--	--	--	--	--	--	--	10
<i>Ranunculus</i> sp.	Buttercup	--	--	--	--	--	20	--	--
<i>Saussurea angustifolia</i>	Saussurea	0.25	10	--	--	2	30	--	--
<i>Saxifraga punctata</i>	Brook saxifrage	--	10	--	--	0.25	40	--	--
<i>Senecio</i> sp.	Senecio	--	--	--	--	0.25	40	--	--
<i>Stellaria laeta</i>	Chickweed	--	20	--	--	--	20	--	10
<i>Valeriana capitata</i>	Valerian	--	--	0.25	20	--	--	--	10

Table 6-12. (cont.)

Species	Common Name	Site						Reference	
		TT60010		TT60100		TT61000		TS-REF-11	
		C	F	C	F	C	F	C	F
<b>Graminoids</b>									
<i>Arctagrostis latifolia</i> var. <i>latifolia</i>	Polar grass	0.25	40	0.25	20	--	10	0.75	60
<i>Caryx bigelowii</i>	Bigelow's sedge	--	--	10.8	100	7	70	3.5	80
<i>Carex microchaeta</i>	Sedge	1.25	50	--	--	--	--	0.25	30
<i>Carex podocarpa</i>	Sedge	0.25	10	--	--	--	--	--	--
<i>Carex saxatilis laxa</i>	Sedge	--	--	--	--	1.5	20	--	--
<i>Eriophorum vaginatum</i>	Cottongrass	--	--	0.25	10	--	--	--	--
<i>Festuca altaica</i>	Fescue grass	0.75	30	--	--	0.25	10	0.25	10
Unknown grass	Unknown grass	--	--	--	--	1.75	20	--	--
<i>Poa glauca</i>	Tundra bluegrass	--	--	--	--	--	20	--	--
<i>Poa</i> sp.	Bluegrass	0.5	50	0.25	30	--	--	--	10
Unknown sedge, CYP Family	Unknown sedge	--	--	--	--	0.25	10	--	--
<b>Deciduous Shrubs</b>									
<i>Betula nana exilis</i>	Dwarf birch	26.5	100	16.8	60	30.8	60	10.5	40
<i>Rubus chamaemorus</i>	Salmonberry	--	--	1.5	70	--	--	0.75	60
<i>Salix glauca</i>	Grayleaf willow	--	--	32.3	60	--	--	--	--
<i>Salix lanata richardsonii</i>	Richardson willow	1.75	20	3.5	40	10.8	40	--	--
<i>Salix planifolia pulchra</i>	Diamondleaf willow	6	60	--	--	1.5	10	23	70
<i>Salix reticulata</i>	Netleaf willow	--	--	1.5	10	15.8	60	--	--
<i>Salix</i> sp.	Dwarf willow	--	--	--	--	--	--	0.25	10
<i>Vaccinium uliginosum alpinum</i>	Alpine blueberry	30.8	90	7	40	9.25	40	40.5	100
<b>Evergreen Shrubs</b>									
<i>Andromeda polifolia</i>	Bog rosemary	--	--	--	10	--	--	--	--
<i>Arctostaphylos alpina</i> <sup>b</sup>	Bearberry	4	20	--	--	--	--	--	--
<i>Cassiope tetragona</i>	Heather	--	--	--	--	1.5	10	--	10
<i>Dryas integrifolia integrifolia</i>	Dryas	--	--	1.75	20	7	40	--	--
<i>Empitrum nigrum hermaphroditum</i>	Crowberry	4.75	40	0.25	20	--	20	10.3	100
<i>Ledum palustre decumbens</i>	Labrador tea	3.5	100	0.75	50	0.25	10	9.25	70
<i>Linnaea borealis borealis</i>	Twin flower	0.5	20	--	--	--	--	--	--
<i>Rhododendron lapponicum</i>	Lapland rosebay	--	--	--	--	1.5	10	--	--
<i>Vaccinium vitis-idaea minus</i>	Lingonberry	6	100	0.75	60	0.25	30	1.75	90

Table 6-12. (cont.)

Species	Common Name	Site						Reference	
		TT60010		TT60100		TT61000		TS-REF-11	
		C	F	C	F	C	F	C	F
<b>Other</b>									
Broadleaf litter	Broadleaf litter	13.5	100	8.75	100	19.5	100	17	100
Dry blades	Dry blades	4.5	100	11.75	100	10.5	100	11	100
Lichen	Lichen	10.8	90	10	100	14.3	90	18	80
Moss	Moss	79.3	100	71.5	100	70	100	50.3	100
Water	Water	--	--	--	--	--	--	0.25	10
<b>Unvegetated</b>									
Road gravel	Road gravel	0.25	10	--	--	--	--	--	--
Rock	Rock	0.25	20	--	--	0.25	10	0.25	10

**Note:** -- - not identified in any 1-m<sup>2</sup> microplot  
 C - average 1-m<sup>2</sup> microplot cover percentage  
 F - percent frequency in ten 1-m<sup>2</sup> microplots

<sup>a</sup> Hillslope mesic open shrubland community.

<sup>b</sup> Nondeciduous shrub per Walker (2000).

Table 6-13. Average percent cover and frequency results at coastal lagoon<sup>a</sup> stations

Species	Species Code	Common Name	Site				Reference			
			PLNL		NLK		CL-REF-1		CL-REF-2	
			C	F	C	F	C	F	C	F
<b>Forbs</b>										
<i>Hippuris vulgaris</i>	HIVU	Mare's tail	65.3	100	--	--	6.75	100	--	--
<i>Polygonum bistorta plumosum</i>	POBI	Bistort	--	--	--	--	--	--	--	--
<i>Potentilla egedii egedii</i>	POEG	Beach cinquefoil	--	--	--	--	--	--	0.25	30
<i>Ranunculus confervoides</i>	RACO	Buttercup	--	--	1.75	40	--	--	--	--
<i>Ranunculus hyperborealis hyperborealis</i>	RAHY	Buttercup	0.5	50	--	--	--	--	--	--
<i>Rumex arcticus</i>	RUAR	Sourdock	--	--	0.25	10	--	--	--	--
<i>Stellaria crassifolia</i>	STCR	Chickweed	0.25	40	1.75	90	0.25	30	--	20
Unknown forb, CRU Family	FONLK2	Unknown mustard	--	--	0.75	50	--	--	--	--
Unknown forb, RAN Family	FONLK3	Unknown buttercup	--	--	0.5	40	--	--	--	--
<b>Graminoids</b>										
<i>Arctophila fulva</i>	ARFU	Pendent grass	3.75	50	--	--	26.5	100	1.75	60
<i>Calamagrostis deschampsiioides</i>	CADE	Bluejoint grass	--	--	2.5	50	--	--	3.25	30
<i>Carex aquatilis</i>	CAAQ	Carex	--	--	37.8	100	--	--	49.8	100
<i>Carex canescens</i>	CACA2	Sedge	--	--	--	--	8	80	2	30
<i>Deschampsia caespitosa</i>	DECA	Tufted hairgrass	0.25	10	0.75	40	0.5	30	0.75	30
<i>Dupontia fischeri psilosantha</i>	DUFI	Tundra grass	10.5	50	37.8	100	19.8	80	22.8	60
<i>Eriophorum angustifolium subarcticum</i>	ERAN	Cottongrass	4.5	40	--	--	15	90	--	--
<i>Poa alpigena</i>	POAL	Bluegrass	--	--	--	20	--	--	--	--
<b>Deciduous Shrubs</b>										
<i>Salix ovalifolia</i>	SAOV	Ovaleaf willow	--	--	--	--	--	10	--	--
<b>Other</b>										
Broadleaf litter	Broadleaf litter	Broadleaf litter	--	--	--	--	17	100	--	--
Detritus/fines	Detritus/fines	Detritus/fines	4.25	70	2.5	100	7.5	100	12.3	100
Dry blades	Dry blades	Dry blades	16.5	50	66.5	100	11	100	15.5	100
Lichen	Lichen	Lichen	--	--	--	--	18	80	--	--
Littoral matter	Littoral matter	Littoral matter	--	--	2.25	50	--	--	0.25	10
Moss	Moss	Moss	3.25	40	38	100	50.3	100	34.5	100
Water	Water	Water	68.3	80	--	--	0.25	10	--	--
<b>Unvegetated</b>										
Rock	Rock	Rock	--	--	--	--	0.25	10	1.5	10
Sand/gravel	Sand/gravel	Sand/gravel	--	--	1.5	10	--	--	3.75	10

Note: -- - not identified in any 1-m<sup>2</sup> microplot

C - average 1-m<sup>2</sup> microplot cover percentage

F - percent frequency in ten 1-m<sup>2</sup> microplots

<sup>a</sup> Coastal lagoon fringe emergent community.

**Table 6-14. Vascular plant species diversity, evenness, and richness at terrestrial and coastal lagoon plant community survey stations**

Station	Diversity	Evenness	Richness	Area Richness
<b>Coastal Lagoon Fringe Emergent</b>				
<b>Site</b>				
PLNL	1.2	0.42	7	12
NLK	1.6	0.51	10	11
<b>Reference</b>				
CL-REF-1	2.2	0.79	8	8
CL-REF-2	1.5	0.52	8	11
<b>Coastal Plain Mesic Tussock Tundra</b>				
<b>Site</b>				
TT50010	2.7	0.67	25	33
TT50100	2.8	0.74	15	22
TT51000	2.8	0.88	11	13
TT52000	2.8	0.87	10	10
<b>Reference</b>				
TS-REF-12	2.7	0.77	13	13
<b>Foothills Mesic Tussock Tundra</b>				
<b>Site</b>				
TT30010	2.5	0.72	12	16
TT80010	2.5	0.71	13	19
TT30100	2.9	0.81	12	15
TT80100	2.7	0.78	12	14
TT31000	2.6	0.82	10	10
TT81000	2.8	0.84	10	11
<b>Reference</b>				
TS-REF-5	2.5	0.73	12	13
TS-REF-7	3.0	0.78	14	14
<b>Hillslope Mesic Open Shrubland</b>				
<b>Site</b>				
TT60010	2.8	0.66	25	29
TT60100	2.6	0.63	23	29
TT61000	3.4	0.69	38	53
<b>Reference</b>				
TS-REF-11	2.7	0.66	24	35

**Note:** Shannon-Weiner diversity index and evenness index calculated using average percent covers for vascular plant species.

Species richness represents the number of vascular plant species observed in ten 1-m<sup>2</sup> microplots.

Area richness represents the number of vascular plant species observed in the near vicinity of the survey line.



Table 6-15. CoPC concentrations, pH, and total solids in tundra soil at terrestrial and coastal lagoon plant community survey stations

Station	pH (wet)	Total Solids (% wet)	Antimony	Arsenic	Barium	Cadmium	Cobalt	Copper	Lead	Manganese	Mercury	Molybdenum	Selenium	Silver	Thallium	Vanadium	Zinc
<b>Coastal Lagoon<sup>a</sup></b>																	
<b>Site</b>																	
PLNL	6.5	13.5	2.8 <i>J</i>	13	429	17.7	8.84	37.1	532	160	0.424	1.96 <i>J</i>	2.2	1.87	0.239	30.4	3,010
NLK	5.6	31	0.31 <i>J</i>	16.6	264	2.47	7.54	16.1	156	249	0.141	2.58 <i>J</i>	1.3	0.151	0.109	31.3	446
<b>Reference</b>																	
CL-REF-1	5.4	15.3	0.17 <i>J</i>	8.57	123	0.204	3.7	9.55	11.5	78.4	0.069	1.13 <i>J</i>	1.1	0.069	0.021	25.7	42.4
CL-REF-2	4.7	34.2	0.2 <i>J</i>	11.4	206	0.524	9.7	15.4	36	331	0.06	3.08 <i>J</i>	0.9	0.129	0.093	32.8	158
<b>Coastal Plain<sup>b</sup></b>																	
<b>Site</b>																	
TT5_0010	6.7	42.9	2.75	8	1,200	20.6	18.6	20.6	1,210	1,860 <i>J</i>	1.75	0.89	1.5	1.91	0.455	11.6	4,330 <i>J</i>
TT5_0100	6.4	34.7	2.46	5.3	1,200	24	8.18	19.8	1,060	1,560 <i>J</i>	0.25	0.84	1.9	1.25	0.368	8.25	5,120 <i>J</i>
TT5_1000	4.5	19	0.83	1.8	15.3	4.08	6.82	7.93	8.62	32.5 <i>J</i>	0.33	1.16	0.9	0.264	0.072	4.64	38.9 <i>J</i>
TT5_2000	4.5	18.9	0.55	0.5	96	1.31	1.97	4.82	54.1	290 <i>J</i>	0.27	0.8	0.5	0.12	0.036	0.98	286 <i>J</i>
<b>Reference</b>																	
TS-REF12	3.6	22	0.21 <i>J</i>	0.28	24.5	0.215	0.49	3.09	5.83	137	0.176	0.506 <i>J</i>	0.2	0.03	0.012	0.85	58.6
<b>Tundra<sup>c</sup></b>																	
<b>Site</b>																	
TT3-0010	7.1	42.8	0.93 <i>J</i>	5.3 <i>J</i>	2,280	7.07 <i>J</i>	8.3 <i>J</i>	19.0 <i>J</i>	385	2,110 <i>J</i>	0.3 <i>J</i>	1.08	1.0 <i>J</i>	0.45	0.30 <i>J</i>	14.2 <i>J</i>	1,350 <i>J</i>
TT8_0010	7.1	66.5	1.07	4.1	1,900	4.72	6.5	15.3	226	494 <i>J</i>	0.13	0.93	1.4	0.453	0.245	15.9	976 <i>J</i>
TT3-0100	6.3	25	0.91 <i>J</i>	2.2 <i>J</i>	690 <i>J</i>	2.1 <i>J</i>	2.69 <i>J</i>	7.5 <i>J</i>	119	348 <i>J</i>	0.12 <i>J</i>	0.48 <i>J</i>	0.5 <i>J</i>	0.32	0.088 <i>J</i>	5.0 <i>J</i>	465
TT8_0100	6.9	19.2	1.25	3	1,470	3.83	5.48	14.9	189	2,380 <i>J</i>	0.2	1.16	1.8	0.375	0.214	11.2	908 <i>J</i>
TT3-1000	4	18.8	0.4 <i>J</i>	0.8 <i>J</i>	131 <i>J</i>	0.55 <i>J</i>	0.6 <i>J</i>	5.87 <i>J</i>	16.1 <i>J</i>	250 <i>J</i>	0.1	0.79 <i>J</i>	0.4 <i>J</i>	0.07	0.049 <i>J</i>	1.5 <i>J</i>	78.4
TT8_1000	4.5	15.3	0.42	0.8	275	0.408	3.57	6.15	4.23 <i>U</i>	139 <i>J</i>	0.15	1.68	0.6	0.062	0.02	4.8	89.3 <i>J</i>
<b>Reference</b>																	
TS-REF-5	3.9	38	0.22 <i>J</i>	3.5 <i>J</i>	383 <i>J</i>	0.29 <i>J</i>	15.3 <i>J</i>	9.5 <i>J</i>	13.4	365 <i>J</i>	0.11 <i>J</i>	0.80 <i>J</i>	0.6	0.12	0.057 <i>J</i>	12.7 <i>J</i>	57.4
TS-REF-7	4.5	14.4	0.18 <i>J</i>	0.4 <i>J</i>	195 <i>J</i>	0.27 <i>J</i>	6.02 <i>J</i>	6.2 <i>J</i>	6.9 <i>J</i>	541 <i>J</i>	0.09 <i>J</i>	2.01 <i>J</i>	0.3 <i>J</i>	0.02	0.027 <i>J</i>	1.1 <i>J</i>	50.3
<b>Hillslope<sup>d</sup></b>																	
<b>Site</b>																	
TT6_0010	6.9	54.1	1.92	9.1	6950 <i>J</i>	5.47	9.11 <i>J</i>	45.6	349	1,020 <i>J</i>	0.25	1.95 <i>J</i>	1.5	0.726	1.29	19.7 <i>J</i>	1,020
TT6_0100	5.6	26.3	2.03	4.9	6360 <i>J</i>	5.06	3.3 <i>J</i>	20.1	281	534 <i>J</i>	0.27	2.47 <i>J</i>	0.9	0.486	0.755	7.51 <i>J</i>	764
TT6_1000	6.6	20.8	1.22	2.9	1290 <i>J</i>	6.11	1.87 <i>J</i>	15.8	145	429 <i>J</i>	0.22	2.09 <i>J</i>	1.6	0.697	0.38	16.0 <i>J</i>	592
<b>Reference</b>																	
TS-REF11	5.3	21	0.3	3.2	293 <i>J</i>	0.414	7.81 <i>J</i>	12.2	12.7	3,490 <i>J</i>	0.12	0.348 <i>J</i>	0.5	0.204	0.105	11.0 <i>J</i>	56.4

**Note:** Metals reported as mg/kg, dry weight basis.

Field replicates averaged.

Results averaged across sampling events (2003 and 2004).

CoPC - chemical of potential concern

*J* - estimated value

*U* - undetected; value reported is one-half the detection limit

<sup>a</sup> Coastal lagoon fringe emergent community.

<sup>b</sup> Coastal plain mesic tussock tundra community.

<sup>c</sup> Foothills tussock tundra community.

<sup>d</sup> Hillslope mesic open shrubland community.

Table 6-16. Comparison of CoPC concentrations in coastal lagoon sedge and grass samples against phytotoxicity thresholds reported in the literature

Station	Sample ID	Species	Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Lead	Mercury	Molybdenum	Selenium	Thallium	Vanadium	Zinc
<b>Phytotoxicity Reference Value</b>																
a			50–200	150	5–20	500	5–30	5–30	15–50	30–300	1–3	10–50	5–30	20	5–10	100–400
b			--	--	3–10	--	5–700	20	25–100	--	--	100	100	--	10	500–1,500
c			--	--	11–26	400–800	14–16	5–20	3–9	20–35	2–5	130–140	7–90	11–45	--	160–320
<b>Site</b>																
NLK	SE0049-D	<i>Carex aquatilis</i>	26.2	0.02 U	0.38	22.0	0.114	0.4 J	0.47	3.3	0.031	0.143	0.3	0.008	0.2 U	44.8
PLNL	SE0045	<i>Carex aquatilis</i>	17.4	0.05	0.13	19.5	0.078	0.1 U	0.15	1.81	0.026	0.231	0.2	0.004	0.2 U	40.5
PLNL	SE0046	<i>Eriophorum angustifolium</i>	2.0 U	0.027 U	0.03 U	14.5	0.034	0.4	0.03	0.76	0.045	0.077	0.05 U	0.004	0.2 U	49.7
<b>Reference</b>																
CL-REF-1	SE0057	<i>Eriophorum angustifolium</i>	8.0	0.025 U	0.03 U	16.4	0.034	0.3	0.1	0.82	0.052	0.104	0.05 U	0.004	0.2 U	35.2
CL-REF-1	SE0058	<i>Carex aquatilis</i>	13.2	0.020 U	0.03 U	18.7	0.072	0.4	0.31	0.69	0.055	0.072	0.05 U	0.0010 U	0.2 U	35.6
CL-REF-2	SE0060	<i>Carex aquatilis</i>	23.5	0.03 U	0.2	32.4	0.197	0.4	0.44	3.28	0.029	0.45	0.2	0.007	0.2 U	49.3

**Note:** Phytotoxicity reference values and tissue concentrations are in mg/kg dry weight.

Samples consisted primarily of above-ground tissue with some root material.

Plant shoots were unwashed, and roots were rinsed in site water.

Boxed concentrations exceed the minimum phytotoxicity reference value.

-- - not reported

CoPC- chemical of potential concern

U - undetected; value reported is one-half the detection limit

J - estimated value

<sup>a</sup> Leaf tissue concentration at which toxicity was observed in plants that are neither sensitive nor tolerant (McBride 1994).

<sup>b</sup> Phytotoxic concentrations in plant foliage (Langmuir et al. 2004).

<sup>c</sup> Upper critical level in leaves and shoots of spring barley (*Hordeum vulgare*) associated with reduced yield (Davis et al. 1978).

Table 6-17. Comparison of CoPC concentrations in unwashed willow and birch leaves against phytotoxicity thresholds reported in the literature

Station	Sample ID	Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Lead	Mercury	Molybdenum	Selenium	Thallium	Vanadium	Zinc
<b>Phytotoxicity Reference Value</b>															
a		50–200	150	5–20	500	5–30	5–30	15–50	30–300	1–3	10–50	5–30	20	5–10	100–400
b		--	--	3–10	--	5–700	20	25–100	--	--	100	100	--	10	500–1,500
c		--	--	11–26	400–800	14–16	5–20	3–9	20–35	2–5	130–140	7–90	11–45	--	160–320
<b>Site</b>															
TT2-0010	WI0006	93.6	0.15	0.13	77.2	16.5	0.2 U	0.456	5.76	0.05	0.174	0.2	0.013	0.2 U	548
TT2-0100	WI0005	10.8	0.14	0.04 U	14.1	1.72	0.2 U	0.738	0.89	0.04	0.149	0.1 U	0.002 U	0.2 U	234
TT2-1000	WI0004	7.5	0.03 U	0.04 U	9.1 J	0.26	0.2 U	0.41	0.35	0.04	0.078	0.1 U	0.002	0.3 U	244
TT3-0010	WI0007	154	0.18	0.19	154	4.43	0.55 U	0.426	7.74	0.06	0.342	0.1 U	0.025	0.2 U	345
TT3-0100	BR0004	28.4 J	0.03 UJ	0.04 U	123	0.136	0.2 U	0.306	1.91	0.03	0.369	0.1 U	0.002 U	0.2 U	138
TT3-0100	WI0011	9.1 J	0.02 UJ	0.04 U	54.4	4.71	0.2 U	3.11	0.48	0.03	0.48	0.1 U	0.0010 U	0.3 U	246
TT3-1000	BR0003	6.6 J	0.02 UJ	0.04 U	77.9	0.155	0.2 U	0.313	0.58	0.04	0.08	0.1 U	0.0010 U	0.3 U	275
TT5-0010	WI0001	66.7	0.17	0.1	57.5 J	10.3	0.2 U	0.604	6.64	0.05	0.122	0.1 U	0.006	0.2 U	592
TT5-0100	WI0003	43.0	0.04 U	0.04 U	35.3	2.14	0.4 U	0.965	4.85	0.04	0.075	0.1 U	0.005	0.2 U	267
TT5-1000	BR0001	8.4	0.12	0.04 U	42.3 J	0.728	0.4 U	0.437	3.77	0.04	0.084	0.1 U	0.0010 U	0.3 U	284
TT5-1000	WI0002	11.3	0.04 U	0.04 U	17.1 J	0.817	0.2 U	2.91	1.07	0.03	0.062	0.1 U	0.0010 U	0.3 U	163
TT5-2000	BR0002	5.7	0.03 U	0.04 U	28.0	0.323	0.2 U	0.76	0.42	0.03	0.222	0.1 U	0.0010 U	0.2 U	225
TT6-0010	WI0024	21.9	0.055 U	0.03 U	40.2	2.69	0.3 UJ	1.17	1.12	0.057	0.16	0.05 U	0.004	0.3 U	127 J
TT6-0100	WI0013-D	10.8 J	0.03 U	0.03 U	31.2 J	1.78	0.2 UJ	0.22	1.07	0.047	0.316	0.05 U	0.005	0.2 U	223 J
TT6-1000	WI0012	6.5 J	0.03 U	0.03 U	14.8 J	4.65	0.2 UJ	0.2	0.41	0.047	0.373	0.05 U	0.003	0.3 U	79.8 J
TT6-2000	WI0017	10.5 J	0.085 U	0.03 U	40.9 J	1.91	0.2 UJ	0.76	0.37	0.047	0.19	0.05 U	0.0010 U	0.2 U	103 J
TT7-0010	WI0016	41.5 J	0.11 U	0.03 U	144 J	2.66	0.2 UJ	1.86	6.89	0.048	0.089	0.05 U	0.006	0.2 U	145 J
TT7-1000	WI0015	2.5 UJ	0.03 U	0.03 U	17.1 J	4.88	0.2 UJ	0.13	0.52	0.034	0.883	0.05 U	0.011	0.2 U	227 J
TT7-2000	WI0029	6.5	0.026 U	0.03 U	54.0	4.48	0.4	0.25	1.4	0.042	0.284	0.05 U	0.007	0.2 U	330
TT7-2000	WI0030	5.5	0.01 U	0.03 U	15.5	3.68	0.4	0.22	1.09	0.056	0.992	0.05 U	0.004	0.2 U	225
TT8-0010	WI0010	108	0.12	0.11	57.2	6.73	0.5 U	1.6	2.91	0.05	0.312	0.2	0.007	0.5	239
TT8-0100	WI0009	20.4	0.05 U	0.04 U	54.6	1.77	0.2 U	3.23	0.79	0.04	0.316	0.1 U	0.0010 U	0.2 U	143
TT8-1000	WI0008	18.0	0.51	0.04 U	39.2	0.714	0.3 U	4.96	1.47	0.04	0.207	0.1 U	0.0010 U	0.3 U	126
<b>Reference</b>															
TS-REF-5	WI0019	14.8	0.03 U	0.03 U	76.1	0.673	0.2 U	8.03	0.4	0.044	0.166	0.1	0.0010 U	0.3 U	78.8
TS-REF-7	BR0005	2.5 U	0.02 U	0.03 U	69.9	0.16	0.4 U	0.54	0.13	0.047	0.15	0.05 U	0.003	0.2 U	107
TS-REF-7	WI0020	9.0	0.03 U	0.03 U	26.7	0.389	0.3 U	2.47	0.11	0.042	0.312	0.05 U	0.0010 U	0.2 U	62.1
TS-REF11	BR0006	2.5 U	0.02 U	0.03 U	74.4	0.108	0.4 UJ	0.1	0.08	0.049	0.005 U	0.05 U	0.0010 U	0.3 U	145 J
TS-REF11	WI0023	22.6	0.04 U	0.03 U	51.1	0.376	0.4 UJ	0.84	0.77	0.055	0.013 U	0.05 U	0.003	0.3 U	47.2 J

**Note:** Phytotoxicity reference values and tissue concentrations are in mg/kg dry weight.  
 "WI" samples are willow leaves (*Salix* spp.); "BR" samples are dwarf birch leaves (*Betula nana*).  
 Boxed concentrations exceed the minimum phytotoxicity reference value.  
 Plant tissue data were not available for the following CoPCs in tundra soil: copper, manganese, and silver.  
 -- - not reported  
 CoPC - chemical of potential concern  
 U - undetected; value reported is one-half the detection limit  
 J - estimated value

<sup>a</sup> Leaf tissue concentration at which toxicity was observed in plants that are neither sensitive nor tolerant (McBride 1994).  
<sup>b</sup> Phytotoxic concentrations in plant foliage (Langmuir et al. 2004).  
<sup>c</sup> Upper critical level in leaves and shoots of spring barley (*Hordeum vulgare*) associated with reduced yield (Davis et al. 1978).

Table 6-18. Comparison of CoPC concentrations in unwashed sedge blades against phytotoxicity thresholds reported in the literature

Station	Sample ID	Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Lead	Mercury	Molybdenum	Selenium	Thallium	Vanadium	Zinc
<b>Phytotoxicity Reference Value</b>															
a		50–200	150	5–20	500	5–30	5–30	15–50	30–300	1–3	10–50	5–30	20	5–10	100–400
b		--	--	3–10	--	5–700	20	25–100	--	--	100	100	--	10	500–1,500
c		--	--	11–26	400–800	14–16	5–20	3–9	20–35	2–5	130–140	7–90	11–45	--	160–320
<b>Site</b>															
TT2-0010	SE0010	69.2	0.19	0.08	85.0	0.256	0.70 U	0.133	5.63	0.05	0.322	0.1 U	0.007	0.2 U	89.4
TT2-0100	SE0006	17.4	0.21	0.04 U	30.7	0.05	0.4 U	0.028	1.01	0.03	0.345	0.1 U	0.0010 U	0.2 U	78.4
TT2-1000	SE0005	7	0.13	0.12	20	0.021	0.2 U	0.033	0.16	0.03	0.379	0.1 U	0.0010 U	0.2 U	52.5
TT3-0010	SE0013	72	0.13	0.09	84.3	0.138	3.3	0.162	4.06	0.03	0.177	0.1 U	0.032	0.2 U	40.4
TT3-0100	SE0022	17.9 J	0.02 UJ	0.04 U	44.5	0.052	0.2 U	0.048	0.91	0.04	0.551	0.1 U	0.0010 U	0.2 U	41.3
TT3-1000	SE0021	5.4 J	0.02 UJ	0.04 U	34.0	0.02	0.2 U	0.076	0.18	0.03	0.403	0.1 U	0.0010 U	0.3 U	51.3
TT5-0010	SE0001	93.8	0.22	0.13	85.8 J	0.567	0.65 U	0.159	10.8	0.05	0.239	0.1 U	0.01	0.2 U	209
TT5-0100	SE0004	26.1	0.17	0.04 U	39.7	0.104	0.3 U	0.046	2.33	0.03	0.172	0.1 U	0.0010 U	0.2 U	67.7
TT5-1000	SE0002-D	8.8	0.19	0.04 U	26.6	0.09	0.3	0.04	2.10	0.04	0.490	0.1 U	0.005	0.3 U	74.9
TT5-1000	SE0002-D	8.0	0.05 U	0.04 U	22.3	0.076	0.2 U	0.038	1.57	0.04	0.391	0.1 U	0.0010 U	0.2 U	64
TT5-2000	SE0003	2.5 U	0.04 U	0.04 U	18.6 J	0.036	0.2 U	0.016	0.33	0.04	0.618	0.1 U	0.009	0.3 U	72.8
TT6-0010	SE0042	13.1	0.03 U	0.03 U	128	0.121	0.2 UJ	0.09	0.71	0.04	0.463	0.3	0.004	0.3 U	58.9 J
TT6-0100	SE0024	13.8 J	0.03 U	0.03 U	47.4 J	0.053	0.2 UJ	0.005 U	0.33	0.017	1.15	0.05 U	0.002	0.3 U	47.4 J
TT6-0100	SE0025	8.6 J	0.03 U	0.03 U	87.7 J	0.122	0.2 UJ	0.04	1.32	0.026	1.72	0.1	0.007	0.2 U	53 J
TT6-1000	SE0023	2.8 UJ	0.03 U	0.03 U	33.6 J	0.038	0.3 UJ	0.03	0.3 U	0.025	0.454	0.4	0.002	0.3 U	33 J
TT6-2000	SE0028	8.6 J	0.070 U	0.03 U	66.7 J	0.398	0.2 UJ	0.09	1.10	0.03	0.134	0.05 U	0.007	0.2 U	71.4 J
TT7-0010	SE0027	12.5 J	0.05 U	0.08	77.0 J	0.403	0.2 UJ	0.11	2.24	0.022	0.556	0.05 U	0.035	0.3 U	166 J
TT7-1000	SE0026	12.2 J	0.04 U	0.03 U	78.4 J	0.172	0.3 UJ	0.04	5.67	0.036	0.888	0.05 U	0.006	0.3 U	43.4 J
TT7-2000	SE0061	7.2	0.02 U	0.03 U	62.3	0.197	0.3	0.05	1.95	0.045	1.26	0.2	0.009	0.2 U	49.7
TT7-2000	SE0062	14.9	0.031 U	0.04 U	93.6	0.378	0.2 UJ	0.07	7.96	0.035	0.669	0.05 U	0.007	0.2 U	65.2
TT8-0010	SE0017	104	0.16	0.11	109	0.164	4.5	0.209	4.89	0.04	0.816	0.3	0.007	0.2 U	51
TT8-0100	SE0015-D	23.5	0.04 U	0.04 U	70.8	0.071	0.4 U	0.061	1.3	0.04	0.767	0.1 U	0.001 U	0.2 U	35.7
TT8-1000	SE0014	6.9	0.04 U	0.04 U	32.3	0.034	0.2 U	0.019	0.34	0.01 U	0.44	0.1 U	0.01	0.2 U	48.4
<b>Reference</b>															
TS-REF-5	SE0031	8.3	0.02 U	0.03 U	34.7	0.071	0.2 U	0.04	0.52	0.023	0.422	0.05 U	0.004	0.3 U	35.9
TS-REF-5	SE0056	8.2	0.037 U	0.03 U	25.6	0.035	0.6	0.05	0.46	0.04	0.549	0.05 U	0.0010 U	0.2 U	30.6
TS-REF-7	SE0032	6.8	0.02 U	0.03 U	28.3	0.032	0.4 U	0.06	0.28	0.029	0.651	0.05 U	0.0010 U	0.3 U	37.9
TS-REF11	SE0041	6.8	0.02 U	0.05	16.7	0.025	0.3 UJ	0.08	0.39	0.036	0.236	0.3	0.0010 U	0.3 U	25.6 J

**Note:** Phytotoxicity reference values and tissue concentrations are in mg/kg dry weight.  
 Sedge samples include *Eriophorum vaginatum* and *Carex* species.  
 Boxed concentrations exceed the minimum phytotoxicity reference value.  
 Plant tissue data were not available for the following CoPCs in tundra soil: copper, manganese, and silver.  
 -- - not reported  
 CoPC - chemical of potential concern  
 U - undetected; value reported is one-half the detection limit  
 J - estimated value

<sup>a</sup> Leaf tissue concentration at which toxicity was observed in plants that are neither sensitive nor tolerant (McBride 1994).  
<sup>b</sup> Phytotoxic concentrations in plant foliage (Langmuir et al. 2004).  
<sup>c</sup> Upper critical level in leaves and shoots of spring barley (*Hordeum vulgare*) associated with reduced yield (Davis et al. 1978).

Table 6-19. Summary of sampling characteristics for site and reference stations

Station	Replicate	Water Temperature (°C)	Water Depth (cm)	Sampling Area (m <sup>2</sup> )	Mean Water Velocity (m/sec)	Sampling Period (min.)	Volume Sampled (m <sup>3</sup> )
<b>Site Stations</b>							
AC-R	A	10.5	28	0.13	0.56	60	266
AC-R	B	10.5	28	0.13	2.69	62	1,315
AC-R	C	10.4	27	0.13	2.26	69	1,201
AC-R	D	10.4	28	0.13	1.54	74	895
AC-R	E	10.4	28	0.13	1.71	76	1,023
ARC-R	A	8.9	25	0.12	1.46	55	560
ARC-R	B	--	27	0.13	2.53	60	1,140
ARC-R	C	9.0	28	0.13	1.92	65	985
ARC-R	D	8.9	28	0.13	1.71	67	900
ARC-R	E	--	28	0.13	0.66	70	364
OR-R	A	8.8	28	0.13	1.55	58	707
OR-R	B	8.8	28	0.13	1.50	60	709
OR-R	C	8.8	28	0.13	2.44	63	1,209
OR-R	D	8.8	28	0.13	1.64	68	876
OR-R	E	8.8	28	0.13	0.64	71	356
<b>Reference Stations</b>							
ST-REF-3	A	7.7	28	0.13	0.55	60	259
ST-REF-3	B	7.7	28	0.13	1.36	61	652
ST-REF-3	C	7.7	28	0.13	1.05	65	538
ST-REF-3	D	7.7	28	0.13	0.68	70	376
ST-REF-3	E	7.7	28	0.13	0.77	78	476
ST-REF-5	A	7.7	28	0.13	0.58	58	267
ST-REF-5	B	7.7	28	0.13	0.88	67	465
ST-REF-5	C	7.7	28	0.13	1.03	74	600
ST-REF-5	D	7.7	28	0.13	0.42	80	266
ST-REF-5	E	7.5	28	0.13	1.07	83	698
ST-REF-6	A	7.6	25	0.12	0.84	58	341
ST-REF-6	B	--	28	0.13	1.86	64	940
ST-REF-6	C	--	28	0.13	1.71	70	942
ST-REF-6	D	--	28	0.13	1.87	79	1,166
ST-REF-6	E	--	28	0.13	1.69	80	1,067

**Note:** -- - data not collected

Table 6-20. Summary of abundances of macroinvertebrates (per m<sup>3</sup>) in drift samples from site and reference stations

Phylum	Class	Order	Family	Genus	Site Stations			Reference Stations		
					ARC-R	OR-R	AC-R	ST-REF-3	ST-REF-5	ST-REF-6
Platyhelminthes	Turbellaria						0.0017	0.0027		
Nematoda	Nematoda					0.0028	0.0017		0.0014	
Annelida	Oligochaeta						0.054	0.081	0.012	0.13
Arthropoda	Arachnida	Acarina			0.33	0.66	0.18	0.12		0.34
	Crustacea	Ostracoda					0.020			
	Insecta	Coleoptera (Beetles)	Dytiscidae	<i>Agabinus</i>				0.00083		
				<i>Dytiscus</i>	0.0035	0.012	0.027			
				<i>Hydaticus</i>						0.0042
				<i>Oreodytes</i>		0.017				
			Hydrophilidae		0.0027			0.0039		
				<i>Ametor</i>			0.023			
				<i>Hydrophilus</i>						0.0037
		Collembola (Springtails)	Sminthuridae		0.0041		0.0039	0.0082		0.015
				<i>Sminthuris</i>	0.0044					
			Isotomidae		0.023	0.0090		0.041		0.19
			Onychiuridae		0.015		0.030	0.014		0.0069
			Poduridae			0.043	0.010	0.012		0.046
	Diptera (True flies)		Chironomidae		3.8	7.1	3.6	1.1	0.15	15
			Culicidae		0.034	0.012		0.010	0.0045	0.021
			Dolichopodidae				0.018	0.010	0.0017	0.0037
			Ephydriidae	<i>Scatella</i>		0.0046	0.0045			
			Empididae				0.0045			
				<i>Chelifera</i>	0.0035	0.017				
				<i>Oreogeton</i>	0.071					0.019
			Dixidae	<i>Dixella</i>					0.0038	
			Simuliidae	<i>Simulium</i>	1.0	1.7	1.1	0.63		1.7
			Tipulidae		0.0035			0.0074		0.012
				<i>Dicranota</i>	0.0071					0.0077
				<i>Tipula</i>	0.0035	0.0057	0.025	0.0074		0.020
	Ephemeroptera (Mayflies)		Ameletidae	<i>Ameletus</i>				0.0039		0.0037
			Baetidae		<i>Baetis</i>	0.75	0.97	16.5	3.3	0.0014
			Heptageniidae	<i>Cinygmula</i>	1.5		0.33	0.62		0.69
				<i>Epeorus</i>				0.0053		0.093

Table 6-20. (cont.)

Phylum	Class	Order	Family	Genus	Site Stations			Reference Stations		
					ARC-R	OR-R	AC-R	ST-REF-3	ST-REF-5	ST-REF-6
		Plecoptera (Stoneflies)	Capniidae	<i>Capnia</i>	2.6	0.23	5.3	0.33		0.31
			Chloroperlidae		0.0035	0.034		0.015		0.11
				<i>Neaviperla</i>	0.0027	0.071		0.11	0.0022	0.087
				<i>Utaperla</i>						0.0043
			Nemouridae		0.023	0.024	0.072	0.022		
				<i>Nemoura</i>	0.087	0.0028	0.11	0.052		0.0080
				<i>Zapada</i>			0.050	0.0021		0.080
		Trichoptera (Caddis flies)	Brachycentridae					0.0074		
				<i>Amiocentris</i>		0.0089	0.0025			0.0075
				<i>Brachycentrus</i>		0.0056	0.0039	0.0039		0.0043
			Limnephilidae				0.0039			
				<i>Grensia</i>	0.0085		0.00083	0.020		0.020
				<i>Hydatophylax</i>	0.023	0.37		0.16		1.8
<b>Grand Total</b>					<b>10</b>	<b>11</b>	<b>28</b>	<b>6.6</b>	<b>0.18</b>	<b>24</b>

**Table 6-21. Summary of benthic metrics at site and reference stations**

Station	Total Abundance (per m <sup>2</sup> )	Percent Dominance	Taxa Richness		Relative Abundance (percent)	
			Total	EPT	EPT	Chironomids
<b>Site Stations</b>						
ARC-R	10	37	24	9	49	37
OR-R	11	63	23	9	15	63
AC-R	28	60	26	10	82	13
<b>Reference Stations</b>						
ST-REF-3	6.6	50	27	14	71	17
ST-REF-6	24	61	31	15	29	62



Table 6-22. Comparison of CoPC concentrations in stream sedge and willow samples against phytotoxicity thresholds reported in the literature

Station	Sample ID	Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Lead	Mercury	Molybdenum	Selenium	Thallium	Vanadium	Zinc
<b>Phytotoxicity Reference Value</b>															
a		50–200	150	5–20	500	5–30	5–30	15–50	30–300	1–3	10–50	5–30	20	5–10	100–400
b		--	--	3–10	--	5–700	20	25–100	--	--	100	100	--	10	500–1,500
c		--	--	11–26	400–800	14–16	5–20	3–9	20–35	2–5	130–140	7–90	11–45	--	160–320
<b>Site</b>															
AC-R	WI0018	156	0.070 <i>U</i>	0.21	81.0	1.99	0.8	0.65	10.9	0.051	0.21	0.2	0.015	0.55 <i>U</i>	185
ARC-R	SE0055	307	0.040 <i>U</i>	1.13	250	0.638	3.1	0.92	14.3	0.06	0.309	0.3	0.027	0.7	87.4
ARC-R	WI0028	142	0.0775 <i>U</i>	0.24	308	3.9	1.8	1.04	11.8	0.05	0.411	0.3	0.022	0.7	198
OR-R	SE0051	1,900	0.0515 <i>U</i>	1.87	208	0.492	16.2	2.96	8.27	0.042	0.238	0.3	0.033	4.8	59.4
OR-R	WI0026-D	134	0.0615 <i>U</i>	0.2	65.2	0.528	1.0	1.3	4.86	0.050	0.36	0.1	0.006	0.3	61.7
<b>Reference</b>															
ST-REF-3	SE0043	261	0.05 <i>U</i>	2.93	50.6	0.18	3.7 <i>J</i>	2.17	1.28	0.032	0.231	0.5	0.023	0.65 <i>U</i>	47.7 <i>J</i>
ST-REF-3	WI0025	11.0	0.035 <i>U</i>	0.04	26.4	0.356	0.2 <i>UJ</i>	2.40	0.14	0.068	0.112	0.1	0.003	0.2 <i>U</i>	97.6 <i>J</i>
ST-REF-5	SE0035	290	0.060 <i>U</i>	0.32	73.3	0.132	2.6	1.13	0.47	0.034	0.378	0.05 <i>U</i>	0.05	0.65 <i>U</i>	29.6
ST-REF-5	WI0021	15.5	0.04 <i>U</i>	0.03 <i>U</i>	57.2	0.401	0.3 <i>U</i>	2.43	0.62	0.037	0.435	0.05 <i>U</i>	0.003	0.3 <i>U</i>	79.2
ST-REF-6	SE0039	396	0.05 <i>U</i>	1.08	64.0	0.057	4.1	1.62	0.74	0.025	0.147	0.2	0.009	0.85 <i>U</i>	30
ST-REF-6	WI0022	2.5 <i>U</i>	0.04 <i>U</i>	0.03 <i>U</i>	24.1	0.558	0.2 <i>UJ</i>	2.06	0.09	0.065	0.09	0.05 <i>U</i>	0.002	0.2 <i>U</i>	92.2

**Note:** Phytotoxicity reference values and tissue concentrations are in mg/kg dry weight.

"WI" samples are willow leaves (*Salix planifolia*); "SE" samples are whole sedge plants (*Carex aquatilis*).

Sedge shoots were unwashed, and roots were rinsed in site water.

Boxed concentrations exceed the minimum phytotoxicity reference value.

-- - not reported

CoPC - chemical of potential concern

*U* - undetected; value reported is one-half the detection limit

*J* - estimated value

<sup>a</sup> Leaf tissue concentration at which toxicity was observed in plants that are neither sensitive nor tolerant (McBride 1994).

<sup>b</sup> Phytotoxic concentrations in plant foliage (Langmuir et al. 2004).

<sup>c</sup> Upper critical level in leaves and shoots of spring barley (*Hordeum vulgare*) associated with reduced yield (Davis et al. 1978).

Table 6-23. Comparison of CoPC concentrations in tundra pond sedge samples against phytotoxicity thresholds reported in the literature

Station	Sample ID	Aluminum	Antimony	Arsenic	Barium	Cadmium	Chromium	Cobalt	Lead	Mercury	Molybdenum	Selenium	Thallium	Vanadium	Zinc
<b>Phytotoxicity Reference Value</b>															
a		50–200	150	5–20	500	5–30	5–30	15–50	30–300	1–3	10–50	5–30	20	5–10	100–400
b		--	--	3–10	--	5–700	20	25–100	--	--	100	100	--	10	500–1,500
c		--	--	11–26	400–800	14–16	5–20	3–9	20–35	2–5	130–140	7–90	11–45	--	160–320
<b>Site</b>															
TP1-0100	SE0009	68.1	0.740	0.43	21.4	1.71	6.8	2.23	48.1	0.06	0.261	0.3	0.085	0.2 U	351
TP1-1000	SE0008	69.8	0.37	0.31	48.2	0.74	3.1	22.5	16.1	0.05	0.108	0.1 U	0.02	0.2 U	87.6
TP3	SE0018-D	225 J	0.17 J	0.28	70.0	0.219	2.3	1.1	3.49	0.07	1.31	0.1 U	0.017	1.0 J	74.8
TP4	SE0011	190	0.510	0.88	289	0.559	18.4	1.38	21.1	0.05	0.321	0.1 U	0.283	0.5	104
<b>Reference</b>															
TP-REF-2	SE0037	48.2	0.030 U	0.50	60.8	0.026	1.5	1.63	0.4	0.041	0.211	0.05 U	0.01	0.3 U	25.4
TP-REF-3	SE0029	1,300	0.095 U	1.96	74.9	0.081	55.0	1.72	2.3	0.03	0.9	0.1	0.13	7.6	36.6
TP-REF-5	SE0033	714	0.08 U	9.4	117	0.179	6.2	4.56	1.1	0.03	0.38	0.2	0.049	3.9	32.0

**Note:** Phytotoxicity reference values and tissue concentrations are in mg/kg dry weight.

Samples are whole sedge plants (*Carex aquatilis*).

Plant shoots were unwashed, and roots were rinsed in site water.

Boxed concentrations exceed the minimum phytotoxicity reference value.

-- - not reported

CoPC - chemical of potential concern

U - undetected; value reported is one-half the detection limit

J - estimated value

<sup>a</sup> Leaf tissue concentration at which toxicity was observed in plants that are neither sensitive nor tolerant (McBride 1994).

<sup>b</sup> Phytotoxic concentrations in plant foliage (Langmuir et al. 2004).

<sup>c</sup> Upper critical level in leaves and shoots of spring barley (*Hordeum vulgare*) associated with reduced yield (Davis et al. 1978).

**Table 6-24. Metals concentrations in site and reference stream sediments and invertebrates**

Analyte	Stream Sediment					Stream Invertebrates				
	Site			Reference		Site <sup>a</sup>			Reference <sup>b</sup>	
	AC-R	ARC-R	OR-R	ST-REF-3	ST-REF-6	AC-R	ARC-R	OR-R	ST-REF-3	ST-REF-6
Cadmium	0.49	1.06	0.44 <i>J</i>	0.25	0.19	0.228	0.803	0.365	0.696	0.347
Lead	29.2	117	22	9.5	5.71	4.43 <i>J</i>	10.9 <i>J</i>	5.16 <i>J</i>	8.14 <i>J</i>	2.73 <i>J</i>
Zinc	125	148	107	66.9	33.1	87.8 <i>J</i>	96.2 <i>J</i>	79 <i>J</i>	137 <i>J</i>	91.3 <i>J</i>

**Note:** Concentrations in mg/kg dry weight  
 Field replicates averaged  
*J* - estimated value

<sup>a</sup> Predominantly crane fly larvae, with small proportions of stone fly larvae, caddis fly larvae, and/or amphipods.

<sup>b</sup> Composite of crane fly larvae, caddis fly larvae, and stone fly larvae.

Table 6-25. Summary of water quality parameters for surface waters

Station	Date	Time	Dissolved Oxygen (mg/L)	Temperature (°C)	pH	Conductivity ( $\mu$ S/cm)	Salinity (ppt)
<b>Site Stream Surface Water</b>							
AC	06/27/04	15:10	9.21	11.4	7.6	122.9	0.1
ARC	06/27/04	11:55	10.06	8.5	7.0	--	--
OR	06/27/04	13:50	10.22	10.5	7.4	99.2	0.1
<b>Reference Stream Surface Water</b>							
ST-REF-3	07/01/04	13:09	5.22	15.4	8.1	158.2	0.1
ST-REF-5	07/01/04	14:30	2.19	17.0	7.6	38.3	0.0
ST-REF-6	07/01/04	16:08	4.08	14.5	7.2	60.6	0.0
<b>Site Tundra Pond Surface Water</b>							
TP1_100m	06/17/04	12:15	--	9.1		--	--
TP1_100m	07/01/04	14:20	2.51	23.53	7.13	715	--
TP1_1000m	06/17/04	10:45	--	11.5		--	--
TP1_1000m	07/01/04	13:25	3.36	23.25	6.2	42	--
TP3	07/01/04	--	3.03	21.55	6.38	329	--
TP4	06/17/04	17:03	--	11.2		--	--
TP4	07/01/04	18:16	2.96	21.93	6.8	119	--
<b>Reference Tundra Pond Surface Water</b>							
TP-REF-2	07/01/04	12:40	5.44	19.5	7.9	111.6	0.1
TP-REF-3	07/01/04	11:50	5.4	22.1	7.7	23.4	0.0
TP-REF-5	07/01/04	12:45	3.5	14.5	7.2	63.2	0.0
<b>Site Lagoon Surface Water</b>							
PLNL	06/29/04	10:30	3.88	17.9	7.2	2,724	1.7
NLK	06/30/04	16:09	8.19	23.6	8.5	958	0.5
NLF	07/02/04	14:35	4.07	20.9	8.6	866	0.5
<b>Reference Lagoon Surface Water</b>							
CL-REF-1	07/03/04	12:05	2.81	18.4	8.5	210.3	0.0
CL-REF-2	07/04/04	12:00	8.45	15.7	8.0	357.8	0.2
CL-REF-3	07/04/04	15:31	8.56	15.5	8.3	361.5	0.2

Table 6-26. Food-web exposure model parameters

Representative Receptor	Community	Body Weight (kg)	Food Ingestion Rate (kg/day(dry wt))	Soil/Sediment Ingestion Rate (kg/day dry wt)	Water Ingestion Rate (L/day) <sup>a</sup>	Diet Composition (percent)	Time Use (days)	Home Range (ha)
<b>Terrestrial</b>								
Willow ptarmigan	Terrestrial avian herbivores	0.53 <sup>b</sup>	0.060 <sup>c</sup>	0.0056 <sup>d</sup>	0.038	90% shrubs, 10% herbaceous plants <sup>e</sup>	365 <sup>f</sup>	3.93 <sup>g</sup>
Tundra vole	Terrestrial mammalian herbivores	0.047 <sup>h</sup>	0.0085 <sup>i</sup>	0.00020 <sup>j</sup>	0.0063	90% herbaceous plants, 5% moss, 5% lichen <sup>k</sup>	365 <sup>f</sup>	0.1087 <sup>l</sup>
Caribou	Terrestrial mammalian herbivores	107 <sup>m</sup>	5.0 <sup>n</sup>	0.34 <sup>o</sup>	6.6	70% lichen, 10% shrubs, 10% herbaceous plants, 10% moss <sup>p</sup>	150 <sup>q</sup>	NA
Moose	Terrestrial mammalian herbivores	339 <sup>r</sup>	6.4 <sup>s</sup>	0.13 <sup>t</sup>	19	90% shrubs, 10% herbaceous plants <sup>u</sup>	365 <sup>f</sup>	2,849–29,008 <sup>v</sup>
Lapland longspur	Terrestrial avian invertevores	0.0254 <sup>w</sup>	0.0053 <sup>x</sup>	0.000074 <sup>y</sup>	0.0050	90% invertebrates, 10% herbaceous plants <sup>z</sup>	150 <sup>aa</sup>	1.76 <sup>bb</sup>
Tundra shrew	Terrestrial mammalian invertevores	0.0064 <sup>cc</sup>	0.0021 <sup>dd</sup>	0.00011 <sup>ee</sup>	0.0011	100% invertebrates <sup>ff</sup>	365 <sup>f</sup>	0.22 <sup>gg</sup>
Snowy owl	Terrestrial avian carnivores	2.28 <sup>hh</sup>	0.10 <sup>ii</sup>	0.0020 <sup>tt</sup>	0.10	100% small mammals <sup>jj</sup>	365 <sup>f</sup>	777 <sup>kk</sup>
Arctic fox	Terrestrial mammalian carnivores	3.2 <sup>ll</sup>	0.11 <sup>mm</sup>	0.0031 <sup>nn</sup>	0.28	100% small mammals <sup>oo</sup>	365 <sup>f</sup>	407 <sup>pp</sup>
<b>Freshwater Aquatic</b>								
Green-winged teal	Freshwater aquatic avian herbivores	0.32 <sup>qq</sup>	0.053 <sup>rr</sup>	0.0010 <sup>ss</sup>	0.027	100% herbaceous plants <sup>tt</sup>	123 <sup>uu</sup>	243 <sup>vv</sup>
Muskrat	Freshwater aquatic mammalian herbivores	0.932 <sup>ww</sup>	0.070 <sup>xx</sup>	0.0014 <sup>t</sup>	0.093	100% herbaceous plants <sup>yy</sup>	365 <sup>f</sup>	0.17 <sup>zz</sup>
Common snipe	Freshwater aquatic avian invertevores	0.116 <sup>qq</sup>	0.015 <sup>aaa</sup>	0.0016 <sup>bbb</sup>	0.014	90% invertebrates, 10% herbaceous plants <sup>ccc</sup>	109 <sup>ddd</sup>	0.0908–47.7 <sup>eee</sup>
<b>Coastal Lagoon</b>								
Brant	Marine avian herbivores	1.23 <sup>qq</sup>	0.13 <sup>rr</sup>	0.011 <sup>fff</sup>	0.068	95% herbaceous plants, 5% moss <sup>ggg</sup>	126 <sup>hhh</sup>	201.06 <sup>iii</sup>
Black-bellied plover	Marine avian invertevores	0.214 <sup>jjj</sup>	0.028 <sup>kkk</sup>	0.0082 <sup>lll</sup>	0.021	100% invertebrates <sup>mmm</sup>	124 <sup>nnn</sup>	53 <sup>ooo</sup>

<sup>a</sup> Based on U.S. EPA (1993) drinking water ingestion equations for all birds or all mammals.

<sup>b</sup> Mean female body weight from West et al. (1970).

<sup>c</sup> Estimated from Andreev (1991).

<sup>d</sup> Based on 9.3 percent soil in wild turkey diet from Beyer et al. (1994).

<sup>e</sup> Estimated from diets reported for Alaska in Hannon et al. (1998).

<sup>f</sup> Assumes receptor is present year-round at the site.

<sup>g</sup> Mean territory size for monogamous males (Hannon and Dobush 1997).

<sup>h</sup> Mean female body weight from Bee and Hall (1956).

<sup>i</sup> Based on Nagy et al. (1999) allometric equation for Rodentia.

<sup>j</sup> Based on 2.4 percent soil in meadow vole diet from Beyer et al. (1994).

<sup>k</sup> Estimated from summer and winter diets at Pearce Point, NWT (Bergman and Krebs 1993).

<sup>l</sup> Mean home range for reproductive females at Pearce Point, NWT (Lambin et al. 1992).

Table 6-26. (cont.)

- 
- <sup>m</sup> Mean female in Alaska from Silva and Downing (1995).
- <sup>n</sup> Based on mean value from Hanson et al. (1975).
- <sup>o</sup> Based on 6.8 percent soil in bison diet from Beyer et al. (1994).
- <sup>p</sup> Based on diets reported in Miller (1976), Boertje (1990), and Scotter (1967).
- <sup>q</sup> Best professional judgment based on Lent (1966), Hemming (1987, 1988, 1989, 1991), and Pollard (1994a,b).
- <sup>r</sup> Mean body weight for female Alaskan moose measured at the Kenai Moose Research Center, Soldotna, AK (Franzmann et al. 1978).
- <sup>s</sup> Average daily ingestion rate for all female moose; 1.9% of body weight per day on a dry weight basis (Schwartz et al. 1984).
- <sup>t</sup> Based on minimum soil ingestion rate from Beyer et al. (1994).
- <sup>u</sup> Estimated from diets reported for Alaska in Franzmann and Schwartz (1997).
- <sup>v</sup> Mean home ranges of nonmigratory individuals in Alaska (Franzmann and Schwartz 1997).
- <sup>w</sup> Mean female body weight from Irving (1960).
- <sup>x</sup> Calculated using an average female daily energy budget of 118 kJ/day and average prey caloric value of 22.16 kJ/g from Custer et al. (1986).
- <sup>y</sup> Based on 1.4 percent soil in Lapland longspur diet reported by URS Team (1996).
- <sup>z</sup> Estimated from summer diets near Barrow, AK (Custer and Pitelka 1978).
- <sup>aa</sup> Based on 150 days from first to last sighting in Cape Thompson area reported by Williamson et al. (1966).
- <sup>bb</sup> Mean male breeding territory near Barrow, AK (Seastedt and MacLean 1979).
- <sup>cc</sup> Mean body weight from Bee and Hall (1956) and Martell and Pearson (1978).
- <sup>dd</sup> Based on measured food consumption from Buckner (1964), assuming a mid-range moisture content of 75 percent in invertebrates from U.S. EPA (1993).
- <sup>ee</sup> Best professional judgment based on Beyer et al. (1994).
- <sup>ff</sup> Based on Yudin (1962, as cited in Aitchison 1987 and Buckner 1964).
- <sup>gg</sup> Mean home range for breeding females (*Sorex vagrans* and *Sorex obscurus*) in British Columbia, Canada (Hawes 1977).
- <sup>hh</sup> Mean female body weight from Kerlinger and Lein (1988).
- <sup>ii</sup> Estimated from Gessaman (1972) and Pitelka et al. (1955), assuming a moisture content of 68 percent in diet from U.S. EPA (1993).
- <sup>jj</sup> Simplified from Parmelee (1992).
- <sup>kk</sup> Mean nesting territory near Barrow, AK (Pitelka et al. 1955).
- <sup>ll</sup> Mean female body weight from Anthony (1997).
- <sup>mmm</sup> Based on Nagy et al. (1999) allometric equation for Carnivora.
- <sup>nn</sup> Based on 2.8 percent soil in red fox diet from Beyer et al. (1994).
- <sup>oo</sup> Simplified from Anthony et al. (2000).
- <sup>pp</sup> Mean female home range in western Alaska (Anthony 1997).
- <sup>qq</sup> Mean female body weight from Dunning (1993).
- <sup>rr</sup> Based on Nagy et al. (1999) allometric equation for all birds.

Table 6-26. (cont.)

---



---

<sup>ss</sup>	Based on 1.9 percent sediment in green-winged teal diet from Beyer et al. (1999).
<sup>tt</sup>	Estimated from autumn diet in southeastern Alaska (Hughes and Young 1982).
<sup>uu</sup>	Based on 123 days from first to last sighting in Cape Thompson area reported by Williamson et al. (1966).
<sup>vv</sup>	Home range for one pair in South Dakota (Drewien 1967, as cited in Granholm 2003).
<sup>ww</sup>	Mean body weight from Fuller (1951).
<sup>xx</sup>	Estimated from Campbell et al. (1998).
<sup>yy</sup>	Based on diets reported in U.S. EPA (1993).
<sup>zz</sup>	Mean female home range in Iowa (Neal 1968, as cited in U.S. EPA 1993).
<sup>aaa</sup>	Based on Nagy et al. (1999) allometric equation for Insectivores.
<sup>bbb</sup>	Based on 10.4 percent soil in American woodcock diet from Beyer et al. (1994).
<sup>ccc</sup>	Based on diets reported in Mueller (1999).
<sup>ddd</sup>	Based on 109 days from first to last sighting in Cape Thompson area reported by Williamson et al. (1966).
<sup>eee</sup>	Estimated area based on a 17–390 m mean distance (radius) females traveled from nest to feeding sites during incubation period (Green et al. 1990).
<sup>fff</sup>	Based on 8.2 percent soil in Canada goose diet from Beyer et al. (1994).
<sup>ggg</sup>	Based on breeding season diets reported in Reed et al. (1998).
<sup>hhh</sup>	Based on 126 days from first to last sighting in Cape Thompson area reported by Williamson et al. (1966).
<sup>iii</sup>	Estimated assuming a maximum foraging distance (radius) of 800 m from nest (Reed et al. 1998).
<sup>jjj</sup>	Mean female body weight for Alaska from Paulson (1995).
<sup>kkk</sup>	Based on Nagy et al. (1999) allometric equation for Charadriiformes.
<sup>lll</sup>	Based on 29% sediment in black-bellied plover diet from Hui and Beyer (1998).
<sup>mmm</sup>	Based on breeding season diets reported in Paulson (1995).
<sup>nnn</sup>	Based on 124 days from first to last sighting of American golden plover in Cape Thompson area reported by Williamson et al. (1966).
<sup>ooo</sup>	Estimated based on average radius of breeding territory in northern Alaska (412 m) (Moitoret pers. comm., as cited in Paulson 1995).

Table 6-27. Toxicity reference values for risk evaluation for wildlife receptors

CoPC	TRVs (mg/kg-day)					
	Avian			Mammalian		
	NOAEL	LOAEL	Citation	NOAEL	LOAEL	Citation
Aluminum	120	NA	Carriere et al. (1986)	1.9	19	Ondreicka et al. (1966)
Antimony	NA	NA	NA	0.66	NA	Schroeder et al. (1970)
Arsenic (arsenate)	10	40	Stanley et al. (1994)	0.40	1.6	Nemec et al. (1998)
Arsenic (arsenite)	20	50	USFWS (1964)	0.13	1.3	Schroeder and Mitchener (1971)
Barium	21	42	Johnson et al. (1960)	5.1	--	Perry et al. (1983)
	--	--	--	--	20	Borzelleca et al. (1988)
Cadmium	1.5	20	White and Finley (1978)	1.0	10	Sutou et al. (1980)
Chromium	0.86	4.3	Haseltine et al. (1985), as cited in Sample et al. (1996)	3.3	--	Mackenzie et al. (1958)
	--	--	--	--	69	Gross and Heller (1946)
Cobalt	NA	NA	NA	0.5	2.0	Nation et al. (1983)
Lead	3.9	--	Pattee (1984)	11	90	Azar et al. (1973)
	--	11	Edens et al. (1976)	--	--	--
Mercury <sup>a</sup>	0.032	0.064	Heinz (1974, 1976a,b, 1979)	0.032	0.16	Verschuuren et al. (1976)
Molybdenum	3.5	35	Lepore and Miller (1965)	0.26	2.6	Schroeder and Mitchener (1971)
Selenium	0.40	0.80	Heinz et al. (1989)	0.20	0.33	Rosenfeld and Beath (1954)
Thallium	0.24	24	Hudson et al. (1984)	0.074	0.74	Formigli et al. (1986)
Vanadium	11	NA	White and Dieter (1978)	0.21	2.1	Domingo et al. (1986)
Zinc (TRV1)	130	NA	Stahl et al. (1990)	160	320	Schlicker and Cox (1968)
Zinc (TRV2)	70	120	Jackson et al. (1986)	--	--	--

**Note:** -- - not applicable  
 CoPC - chemical of potential concern  
 LOAEL - lowest-observed-adverse-effect level  
 NA - not available; no suitable TRV was derived  
 NOAEL - no-observed-adverse-effect level  
 TRV - toxicity reference value

<sup>a</sup> Mercury TRVs were based on exposure to methylmercury.



Table 6-28. Food-web exposure modeling results for caribou

Assessment Unit	Chemical	NOAEL Hazard Quotient		LOAEL Hazard Quotient	
		Mean	95% UCL	Mean	95% UCL
Reference area	Aluminum	8.9	16	0.89	1.6
Port	Aluminum	22	44	2.2	4.4
Port	Barium	1.6	2.6	0.41	0.66
Road	Aluminum	23	43	2.3	4.3
Road	Barium	2.4	4.1	0.62	1.0
Mine	Aluminum	24	67	2.4	6.7
Mine	Barium	5.1	7.9	1.3	2.0
Mine	Lead	0.52	1.1	0.064	0.13
Whole site	Aluminum	25	40	2.5	4.0
Whole site	Barium	3.1	4.7	0.80	1.2

**Note:** Results shown only for chemicals with NOAEL-based hazard quotients >1.0.

For 11 CoPCs (antimony, arsenic, cadmium, chromium, cobalt, mercury, molybdenum, selenium, thallium, vanadium, and zinc) all hazard quotients were less than 1.0.

95% UCL - 95 percent upper confidence limit on the mean

LOAEL - lowest-observed-adverse-effect level

NOAEL - no-observed-adverse-effect level

**Table 6-29. Food-web exposure modeling results for Arctic fox**

Assessment Unit	Chemical	NOAEL Hazard Quotient		LOAEL Hazard Quotient	
		Mean	95% UCL	Mean	95% UCL
Reference area	Aluminum	2.3	4.4	0.23	0.44
Port	Aluminum	11	21	1.1	2.1
Road	Aluminum	2.8	5.0	0.28	0.50
Road	Mercury	2.6	11	0.51	2.2

**Note:** Results shown only for chemicals with NOAEL-based hazard quotients >1.0.

For 12 CoPCs (antimony, arsenic, barium, cadmium, chromium, cobalt, lead, molybdenum, selenium, thallium, vanadium, and zinc) all hazard quotients were less than 1.0.

95% UCL - 95 percent upper confidence limit on the mean

LOAEL - lowest-observed-adverse-effect level

NOAEL - no-observed-adverse-effect level