

# Human Health Conceptual Site Model Scoping Form

Site Name: \_\_\_\_\_

File Number: \_\_\_\_\_

Completed by: \_\_\_\_\_

## Introduction

The form should be used to reach agreement with the Alaska Department of Environmental Conservation (DEC) about which exposure pathways should be further investigated during site characterization. From this information, a CSM graphic and text must be submitted with the site characterization work plan.

*General Instructions: Follow the italicized instructions in each section below.*

## 1. General Information:

**Sources** (*check potential sources at the site*)

- |  |                                       |
|--|---------------------------------------|
| <input type="checkbox"/> USTs                          | <input type="checkbox"/> Vehicles     |
| <input type="checkbox"/> ASTs                          | <input type="checkbox"/> Landfills    |
| <input type="checkbox"/> Dispensers/fuel loading racks | <input type="checkbox"/> Transformers |
| <input type="checkbox"/> Drums                         | <input type="checkbox"/> Other: _____ |

**Release Mechanisms** (*check potential release mechanisms at the site*)

- |                                 |   |
|---------------------------------|---|
| <input type="checkbox"/> Spills | <input type="checkbox"/> Direct discharge |
| <input type="checkbox"/> Leaks  | <input type="checkbox"/> Burning          |
|                                 | <input type="checkbox"/> Other: _____     |

**Impacted Media** (*check potentially-impacted media at the site*)

- |  |  |
|--|--|
| <input type="checkbox"/> Surface soil (0-2 feet bgs*)  | <input type="checkbox"/> Groundwater   |
| <input type="checkbox"/> Subsurface Soil (>2 feet bgs) | <input type="checkbox"/> Surface water |
| <input type="checkbox"/> Air                           | <input type="checkbox"/> Other: _____  |

**Receptors** (*check receptors that could be affected by contamination at the site*)

- |   |  |
|---|--|
| <input type="checkbox"/> Residents (adult or child)                       | <input type="checkbox"/> Site visitor      |
| <input type="checkbox"/> Commercial or industrial worker                  | <input type="checkbox"/> Trespasser        |
| <input type="checkbox"/> Construction worker                              | <input type="checkbox"/> Recreational user |
| <input type="checkbox"/> Subsistence harvester (i.e., gathers wild foods) | <input type="checkbox"/> Farmer            |
| <input type="checkbox"/> Subsistence consumer (i.e., eats wild foods)     | <input type="checkbox"/> Other: _____      |

\* bgs – below ground surface

**2. Exposure Pathways:** (The answers to the following questions will identify complete exposure pathways at the site. Check each box where the answer to the question is "yes".)

**a) Direct Contact –**

**1 Incidental Soil Ingestion**

Is soil contaminated anywhere between 0 and 15 feet bgs?

Do people use the site or is there a chance they will use the site in the future?

If both boxes are checked, label this pathway complete: \_\_\_\_\_

**2 Dermal Absorption of Contaminants from Soil**

Is soil contaminated anywhere between 0 and 15 feet bgs?

Do people use the site or is there a chance they will use the site in the future?

Can the soil contaminants permeate the skin? (Contaminants listed below, or within the groups listed below, should be evaluated for dermal absorption).

- |                                |                   |
|--------------------------------|-------------------|
| Arsenic                        | Lindane           |
| Cadmium                        | PAHs              |
| Chlordane                      | Pentachlorophenol |
| 2,4-dichlorophenoxyacetic acid | PCBs              |
| Dioxins                        | SVOCs             |
| DDT                            |                   |

If all of the boxes are checked, label this pathway complete: \_\_\_\_\_

**b) Ingestion –**

**1 Ingestion of Groundwater**

Have contaminants been detected or are they expected to be detected in the groundwater, OR are contaminants expected to migrate to groundwater in the future?

Could the potentially affected groundwater be used as a current or future drinking water source? Please note, only leave the box unchecked if ADEC has determined the groundwater is not a currently or reasonably expected future source of drinking water according to 18 AAC 75.350.

If both the boxes are checked, label this pathway complete: \_\_\_\_\_

## 2 Ingestion of Surface Water

Have contaminants been detected or are they expected to be detected in surface water OR are contaminants expected to migrate to surface water in the future?

Could potentially affected surface water bodies be used, currently or in the future, as a drinking water source? *Consider both public water systems and private use (i.e., during residential, recreational or subsistence activities).*

*If both boxes are checked, label this pathway complete:* \_\_\_\_\_

## 3 Ingestion of Wild Foods

Is the site in an area that is used or reasonably could be used for hunting, fishing, or harvesting of wild food?

Do the site contaminants have the potential to bioaccumulate (*see Appendix A*)?

Are site contaminants located where they would have the potential to be taken up into biota? (i.e. the top 6 feet of soil, in groundwater that **could be** connected to surface water, etc.)

*If all of the boxes are checked, label this pathway complete:* \_\_\_\_\_

### c) Inhalation

#### 1 Inhalation of Outdoor Air

Is soil contaminated anywhere between 0 and 15 feet bgs?

Do people use the site or is there a chance they will use the site in the future?

Are the contaminants in soil volatile (*See Appendix B*)?

*If all of the boxes are checked, label this pathway complete:* \_\_\_\_\_

#### 2 Inhalation of Indoor Air

Are occupied buildings on the site or reasonably expected to be placed on the site in an area that could be affected by contaminant vapors? (i.e., within 100 feet, horizontally or vertically, of the contaminated soil or groundwater, or subject to “preferential pathways” that promote easy airflow, like utility conduits or rock fractures)

Are volatile compounds present in soil or groundwater (*See Appendix C*)?

*If both boxes are checked, label this pathway complete:* \_\_\_\_\_

**3. Additional Exposure Pathways:** *(Although there are no definitive questions provided in this section, these exposure pathways should also be considered at each site. Use the guidelines provided below to determine if further evaluation of each pathway is warranted.)*

**Dermal Exposure to Contaminants in Groundwater and Surface Water**

Exposure from this pathway may need to be assessed only in cases where DEC water-quality or drinking-water standards are not being applied as cleanup levels. Examples of conditions that may warrant further investigation include:

- Climate permits recreational use of waters for swimming,
- Climate permits exposure to groundwater during activities, such as construction, without protective clothing, or
- Groundwater or surface water is used for household purposes.

*Check the box if further evaluation of this pathway is needed:*

Comments:

**Inhalation of Volatile Compounds in Household Water**

Exposure from this pathway may need to be assessed only in cases where DEC water-quality or drinking-water standards are not being applied as cleanup levels. Examples of conditions that may warrant further investigation include:

- The contaminated water is used for household purposes such as showering, laundering, and dish washing, and
- The contaminants of concern are volatile (common volatile contaminants are listed in Appendix B)

*Check the box if further evaluation of this pathway is needed:*

Comments:

**Inhalation of Fugitive Dust**

Generally DEC soil ingestion cleanup levels in Table B1 of 18 AAC 75 are protective of this pathway, although this is not true in the case of chromium. Examples of conditions that may warrant further investigation include:

- Nonvolatile compounds are found in the top 2 centimeters of soil. The top 2 centimeters of soil are likely to be dispersed in the wind as dust particles.
- Dust particles are less than 10 micrometers. This size can be inhaled and would be of concern for determining if this pathway is complete.

*Check the box if further evaluation of this pathway is needed:*

Comments:

### **Direct Contact with Sediment**

This pathway involves people's hands being exposed to sediment, such as during recreational or some types of subsistence activities. People then incidentally **ingest** sediment from normal hand-to-mouth activities. In addition, **dermal absorption of contaminants** may be of concern if people come in contact with sediment and the contaminants are able to permeate the skin (see dermal exposure to soil section). This type of exposure is rare but it should be investigated if:

- Climate permits recreational activities around sediment, and/or
- Community has identified subsistence or recreational activities that would result in exposure to the sediment, such as clam digging.

ADEC soil ingestion cleanup levels are protective of direct contact with sediment. If they are determined to be over-protective for sediment exposure at a particular site, other screening levels could be adopted or developed.

*Check the box if further evaluation of this pathway is needed:*

Comments:

**4. Other Comments** *(Provide other comments as necessary to support the information provided in this form.)*

## APPENDIX A

### BIOACCUMULATIVE COMPOUNDS

**Table A-1: List of Compounds of Potential Concern for Bioaccumulation**

Organic compounds are identified as bioaccumulative if they have a BCF equal to or greater than 1,000 or a log  $K_{ow}$  greater than 3.5. Inorganic compounds are identified as bioaccumulative if they are listed as such by EPA (2000). Those compounds in Table X of 18 AAC 75.345 that are bioaccumulative, based on the definition above, are listed below.

Aldrin	DDT	Lead
Arsenic	Dibenzo(a,h)anthracene	Mercury
Benzo(a)anthracene	Dieldrin	Methoxychlor
Benzo(a)pyrene	Dioxin	Nickel
Benzo(b)fluoranthene	Endrin	PCBs
Benzo(k)fluoranthene	Fluoranthene	
Cadmium	Heptachlor	Pyrene
Chlordane	Heptachlor epoxide	Selenium
Chrysene	Hexachlorobenzene	Silver
Copper	Hexachlorocyclopentadiene	Toxaphene
DDD	Indeno(1,2,3-c,d)pyrene	Zinc
DDE		

Because BCF values can relatively easily be measured or estimated, the BCF is frequently used to determine the potential for a chemical to bioaccumulate. A compound with a BCF greater than 1,000 is considered to bioaccumulate in tissue (EPA 2004b).

For inorganic compounds, the BCF approach has not been shown to be effective in estimating the compound's ability to bioaccumulate. Information available, either through scientific literature or site-specific data, regarding the bioaccumulative potential of an inorganic site contaminant should be used to determine if the pathway is complete.

The list was developed by including organic compounds that either have a BCF equal to or greater than 1,000 or a log  $K_{ow}$  greater than 3.5 and inorganic compounds that are listed by the United States Environmental Protection Agency (EPA) as being bioaccumulative (EPA 2000). The BCF can also be estimated from a chemical's physical and chemical properties. A chemical's octanol-water partitioning coefficient ( $K_{ow}$ ) along with defined regression equations can be used to estimate the BCF. EPA's Persistent, Bioaccumulative, and Toxic (PBT) Profiler (EPA 2004) can be used to estimate the BCF using the  $K_{ow}$  and linear regressions presented by Meylan et al. (1996). The PBT Profiler is located at <http://www.pbtprofiler.net/>. For compounds not found in the PBT Profiler, DEC recommends using a log  $K_{ow}$  greater than 3.5 to determine if a compound is bioaccumulative.

## APPENDIX B

### VOLATILE COMPOUNDS

**Table B-1: List of Volatile Compounds of Potential Concern**

Common volatile contaminants of concern at contaminated sites. A chemical is defined as volatile if the Henry's Law constant is  $1 \times 10^{-5}$  atm-m<sup>3</sup>/mol or greater and the molecular weight less than 200 g/mole (g/mole; EPA 2004a). Those compounds in Table X of 18 AAC 75.345 that are volatile, based on the definition above, are listed below.

Acenaphthene	1,4-dichlorobenzene	Pyrene
Acetone	1,1-dichloroethane	Styrene
Anthracene	1,2-dichloroethane	1,1,2,2-tetrachloroethane
Benzene	1,1-dichloroethylene	Tetrachloroethylene
Bis(2-chlorethyl)ether	Cis-1,2-dichloroethylene	Toluene
Bromodichloromethane	Trans-1,2-dichloroethylene	1,2,4-trichlorobenzene
Carbon disulfide	1,2-dichloropropane	1,1,1-trichloroethane
Carbon tetrachloride	1,3-dichloropropane	1,1,2-trichloroethane
Chlorobenzene	Ethylbenzene	Trichloroethylene
Chlorodibromomethane	Fluorene	Vinyl acetate
Chloroform	Methyl bromide	Vinyl chloride
2-chlorophenol	Methylene chloride	Xylenes
Cyanide	Naphthalene	GRO
1,2-dichlorobenzene	Nitrobenzene	DRO

## APPENDIX C

### COMPOUNDS OF CONCERN FOR VAPOR MIGRATION

**Table C-1: List of Compounds of Potential Concern for the Vapor Migration**

A chemical is considered sufficiently toxic if the vapor concentration of the pure component poses an incremental lifetime cancer risk greater than  $10^{-6}$  or a non-cancer hazard index greater than 1. A chemical is considered sufficiently volatile if its Henry's Law constant is  $1 \times 10^{-5}$  atm-m<sup>3</sup>/mol or greater.

Acenaphthene	Dibenzofuran	Hexachlorobenzene
Acetaldehyde	1,2-Dibromo-3-chloropropane	Hexachlorocyclopentadiene
Acetone	1,2-Dibromoethane (EDB)	Hexachloroethane
Acetonitrile	1,3-Dichlorobenzene	Hexane
Acetophenone	1,2-Dichlorobenzene	Hydrogen cyanide
Acrolein	1,4-Dichlorobenzene	Isobutanol
Acrylonitrile	2-Nitropropane	Mercury (elemental)
Aldrin	N-Nitroso-di-n-butylamine	Methacrylonitrile
alpha-HCH (alpha-BHC)	n-Propylbenzene	Methoxychlor
Benzaldehyde	o-Nitrotoluene	Methyl acetate
Benzene	o-Xylene	Methyl acrylate
Benzo(b)fluoranthene	p-Xylene	Methyl bromide
Benzylchloride	Pyrene	Methyl chloride chloromethane)
beta-Chloronaphthalene	sec-Butylbenzene	Methylcyclohexane
Biphenyl	Styrene	Methylene bromide
Bis(2-chloroethyl)ether	tert-Butylbenzene	Methylene chloride
Bis(2-chloroisopropyl)ether	1,1,1,2-Tetrachloroethane	Methylethylketone (2-butanone)
Bis(chloromethyl)ether	1,1,2,2-Tetrachloroethane	Methylisobutylketone
Bromodichloromethane	Tetrachloroethylene	Methylmethacrylate
Bromoform	Dichlorodifluoromethane	2-Methylnaphthalene
1,3-Butadiene	1,1-Dichloroethane	MTBE
Carbon disulfide	1,2-Dichloroethane	m-Xylene
Carbon tetrachloride	1,1-Dichloroethylene	Naphthalene
Chlordane	1,2-Dichloropropane	n-Butylbenzene
2-Chloro-1,3-butadiene (chloroprene)	1,3-Dichloropropene	Nitrobenzene
Chlorobenzene	Dieldrin	Toluene
1-Chlorobutane	Endosulfan	trans-1,2-Dichloroethylene
Chlorodibromomethane	Epichlorohydrin	1,1,2-Trichloro-1,2,2-trifluoroethane
Chlorodifluoromethane	Ethyl ether	1,2,4-Trichlorobenzene
Chloroethane (ethyl chloride)	Ethylacetate	1,1,2-Trichloroethane
Chloroform	Ethylbenzene	1,1,1-Trichloroethane
2-Chlorophenol	Ethylene oxide	Trichloroethylene
2-Chloropropane	Ethylmethacrylate	Trichlorofluoromethane
Chrysene	Fluorene	1,2,3-Trichloropropane
cis-1,2-Dichloroethylene	Furan	1,2,4-Trimethylbenzene
Crotonaldehyde (2-butenal)	Gamma-HCH (Lindane)	1,3,5-Trimethylbenzene
Cumene	Heptachlor	Vinyl acetate
DDE	Hexachloro-1,3-butadiene	Vinyl chloride (chloroethene)

Source: EPA 2002.

Guidance on Developing Conceptual Site Models  
January 31, 2005